

Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11) **EP 0 796 913 A2**

(12) **EUROPEAN PATENT APPLICATION**

(43) Date of publication:  
24.09.1997 Bulletin 1997/39

(21) Application number: 97104842.6

(22) Date of filing: 19.03.1997

(51) Int. Cl.<sup>6</sup>: **C12N 15/12**, C12N 15/54,  
C12N 15/55, C07K 14/47,  
C12N 9/12, C12N 9/00,  
C12N 9/64, C12Q 1/68,  
A61K 38/17, A61K 38/45,  
A61K 38/53

(84) Designated Contracting States:  
**AT BE CH DE DK ES FI FR GB GR IE IT LI LU MC  
NL PT SE**

(30) Priority: 19.03.1996 JP 63410/96  
05.03.1997 JP 69163/97

(71) Applicant:  
**OTSUKA PHARMACEUTICAL CO., LTD.**  
Chiyoda-ku Tokyo 101 (JP)

(72) Inventors:  
• **Fujiwara, Tsutomu**  
Naruto-shi, Tokushima-ken (JP)

• **Watanabe, Takeshi**  
Aizumi-cho, Itano-gun, Tokushima-ken (JP)  
• **Horie, Masato**  
Tokushima-shi, Tokushima-ken (JP)

(74) Representative: **Hansen, Bernd, Dr. Dipl.-Chem.**  
et al  
**Hoffmann Eitle,**  
Patent- und Rechtsanwälte,  
Arabellastrasse 4  
81925 München (DE)

Remarks:

The applicant has subsequently filed a sequence listing and declared, that it includes no new matter.

(54) **GDP dissociation stimulating protein, brain-specific nucleosome assembly protein, skeletal muscle specific ubiquitin-conjugating enzyme, cell proliferation protein, phosphatidylinositolkinase, nel related proteins**

(57) The present invention provides novel human genes, for example a novel human gene comprising a nucleotide sequence coding for the amino acid sequence shown under SEQ ID NO:1. The use of the genes makes it possible to detect the expression of the same in various tissues, analyze their structures and functions, and produce the human proteins encoded by the genes by the technology of genetic engineering. Through these, it becomes possible to analyze the corresponding expression products, elucidate the pathology of diseases associated with the genes, for example hereditary diseases and cancer, and diagnose and treat such diseases.

EP 0 796 913 A2

## Description

TECHNICAL FIELD

5 The present invention relates to a gene useful as an indicator in the prophylaxis, diagnosis and treatment of diseases in humans. More particularly, it relates to a novel human gene analogous to rat, mouse, yeast, nematode and known human genes, among others, and utilizable, after cDNA analysis thereof, chromosome mapping of cDNA and function analysis of cDNA, in gene diagnosis using said gene and in developing a novel therapeutic method.

10 BACKGROUND ART

The genetic information of a living thing has been accumulated as sequences (DNA) of four bases, namely A, C, G and T, which exist in cell nuclei. Said genetic information has been preserved for line preservation and ontogeny of each individual living thing.

15 In the case of human being, the number of said bases is said to be about 3 billion ( $3 \times 10^9$ ) and supposedly there are 50 to 100 thousand genes therein. Such genetic information serves to maintain biological phenomena in that regulatory proteins, structural proteins and enzymes are produced via such route that mRNA is transcribed from a gene (DNA) and then translated into a protein. Abnormalities in said route from gene to protein translation are considered to be causative of abnormalities of life supporting systems, for example in cell proliferation and differentiation, hence causative of various diseases.

20 As a result of gene analyses so far made, a number of genes which may be expected to serve as useful materials in drug development, have been found, for example genes for various receptors such as insulin receptor and LDL receptor, genes involved in cell proliferation and differentiation and genes for metabolic enzymes such as proteases, ATPase and superoxide dismutases.

25 However, analysis of human genes and studies of the functions of the genes analyzed and of the relations between the genes analyzed and various diseases have been just begun and many points remain unknown. Further analysis of novel genes, analysis of the functions thereof, studies of the relations between the genes analyzed and diseases, and studies for applying the genes analyzed to gene diagnosis or for medicinal purposes, for instance, are therefore desired in the relevant art.

30 If such a novel human gene as mentioned above can be provided, it will be possible to analyze the level of expression thereof in each cell and the structure and function thereof and, through expression product analysis and other studies, it may become possible to reveal the pathogenesis of a disease associated therewith, for example a genopathy or cancer, or diagnose and treat said disease, for instance. It is an object of the present invention to provide such a novel human gene.

35 For attaining the above object, the present inventors made intensive investigations and obtained the findings mentioned below. Based thereon, the present invention has now been completed.

DISCLOSURE OF INVENTION

40 Thus, the present inventors synthesized cDNAs based on mRNAs extracted from various tissues, inclusive of human fetal brain, adult blood vessels and placenta, constructed libraries by inserting them into vectors, allowing colonies of *Escherichia coli* transformed with said libraries to form on agar medium, picked up colonies at random and transferred to 96-well micro plates and registered a large number of human gene-containing *E. coli* clones.

45 Each clone thus registered was cultivated on a small size, DNA was extracted and purified, the four base-specifically terminating extension reactions were carried out by the dideoxy chain terminator method using the cDNA extracted as a template, and the base sequence of the gene was determined over about 400 bases from the 5' terminus thereof using an automatic DNA sequencer. Based on the thus-obtained base sequence information, a novel family gene analogous to known genes of animal and plant species such as bacteria, yeasts, nematodes, mice and humans was searched for.

50 The method of the above-mentioned cDNA analysis is detailedly described in the literature by Fujiwara, one of the present inventors [Fujiwara, Tsutomu, Saibo Kogaku (Cell Engineering), 14, 645-654 (1995)].

Among this group, there are novel receptors, DNA binding domain-containing transcription regulating factors, signal transmission system factors, metabolic enzymes and so forth. Based on the homology of the novel gene of the present invention as obtained by gene analysis to the genes analogous thereto, the product of the gene, hence the function of the protein, can approximately be estimated by analogy. Furthermore, such functions as enzyme activity and binding ability can be investigated by inserting the candidate gene into an expression vector to give a recombinant.

According to the present invention, there are provided a novel human gene characterized by containing a nucleotide sequence coding for an amino acid sequence defined by SEQ ID NO:1 :4, :7, :10, :13, :16, :19, :22, :25, :28, :31, :34, :37 or 40, a human gene characterized by containing the nucleotide sequence defined by SEQ ID NO:2, :5, :8, :11,

:14, :17, :20, :23, :26, :29, :32, :35, :38 or :41, respectively coding for the amino acid sequence mentioned above, and a novel human gene characterized by the nucleotide sequence defined by SEQ ID NO:3, :6, :9, :12, :15, :18, :21, :24, :27, :30, :33, :36, :39 or :42.

The symbols used herein for indicating amino acids, peptides, nucleotides, nucleotide sequences and so on are those recommended by IUPAC and IUB or in "Guideline for drafting specifications etc. including nucleotide sequences or amino acid sequences" (edited by the Japanese Patent Office), or those in conventional use in the relevant field of art.

As specific examples of such gene of the present invention, there may be mentioned genes deducible from the DNA sequences of the clones designated as "GEN-501D08", "GEN-080G01", "GEN-025F07", "GEN-076C09", "GEN-331G07", "GEN-163D09", "GEN-078D05TA13", "GEN-423A12", "GEN-092E10", "GEN-428B12", "GEN-073E07", "GEN-093E05" and "GEN-077A09" shown later herein in Examples 1 to 11. The respective nucleotide sequences are as shown in the sequence listing.

These clones have an open reading frame comprising nucleotides (nucleic acid) respectively coding for the amino acids shown in the sequence listing. Their molecular weights were calculated at the values shown later herein in the respective examples. Hereinafter, these human genes of the present invention are sometimes referred to as the designation used in Examples 1 to 11.

In the following, the human gene of the present invention is described in further detail.

As mentioned above, each human gene of the present invention is analogous to rat, mouse, yeast, nematode and known human genes, among others, and can be utilized in human gene analysis based on the information about the genes analogous thereto and in studying the function of the gene analyzed and the relation between the gene analyzed and a disease. It is possible to use said gene in gene diagnosis of the disease associated therewith and in exploitation studies of said gene for medicinal purposes.

The gene of the present invention is represented in terms of a single-stranded DNA sequence, as shown under SEQ ID NO:2. It is to be noted, however, that the present invention also includes a DNA sequence complementary to such a single-stranded DNA sequence and a component comprising both. The sequence of the gene of the present invention as shown under SEQ ID NO:3n - 1 (where n is an integer of 1 to 14) is merely an example of the codon combination encoding the respective amino acid residues. The gene of the present invention is not limited thereto but can of course have a DNA sequence in which the codons are arbitrarily selected and combined for the respective amino acid residues. The codon selection can be made in the conventional manner, for example taking into consideration the codon utilization frequencies in the host to be used [Nucl. Acids Res., 9, 43-74 (1981)].

The gene of the present invention further includes DNA sequences coding for functional equivalents derived from the amino acid sequence mentioned above by partial amino acid or amino acid sequence substitution, deletion or addition. These polypeptides may be produced by spontaneous modification (mutation) or may be obtained by posttranslational modification or by modifying the natural gene (of the present invention) by a technique of genetic engineering, for example by site-specific mutagenesis [Methods in Enzymology, 154, p. 350, 367-382 (1987); *ibid.*, 100, p. 468 (1983); Nucleic Acids Research, 12, p. 9441 (1984); Zoku Seikagaku Jikken Koza (Sequel to Experiments in Biochemistry) 1, "Idensi Kenkyu-ho (Methods in Gene Research) II", edited by the Japan Biochemical Society, p. 105 (1986)] or synthesizing mutant DNAs by a chemical synthetic technique such as the phosphotriester method or phosphoramidite method [J. Am. Chem. Soc. 89, p. 4801 (1967); *ibid.*, 91, p. 3350 (1969); Science, 150, p. 178 (1968); Tetrahedron Lett., 22, p. 1859 (1981); *ibid.*, 24, p. 245 (1983)], or by utilizing the techniques mentioned above in combination.

The protein encoded by the gene of the present invention can be expressed readily and stably by utilizing said gene, for example inserting it into a vector for use with a microorganism and cultivating the microorganism thus transformed.

The protein obtained by utilizing the gene of the present invention can be used in specific antibody production. In this case, the protein producible in large quantities by the genetic engineering technique mentioned above can be used as the component to serve as an antigen. The antibody obtained may be polyclonal or monoclonal and can be advantageously used in the purification, assay, discrimination or identification of the corresponding protein.

The gene of the present invention can be readily produced based on the sequence information thereof disclosed herein by using general genetic engineering techniques [cf. e.g. Molecular Cloning, 2nd Ed., Cold Spring Harbor Laboratory Press (1989); Zoku Seikagaku Jikken Koza, "Idensi Kenkyu-ho I, II and III", edited by the Japan Biochemical Society (1986)].

This can be achieved, for example, by selecting a desired clone from a human cDNA library (prepared in the conventional manner from appropriate cells of origin in which the gene is expressed) using a probe or antibody specific to the gene of the present invention [e.g. Proc. Natl. Acad. Sci. USA, 78, 6613 (1981); Science, 222, 778 (1983)].

The cells of origin to be used in the above method are, for example, cells or tissues in which the gene in question is expressed, or cultured cells derived therefrom. Separation of total RNA, separation and purification of mRNA, conversion to (synthesis of) cDNA, cloning thereof and so on can be carried out by conventional methods. cDNA libraries are also commercially available and such cDNA libraries, for example various cDNA libraries available from Clontech Lab. Inc. can also be used in the above method.

Screening of the gene of the present invention from these cDNA libraries can be carried out by the conventional method mentioned above. These screening methods include, for example, the method comprising selecting a cDNA clone by immunological screening using an antibody specific to the protein produced by the corresponding cDNA, the technique of plaque or colony hybridization using probes selectively binding to the desired DNA sequence, or a combination of these. As regards the probe to be used here, a DNA sequence chemically synthesized based on the information about the DNA sequence of the present invention is generally used. It is of course possible to use the gene of the present invention or fragments thereof as the probe.

Furthermore, a sense primer and an antisense primer designed based on the information about the partial amino acid sequence of a natural extract isolated and purified from cells or a tissue can be used as probes for screening.

For obtaining the gene of the present invention, the technique of DNA/RNA amplification by the PCR method [Science, 230, 1350-1354 (1984)] can suitably be employed. Particularly when the full-length cDNA can hardly be obtained from the library, the RACE method (rapid amplification of cDNA ends; Jikken Igaku (Experimental Medicine), 12 (6), 35-38 (1994)), in particular the 5'RACE method [Frohman, M. A., et al., Proc. Natl. Acad. Sci. USA, 85, 8998-9002 (1988)] is preferably employed. The primers to be used in such PCR method can be appropriately designed based on the sequence information of the gene of the present invention as disclosed herein and can be synthesized by a conventional method.

The amplified DNA/RNA fragment can be isolated and purified by a conventional method as mentioned above, for example by gel electrophoresis.

The nucleotide sequence of the thus-obtained gene of the present invention or any of various DNA fragments can be determined by a conventional method, for example the dideoxy method [Proc. Natl. Acad. Sci. USA, 74, 5463-5467 (1977)] or the Maxam-Gilbert method [Methods in Enzymology, 65, 499 (1980)]. Such nucleotide sequence determination can be readily performed using a commercially available sequence kit as well.

When the gene of the present invention is used and conventional techniques of recombinant DNA technology [see e.g. Science, 224, p. 1431 (1984); Biochem. Biophys. Res. Comm., 130, p. 692 (1985); Proc. Natl. Acad. Sci. USA 80, p. 5990 (1983) and the references cited above] are followed, a recombinant protein can be obtained. More detailedly, said protein can be produced by constructing a recombinant DNA enabling the gene of the present invention to be expressed in host cells, introducing it into host cells for transformation thereof and cultivating the resulting transformant.

In that case, the host cells may be eukaryotic or prokaryotic. The eukaryotic cells include vertebrate cells, yeast cells and so on, and the vertebrate cells include, but are not limited to, simian cells named COS cells [Cell, 23, 175-182 (1981)], Chinese hamster ovary cells and a dihydrofolate reductase-deficient cell line derived therefrom [Proc. Natl. Acad. Sci. USA, 77, 4216-4220 (1980)] and the like, which are frequently used.

As regards the expression vector to be used with vertebrate cells, an expression vector having a promoter located upstream of the gene to be expressed, RNA splicing sites, a polyadenylation site and a transcription termination sequence can be generally used. This may further have an origin of replication as necessary. As an example of said expression vector, there may be mentioned pSV2dhfr [Mol. Cell. Biol., 1, 854 (1981)], which has the SV40 early promoter. As for the eukaryotic microorganisms, yeasts are generally and frequently used and, among them, yeasts of the genus *Saccharomyces* can be used with advantage. As regards the expression vector for use with said yeasts and other eukaryotic microorganisms, pAM82 [Proc. Natl. Acad. Sci. USA, 80, 1-5 (1983)], which has the acid phosphatase gene promoter, for instance, can be used.

Furthermore, a prokaryotic gene fused vector can be preferably used as the expression vector for the gene of the present invention. As specific examples of said vector, there may be mentioned pGEX-2TK and pGEX-4T-2 which have a GST domain (derived from *S. japonicum*) with a molecular weight of 26,000.

*Escherichia coli* and *Bacillus subtilis* are generally and preferably used as prokaryotic hosts. When these are used as hosts in the practice of the present invention, an expression plasmid derived from a plasmid vector capable of replicating in said host organisms and provided in this vector with a promoter and the SD (Shine and Dalgarno) sequence upstream of said gene for enabling the expression of the gene of the present invention and further provided with an initiation codon (e.g. ATG) necessary for the initiation of protein synthesis is preferably used. The *Escherichia coli* strain K12, among others, is preferably used as the host *Escherichia coli*, and pBR322 and modified vectors derived therefrom are generally and preferably used as the vector, while various known strains and vectors can also be used. Examples of the promoter which can be used are the tryptophan (trp) promoter, lpp promoter, lac promoter and PL/PR promoter.

The thus-obtained desired recombinant DNA can be introduced into host cells for transformation by using various general methods. The transformant obtained can be cultured by a conventional method and the culture leads to expression and production of the desired protein encoded by the gene of the present invention. The medium to be used in said culture can suitably be selected from among various media in conventional use according to the host cells employed. The host cells can be cultured under conditions suited for the growth thereof.

In the above manner, the desired recombinant protein is expressed and produced and accumulated or secreted within the transformant cells or extracellularly or on the cell membrane.

The recombinant protein can be separated and purified as desired by various separation procedures utilizing the



physical, chemical and other properties thereof [cf. e.g. "Seikagaku (Biochemistry) Data Book II", pages 1175-1259, 1st Edition, 1st Printing, published June 23, 1980 by Tokyo Kagaku Dojin; Biochemistry, 25 (25), 8274-8277 (1986); Eur. J. Biochem., 163, 313-321 (1987)]. Specifically, said procedures include, among others, ordinary reconstitution treatment, treatment with a protein precipitating agent (salting out), centrifugation, osmotic shock treatment, sonication, ultrafiltration, various liquid chromatography techniques such as molecular sieve chromatography (gel filtration), adsorption chromatography, ion exchange chromatography, affinity chromatography and high-performance liquid chromatography (HPLC), dialysis and combinations thereof. Among them, affinity chromatography utilizing a column with the desired protein bound thereto is particularly preferred.

Furthermore, on the basis of the sequence information about the gene of the present invention as revealed by the present invention, for example by utilizing part or the whole of said gene, it is possible to detect the expression of the gene of the present invention in various human tissues. This can be performed by a conventional method, for example by RNA amplification by RT-PCR (reverse transcribed-polymerase chain reaction) [Kawasaki, E. S., et al., Amplification of RNA, in PCR Protocol, A guide to methods and applications, Academic Press, Inc., San Diego, 21-27 (1991)], or by northern blotting analysis [Molecular Cloning, Cold Spring Harbor Laboratory (1989)], with good results.

The primers to be used in employing the above-mentioned PCR method are not limited to any particular ones provided that they are specific to the gene of the present invention and enable the gene of the present invention alone to be specifically amplified. They can be designed or selected appropriately based on the gene information provided by the present invention. They can have a partial sequence comprising about 20 to 30 nucleotides according to the established practice. Suitable examples are as shown in Examples 1 to 11.

Thus, the present invention also provides primers and/or probes useful in specifically detecting such novel gene.

By using the novel gene provided by the present invention, it is possible to detect the expression of said gene in various tissues, analyze the structure and function thereof and, further, produce the human protein encoded by said gene in the manner of genetic engineering. These make it possible to analyze the expression product, reveal the pathology of a disease associated therewith, for example a genopathy or cancer, and diagnose and treat the disease.

The following drawings are referred to in the examples.

Fig. 1 shows the result obtained by testing the PI4 kinase activity of NPIK in Example 9. Fig. 2 shows the effect of Triton X-100 and adenosine on NPIK activity.

## EXAMPLES

The following examples illustrate the present invention in further detail.

### Example 1

#### GDP dissociation stimulator gene

##### (1) Cloning and DNA sequencing of GDP dissociation stimulator gene

mRNAs extracted from the tissues of human fetal brain, adult blood vessels and placenta were purchased from Clontech and used as starting materials.

cDNA was synthesized from each mRNA and inserted into the vector  $\lambda$ ZAPII (Stratagene) to thereby construct a cDNA library (Otsuka GEN Research Institute, Otsuka Pharmaceutical Co., Ltd.)

Human gene-containing *Escherichia coli* colonies were allowed to form on agar medium by the *in vivo* excision technique [Short, J. M., et al., Nucleic Acids Res., 16, 7583-7600 (1988)]. Colonies were picked up at random and human gene-containing *Escherichia coli* clones were registered on 96-well micro plates. The clones registered were stored at -80°C.

Each of the clones registered was cultured overnight in 1.5 ml of LB medium, and DNA was extracted and purified using a model PI-100 automatic plasmid extractor (Kurabo). Contaminant *Escherichia coli* RNA was decomposed and removed by RNase treatment. The DNA was dissolved to a final volume of 30  $\mu$ l. A 2- $\mu$ l portion was used for roughly checking the DNA size and quantity using a minigel, 7  $\mu$ l was used for sequencing reactions and the remaining portion (21  $\mu$ l) was stored as plasmid DNA at 4°C.

This method, after slight changes in the program, enables extraction of the cosmid, which is useful also as a probe for FISH (fluorescence in situ hybridization) shown later in the examples.

Then, the dideoxy terminator method of Sanger et al. [Sanger, F., et al., Proc. Natl. Acad. Sci. USA, 74, 5463-5467 (1977)] using T3, T7 or a synthetic oligonucleotide primer or the cycle sequence method [Carothers, A. M., et al., Bio. Techniques 7, 494-499 (1989)] comprising the dideoxy chain terminator method plus PCR method was carried out. These are methods of terminating the extension reaction specifically to the four bases using a small amount of plasmid DNA (about 0.1 to 0.5  $\mu$ g) as a template.

The sequence primers used were FITC (fluorescein isothiocyanate)-labeled ones. Generally, about 25 cycles of

reaction were performed using Taq polymerase. The PCR products were separated on a polyacrylamide urea gel and the fluorescence-labeled DNA fragments were submitted to an automatic DNA sequencer (ALF™ DNA Sequencer; Pharmacia) for determining the sequence of about 400 bases from the 5' terminus side of cDNA.

Since the 3' nontranslational region is high in heterogeneity for each gene and therefore suited for discriminating individual genes from one another, sequencing was performed on the 3' side as well depending on the situation.

The vast sum of nucleotide sequence information obtained from the DNA sequencer was transferred to a 64-bit DEC 3400 computer for homology analysis by the computer. In the homology analysis, a data base (GenBank, EMBL) was used for searching according to the UWGCG FASTA program [Pearson, W. R. and Lipman, D. J., Proc. Natl. Acad. Sci. USA, **85**, 2444-2448 (1988)].

As a result of arbitrary selection by the above method and of cDNA sequence analysis, a clone designated as GEN-501D08 and having a 0.8 kilobase insert was found to show a high level of homology to the C terminal region of the human Ral guanine nucleotide dissociation stimulator (RalGDS) gene. Since RalGDS is considered to play a certain role in signal transmission pathways, the whole nucleotide sequence of the cDNA insert portion providing the human homolog was further determined.

Low-molecular GTPases play an important role in transmitting signals for a number of cell functions including cell proliferation, differentiation and transformation [Bourne, H. R. et al., Nature, **348**, 125-132 (1990); Bourne et al., Nature, **349**, 117-127 (1991)].

It is well known that, among them, those proteins encoded by the ras gene family function as molecular switches or, in other words, the functions of the ras gene family are regulated by different conditions of binding proteins such as biologically inactive GDP-binding proteins or active GTP-binding proteins, and that these two conditions are induced by GTPase activating proteins (GAPs) or GDS. The former enzymes induce GDP binding by stimulating the hydrolysis of bound GTP and the latter enzyme induces the regular GTP binding by releasing bound GDP [Boguski, M. S. and McCormick, F., Nature, **366**, 643-654 (1993)].

RalGDS was first discovered as a member of the ras gene family lacking in transforming activity and as a GDP dissociation stimulator specific to RAS [Chardin, P. and Tavittian, A., EMBO J., **5**, 2203-2208 (1986); Albright, C. F., et al., EMBO J., **12**, 339-347 (1993)].

In addition to Ral, RalGDS was found to function, through interaction with these proteins, as an effector molecule for N-ras, H-ras, K-ras and Rap [Spaargaren, M. and Bischoff, J. R., Proc. Natl. Acad. Sci. USA, **91**, 12609-12613 (1994)].

The nucleotide sequence of the cDNA clone designated as GEN-501D08 is shown under SEQ ID NO:3, the nucleotide sequence of the coding region of said clone under SEQ ID NO:2, and the amino acid sequence encoded by said nucleotide sequence under SEQ ID NO:1.

This cDNA comprises 842 nucleotides, including an open reading frame comprising 366 nucleotides and coding for 122 amino acids. The translation initiation codon was found to be located at the 28th nucleotide residue.

Comparison between the RalGDS protein known among conventional databases and the amino acid sequence deduced from said cDNA revealed that the protein encoded by this cDNA is homologous to the C terminal domain of human RalGDS. The amino acid sequence encoded by this novel gene was found to be 39.5% identical with the C terminal domain of RalGDS which is thought to be necessary for binding to ras.

Therefore, it is presumable, as mentioned above, that this gene product might interact with the ras family proteins or have influence on the ras-mediated signal transduction pathways. However, this novel gene is lacking in the region coding for the GDS activity domain and the corresponding protein seems to be different in function from the GDS protein. This gene was named human RalGDS by the present inventors.

## (2) Northern blot analysis

The expression of the RalGDS protein mRNA in normal human tissues was evaluated by Northern blotting using, as a probe, the human cDNA clone labeled by the random oligonucleotide priming method.

The Northern blot analysis was carried out with a human MTN blot (Human Multiple Tissue Northern blot; Clontech, Palo Alto, CA, USA) according to the manufacturer's protocol.

Thus, the PCR amplification product from the above GEN-501D08 clone was labeled with [<sup>32</sup>P]-dCTP (random-primed DNA labeling kit, Boehringer-Mannheim) for use as a probe.

For blotting, hybridization was performed overnight at 42°C in a solution comprising 50% formamide/5 x SSC/50 x Denhardt's solution/0.1% SDS (containing 100 µg/ml denatured salmon sperm DNA). After washing with two portions of 2 x SSC/0.01% SDS at room temperature, the membrane filter was further washed three times with 0.1 x SSC/0.05% SDS at 50°C for 40 minutes. An X-ray film (Kodak) was exposed to the filter at -70°C for 18 hours.

As a result, it was revealed that a 900-bp transcript had been expressed in all the human tissues tested. In addition, a 3.2-kb transcript was observed specifically in the heart and skeletal muscle. The expression of these transcripts differing in size may be due either to alternative splicing or to cross hybridization with homologous genes.

## (3) Cosmid clone and chromosome localization by FISH

FISH was performed by screening a library of human chromosomes cloned in the cosmid vector pWE15 using, as a probe, the 0.8-kb insert of the cDNA clone [Sambrook, J., et al., Molecular Cloning, 2nd Ed., pp. 3.1-3.58, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (1989)].

FISH for chromosome assignment was carried out by the method of Inazawa et al. which comprises G-banding pattern comparison for confirmation [Inazawa, J., et al., Genomics, 17, 153-162 (1993)].

For use as a probe, the cosmid DNA (0.5 µg) obtained from chromosome screening and corresponding to GEN-501D08 was labeled with biotin-16-dUTP by nick translation.

To eliminate the background noise due to repetitive sequences, 0.5 µl of sonicated human placenta DNA (10 mg/ml) was added to 9.5 µl of the probe solution. The mixture was denatured at 80°C for 5 minutes and admixed with an equal volume of 4 x SSC containing 20% dextran sulfate. Then, a denatured slide was sown with the hybridization mixture and, after covering with paraffin, incubated in a wet chamber at 37°C for 16 to 18 hours. After washing with 50% formamide/2 x SSC at 37°C for 15 minutes, the slide was washed with 2 x SSC for 15 minutes and further with 1 x SSC for 15 minutes.

The slide was then incubated in 4 x SSC supplemented with "1% Block Ace" (trademark; Dainippon Pharmaceutical) containing avidin-FITC (5 µg/ml) at 37°C for 40 minutes. Then, the slide was washed with 4 x SSC for 10 minutes and with 4 x SSC containing 0.05% Triton X-100 for 10 minutes and immersed in an antifading PPD solution [prepared by adjusting 100 mg of PPD (Wako Catalog No. 164-015321) and 10 ml of PBS(-) (pH 7.4) to pH 8.0 with 0.5 M Na<sub>2</sub>CO<sub>3</sub>/0.5 M NaHCO<sub>3</sub> (9:1, v/v) buffer (pH 9.0) and adding glycerol to make a total volume of 100 ml] containing 1% DABCO (Sigma) in PBS(-):glycerol 1:9 (v:v), followed by counter staining with DAPI (4,6-diamino-2-phenylindole; Sigma).

With more than 100 tested cells in the metaphase, a specific hybridization signal was observed on the chromosome band at 6p21.3, without any signal on other chromosomes. It was thus confirmed that the RalGDS gene is located on the chromosome 6p21.3.

By using the novel human RalGDS-associated gene of the present invention as obtained in this example, the expression of said gene in various tissues can be detected and the human RalGDS protein can be produced in the manner of genetic engineering. These are expected to enable studies on the roles of the expression product protein and ras-mediated signals in transduction pathways as well as pathological investigations of diseases in which these are involved, for example cancer, and the diagnosis and treatment of such diseases. Furthermore, it becomes possible to study the development and progress of diseases involving the same chromosomal translocation of the RalGDS protein gene of the present invention, for example tonic spondylitis, atrial septal defect, pigmentary retinopathy, aphasia and the like.

Example 2

Cytoskeleton-associated protein 2 gene (CKAP2 gene)

## (1) Cytoskeleton-associated protein 2 gene cloning and DNA sequencing

cDNA clones were arbitrarily chosen from a human fetal brain cDNA library in the same manner as in Example 1 were subjected to sequence analysis and, as a result, a clone having a base sequence containing the CAP-glycine domain of the human cytoskeleton-associated protein (CAP) gene and highly homologous to several CAP family genes was found and named GEN-080G01.

Meanwhile, the cytoskeleton occurs in the cytoplasm and just inside the cell membrane of eukaryotic cells and is a network structure comprising complicatedly entangled filaments. Said cytoskeleton is constituted of microtubules composed of tubulin, microfilaments composed of actin, intermediate filaments composed of desmin and vimentin, and so on. The cytoskeleton not only acts as supportive cellular elements but also isokinetically functions to induce morphological changes of cells by polymerization and depolymerization in the fibrous system. The cytoskeleton binds to intracellular organelles, cell membrane receptors and ion channels and thus plays an important role in intracellular movement and locality maintenance thereof and, in addition, is said to have functions in activity regulation and mutual information transmission. Thus it supposedly occupies a very important position in physiological activity regulation of the whole cell. In particular, the relation between canceration of cells and qualitative changes of the cytoskeleton attracts attention since cancer cells differ in morphology and recognition response from normal cells.

The activity of this cytoskeleton is modulated by a number of cytoskeleton-associated proteins (CAPs). One group of CAPs is characterized by a glycine motif highly conserved and supposedly contributing to association with microtubules [CAP-GLY domain; Riehemann, K. and Song, C., Trends Biochem. Sci., 18, 82-83 (1993)].

Among the members of this group of CAPs, there are CLIP-170, 150 kDa DAP (dynein-associated protein, or dynactin), *D. melanogaster* GLUED, *S. cerevisiae* BIK1, restin [Bilbe, G., et al., EMBO J., 11, 2103-2113 (1992)]; Hilliker,

C., et al., Cytogenet. Cell Genet., 65, 172-176 (1994)] and *C. elegans* 13.5 kDa protein [Wilson, R., et al., Nature, 368, 32-38 (1994)]. Except for the last two proteins, direct or indirect evidences have suggested that they could interact with microtubules.

The above-mentioned CLIP-170 is essential for the *in vitro* binding of endocytic vesicles to microtubules and colocalizes with endocytic organelles [Rickard, J. E. and Kreis, T. E., J. Biol. Chem., 18, 82-83 (1990); Pierre, P., et al., Cell, 70, 887-900 (1992)].

The above-mentioned dynactin is one of the factors constituting the cytoplasmic dynein motor, which functions in retrograde vesicle transport [Schroer, T. A. and Sheetz, M. P., J. Cell Biol., 115, 1309-1318 (1991)] or probably in the movement of chromosomes during mitosis [Pfarr, C. M., et al., Nature, 345, 263-265 (1990); Steuer, E. R., et al., Nature, 345, 266-268 (1990); Wordeman, L., et al., J. Cell Biol., 114, 285-294 (1991)].

GLUED, the *Drosophila* homolog of mammalian dynactin, is essential for the viability of almost all cells and for the proper organization of some neurons [Swaroop, A., et al., Proc. Natl. Acad. Sci. USA, 84, 6501-6505 (1987); Holzbaur, E. L. P., et al., Nature, 351, 579-583 (1991)].

BIK1 interacts with microtubules and plays an important role in spindle formation during mitosis in yeasts [Trueheart, J., et al., Mol. Cell. Biol., 7, 2316-2326 (1987); Berlin, V., et al., J. Cell Biol., 111, 2573-2586 (1990)].

At present, these genes are classified under the term CAP family (CAPs).

As a result of database searching, the above-mentioned cDNA clone of 463-bp (excluding the poly-A signal) showed significant homology in nucleotide sequence with the restin and CLIP-170 encoding genes. However, said clone was lacking in the 5' region as compared with the restin gene and, therefore, the technique of 5' RACE [Frohman, M. A., et al., Proc. Natl. Acad. Sci. USA 85, 8998-9002 (1988)] was used to isolate this missing segment.

## (2) 5' RACE (5' rapid amplification of cDNA ends)

A cDNA clone containing the 5' portion of the gene of the present invention was isolated for analysis by the 5' RACE technique using a commercial kit (5'-Rapid AmpliFinder RACE kit, Clontech) according to the manufacturer's protocol with minor modifications, as follows.

The gene-specific primer P1 and primer P2 used here were synthesized by the conventional method and their nucleotide sequences are as shown below in Table 1. The anchor primer used was the one attached to the commercial kit.

Table 1

Primer	Nucleotide sequence
Primer P1	5'-ACACCAATCCAGTAGCCAGGCTTG-3'
Primer P2	5'-CACTCGAGAATCTGTGAGACCTACATACATGACG-3'

cDNA was obtained by reverse transcription of 0.1 µg of human fetal brain poly(A)+RNA by the random hexamer technique using reverse transcriptase (Superscript™ II, Life Technologies) and the cDNA was amplified by the first PCR using the P1 primer and anchor primer according to Watanabe et al. [Watanabe, T., et al., Cell Genet., in press].

Thus, to 0.1 µg of the above-mentioned cDNA were added 2.5 mM dNTP/1 x Taq buffer (Takara Shuzo)/0.2 µM P1 primer, 0.2 µM adaptor primer/0.25 unit ExTaq enzyme (Takara Shuzo) to make a total volume of 50 µl, followed by addition of the anchor primer. The mixture was subjected to PCR. Thus, 35 cycles of amplification were performed under the conditions: 94°C for 45 seconds, 60°C for 45 seconds, and 72°C for 2 minutes. Finally, the mixture was heated at 72°C for 5 minutes.

Then, 1 µl of the 50-µl first PCR product was subjected to amplification by the second PCR using the specific nested P2 primer and anchor primer. The second PCR product was analyzed by 1.5% agarose gel electrophoresis.

Upon agarose gel electrophoresis, a single band, about 650 nucleotides in size, was detected. The product from this band was inserted into a vector (pT7Blue(R)-Vector, Novagen) and a plurality of clones with an insert having an appropriate size were selected.

Six of the 5' RACE clones obtained from the PCR product had the same sequence but had different lengths. By sequencing two overlapping cDNA clones, GEN-080G01 and GEN-080G0149, the protein-encoding sequence and 5' and 3' flanking sequences, 1015 nucleotides in total length, were determined. Said gene was named cytoskeleton-associated protein 2 gene (CKAP2 gene).

The nucleotide sequence obtained from the above-mentioned two overlapping cDNA clones GEN-080G01 and GEN-080G0149 is shown under SEQ ID NO:6, the nucleotide sequence of the coding region of said clone under SEQ ID NO:5, and the amino acid sequence encoded by said nucleotide sequence under SEQ ID NO:4.

As shown under SEQ ID NO:6, the CKAP2 gene had a relatively GC-rich 5' noncoding region, with incomplete triplet repeats, (CAG)<sub>4</sub>(CGG)<sub>4</sub>(CTG)(CGG), occurring at nucleotides 40-69.

ATG located at nucleotides 274-276 is the presumable start codon. A stop codon (TGA) was situated at nucleotides 853-855. A polyadenylation signal (ATTAAA) was followed by 16 nucleotides before the poly(A) start. The estimated open reading frame comprises 579 nucleotides coding for 193 amino acid residues with a calculated molecular weight of 21,800 daltons.

The coding region was further amplified by RT-PCR, to eliminate the possibility of the synthetic sequence obtained being a cDNA chimera.

## (2) Similarity of CKAP2 to other CAPs

While sequencing of CKAP2 revealed homology with the sequences of restin and CLIP-170, the homologous region was limited to a short sequence corresponding to the CAP-GLY domain. On the amino acid level, the deduced CKAP2 was highly homologous to five other CAPs in this domain.

CKAP2 was lacking in such other motif characteristics of some CAPs as the alpha helical rod and zinc finger motif. The alpha helical rod is thought to contribute to dimerization and to increase the microtubule binding capacity [Pierre, P., et al., *Cell*, **70**, 887-900 (1992)]. The lack of the alpha helical domain might mean that CKAP2 be incapable of homo or hetero dimer formation.

Paralleling of the CAP-GLY domains of these proteins revealed that other conserved residues other than glycine residues are also found in CKAP2. CAPs having a CAP-GLY domain are thought to be associated with the activities of cellular organelles and the interactions thereof with microtubules. Since it contains a CAP-GLY domain, as mentioned above, CKAP2 is placed in the family of CAPs.

Studies with mutants of Glued have revealed that the Glued product plays an important role in almost all cells [Swaroop, A., et al., *Proc. Natl. Acad. Sci. USA*, **84**, 6501-6505 (1987)] and that it has other neuron-specific functions in neuronal cells [Meyerowitz, E. M. and Kankel, D. R., *Dev. Biol.*, **62**, 112-142 (1978)]. These microtubule-associated proteins are thought to function in vesicle transport and mitosis. Because of the importance of the vesicle transport system in neuronal cells, defects in these components might lead to aberrant neuronal systems.

In view of the above, CKAP2 might be involved in specific neuronal functions as well as in fundamental cellular functions.

## (3) Northern blot analysis

The expression of human CKAP2 mRNA in normal human tissues was examined by Northern blotting in the same manner as in Example 1 (2) using the GEN-080G01 clone (corresponding to nucleotides 553-1015) as a probe.

As a result, in all the eight tissues tested, namely human heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas, a 1.0 kb transcript agreeing in size with the CKAP2 cDNA was detected. Said 1.0 kb transcript was expressed at significantly higher levels in heart and brain than in the other tissues examined. Two weak bands, 3.4 kb and 4.6 kb, were also detected in all the tissues examined.

According to the Northern blot analysis, the 3.4 kb and 4.6 kb transcripts might possibly be derived from the same gene coding for the 1.0 kb CKAP2 by alternative splicing or transcribed from other related genes. These characteristics of the transcripts may indicate that CKAP2 might also code for a protein having a CAP-GLY domain as well as an alpha helix.

## (4) Cosmid cloning and chromosomal localization by direct R-banding FISH

Two cosmids corresponding to the CKAP2 cDNA were obtained. These two cosmid clones were subjected to direct R-banding FISH in the same manner as in Example 1

## (3) for chromosomal locus mapping of CKAP2.

For suppressing the background due to repetitive sequences, a 20-fold excessive amount of human Cot-I DNA (BRL) was added as described by Lichter et al. [Lichter, P., et al., *Proc. Natl. Acad. Sci. USA*, **87**, 6634-6638 (1990)]. A Provia 100 film (Fuji ISO 100; Fuji Photo Film) was used for photomicrography.

As a result, CKAP2 was mapped on chromosome bands 19q13.11-q13.12.

Two autosomal dominant neurological diseases have been localized to this region by linkage analysis: CADASIL (cerebral autosomal dominant arteriopathy with subcortical infarcts and leukoencephalopathy) between the DNA markers D19S221 and D19S222, and FHM (familial hemiplegic migraine) between D19S215 and D19S216. These two diseases may be allelic disorders in which the same gene is involved [Tournier-Lasserre, E., et al., *Nature Genet.*, **3**, 256-259 (1993); Joutel, A., et al., *Nature Genet.*, **5**, 40-45 (1993)].

Although no evidence is available to support CKAP2 as a candidate gene for FHM or CADASIL, it is conceivable that its mutation might lead to some or other neurological disease.

By using the novel human CKAP2 gene of the present invention as obtained in this example, it is possible to detect the expression of said gene in various tissues or produce the human CKAP2 gene in the manner of genetic engineering. Through these, it becomes possible to analyze the functions of the human CKAP2 system or human CKAP2, which is involved in diverse activities essential to cells, as mentioned above, to diagnose various neurological diseases in which said system or gene is involved, for example familial migraine, and to screen out and evaluate a therapeutic or prophylactic drug therefor.

### Example 3

#### OTK27 gene

##### (1) OTK27 gene cloning and DNA sequencing

As a result of sequence analysis of cDNA clones arbitrarily selected from a human fetal brain cDNA library in the same manner as in Example 1 (1) and database searching, a cDNA clone, GEN-025F07, coding for a protein highly homologous to NHP2, a yeast nucleoprotein [*Saccharomyces cerevisiae*; Kolodrubetz, D. and Burgum, A., YEAST, 7, 79-90 (1991)], was found and named OTK27.

Nucleoproteins are fundamental cellular constituents of chromosomes, ribosomes and so forth and are thought to play an essential role in cell multiplication and viability. The yeast nucleoprotein NHP2, a high-mobility group (HMG)-like protein, like HMG, has reportedly a function essential for cell viability [Kolodrubetz, D. and Burgum, A., YEAST, 7, 79-90 (1991)].

The novel human gene, OTK27 gene, of the present invention, which is highly homologous to the above-mentioned yeast NHP2 gene, is supposed to be similar in function.

The nucleotide sequence of said GEN-025F07 clone was found to comprise 1493 nucleotides, as shown under SEQ ID NO:9, and contain an open reading frame comprising 384 nucleotides, as shown under SEQ ID NO:8, coding for an amino acid sequence comprising 128 amino acid residues, as shown under SEQ ID NO:7. The initiation codon was located at nucleotides 95-97 of the sequence shown under SEQ ID NO:9, and the termination codon at nucleotides 479-481.

At the amino acid level, the OTK27 protein was highly homologous (38%) to NHP2. It was 83% identical with the protein deduced from the cDNA from *Arabidopsis thaliana*;

Newman, T., unpublished; GENEMBL Accession No. T14197).

##### (2) Northern blot analysis

For examining the expression of human OTK27 mRNA in normal human tissues, the insert in the OTK27 cDNA was amplified by PCR, the PCR product was purified and labeled with [<sup>32</sup>P]-dCTP (random-primed DNA labeling kit, Boehringer Mannheim), and Northern blotting was performed using the labeled product as a probe in the same manner as in Example 1 (2).

As a result of the Northern blot analysis, two bands corresponding to possible transcripts from this gene were detected at approximately 1.6 kb and 0.7 kb. Both sizes of transcript were expressed in all normal adult tissues examined. However, the expression of the 0.7 kb transcript was significantly reduced in brain and was of higher levels in heart, skeletal muscle and testicle than in other tissues examined.

For further examination of these two transcripts, eleven cDNA clones were isolated from a testis cDNA library and their DNA sequences were determined in the same manner as in Example 1 (1).

As a result, in six clones, the sequences were found to be in agreement with that of the 0.7 kb transcript, with a poly(A) sequence starting at around the 600th nucleotide, namely at the 598th nucleotide in two of the six clones, at the 606th nucleotide in three clones, and at the 613th nucleotide in one clone.

In these six clones, the "TATAAA" sequence was recognized at nucleotides 583-588 as a probable poly(A) signal. The upstream poly(A) signal "TATAAA" of this gene was recognized as little influencing in brain and more effective in the three tissues mentioned above than in other tissues. The possibility was considered that the stability of each transcript vary from tissue to tissue.

Results of zoo blot analysis indicated that this gene is well conserved also in other vertebrates. Since this gene is expressed ubiquitously in normal adult tissues and conserved among a wide range of species, the gene product is likely to play an important physiological role. The evidence that yeasts lacking in NHP2 are nonviable suggests that the human homolog may also be essential to cell viability.

## (3) Chromosomal localization of OTK27 by direct R-banding FISH

One cosmid clone corresponding to the cDNA OTK27 was isolated from a total human genomic cosmid library (5-genome equivalent) using the OTK27 cDNA insert as a probe and subjected to FISH in the same manner as in Example 1 (3) for chromosomal localization of OTK27.

As a result, two distinct spots were observed on the chromosome band 12q24.3.

The OTK27 gene of the present invention can be used in causing expression thereof and detecting the OTK27 protein, a human nucleoprotein, and thus can be utilized in the diagnosis and pathologic studies of various diseases in which said protein is involved and, because of its involvement in cell proliferation and differentiation, in screening out and evaluating therapeutic and preventive drugs for cancer.

Example 4

## OTK18 gene

## (1) OTK18 gene cloning and DNA sequencing

Zinc finger proteins are defined as constituting a large family of transcription-regulating proteins in eukaryotes and carry evolutionally conserved structural motifs [Kadonaga, J. T., et al., Cell, 51, 1079-1090 (1987); Klung, A. and Rhodes, D., Trends Biol. Sci., 12, 464-469 (1987); Evans, R. M. and Hollenberg, S. M., Cell, 52, 1-3 (1988)].

The zinc finger, a loop-like motif formed by the interaction between the zinc ion and two residues, cysteine and histidine residues, is involved in the sequence-specific binding of a protein to RNA or DNA. The zinc finger motif was first identified within the amino acid sequence of the Xenopus transcription factor IIIA [Miller, J., et al., EMBO J., 4, 1609-1614 (1986)].

The C<sub>2</sub>H<sub>2</sub> zinc finger motif is in general tandemly repeated and contains an evolutionally conserved intervening sequence of 7 or 8 amino acids. This intervening stretch was first identified in the Kruppel segmentation gene of Drosophila [Rosenberg, U. B., et al., Nature, 319, 336-339 (1986)]. Since then, hundreds of C<sub>2</sub>H<sub>2</sub> zinc finger protein-encoding genes have been found in vertebrate genomes.

As a result of sequence analysis of cDNA clones arbitrarily selected from a human fetal brain cDNA library in the same manner as in Example 1 (1) and database searching, several zinc finger structure-containing clones were identified and, further, a clone having a zinc finger structure of the Kruppel type was found.

Since this clone lacked the 5' portion of the transcript, plaque hybridization was performed with a fetal brain cDNA library using, as a probe, an approximately 1.8 kb insert in the cDNA clone, whereby three clones were isolated. The nucleotide sequences of these were determined in the same manner as in Example 1 (1).

Among the three clones, the one having the largest insert spans 3,754 nucleotides including an open reading frame of 2,133 nucleotides coding for 711 amino acids. It was found that said clone contains a novel human gene coding for a peptide highly homologous in the zinc finger domain to those encoded by human ZNF41 and the Drosophila Kruppel gene. This gene was named OTK18 gene (derived from the clone GEN-076C09).

The nucleotide sequence of the cDNA clone of the OTK18 gene is shown under SEQ ID NO:12, the coding region-containing nucleotide sequence under SEQ ID NO:11, and the predicted amino acid sequence encoded by said OTK18 gene under SEQ ID NO:10.

It was found that the amino acid sequence of OTK18 as deduced from SEQ ID NO:12 contains 13 finger motifs on its carboxy side.

## (2) Comparison with other zinc finger motif-containing genes

Comparison among OTK18, human ZNF41 and the Drosophila Kruppel gene revealed that each finger motif is for the most part conserved in the consensus sequence CXECGKAFXQKSXLX<sub>2</sub>HQRXH.

Comparison of the consensus sequence of the zinc finger motifs of OTK18 with those of human ZNF41 and the Drosophila Kruppel gene revealed that the Kruppel type motif is well conserved in the OTK18-encoded protein. However, the sequence similarities were limited to zinc finger domains and no significant homologies were found with regard to other regions.

The zinc finger domain interacts specifically with the target DNA, recognizing an about 5 bp sequence to thereby bind to the DNA helix [Rhodes, D. and Klug, A., Cell 46, 123-132 (1986)].

Based on the idea that, in view of the above, the multiple module (tandem repetitions of zinc finger) can interact with long stretches of DNA, it is presumable that the target DNA of this gene product containing 13 repeated zinc finger units would be a DNA fragment with a length of approximately 65 bp.



## (3) Northern blot analysis

Northern blot analysis was performed as described in Example 1 (2) for checking normal human tissues for expression of the human OTK18 mRNA therein by amplifying the insert of the OTK18 cDNA by PCR, purifying the PCR product, labeling the same with [<sup>32</sup>P]-dCTP (random-primed DNA labeling kit, Boehringer Mannheim) and using an MTN blot with the labeled product as a probe.

The results of Northern blot analysis revealed that the transcript of OTK18 is approximately 4.3 kb long and is expressed ubiquitously in various normal adult tissues. However, the expression level in the liver and in peripheral blood lymphocytes seemed to be lower than in other organs tested.

## (4) Cosmid cloning and chromosomal localization by direct R-banding FISH

Chromosomal localization of OTK18 was carried out as described in Example 1 (3).

As a result, complete twin spots were identified with 8 samples while 23 samples showed an incomplete signal or twin spots on either or both homologs. All signals appeared at the q13.4 band of chromosome 19. No twin spots were observed on any other chromosomes.

The results of FISH thus revealed that this gene is localized on chromosomal band 19q13.4. This region is known to contain many DNA segments that hybridize with oligonucleotides corresponding to zinc finger domains [Hoovers, J. M. N., et al., *Genomics*, **12**, 254-263 (1992)]. In addition, at least one other gene coding for a zinc finger domain has been identified in this region [Marine, J.-C., et al., *Genomics*, **21**, 285-286 (1994)].

Hence, the chromosome 19q13 is presumably a site of grouping of multiple genes coding for transcription-regulating proteins.

When the novel human OTK18 gene provided by this example is used, it becomes possible to detect expression of said gene in various tissues and produce the human OTK18 protein in the manner of genetic engineering. Through these, it is possible to analyze the functions of the human transcription regulating protein gene system or human transcription regulating proteins, which are deeply involved in diverse activities fundamental to cells, as mentioned above, to diagnose various diseases with which said gene is associated, for example malformation or cancer resulting from a developmental or differentiation anomaly, and mental or nervous disorder resulting from a developmental anomaly in the nervous system, and further to screen out and evaluate therapeutic or prophylactic drugs for these diseases.

Example 5

Genes encoding human 26S proteasome constituent P42 protein and P27 protein

## (1) Cloning and DNA sequencing of genes respectively encoding human 26S proteasome constituent P42 protein and P27 protein

Proteasome, which is a multifunctional protease, is an enzyme occurring widely in eukaryotes from yeasts to humans and decomposing ubiquitin-binding proteins in cells in an energy-dependent manner. Structurally, said proteasome is constituted of 20S proteasome composed of various constituents with a molecular weight of 21 to 31 kilodaltons and a group of PA700 regulatory proteins composed of various constituents with a molecular weight of 30 to 112 kilodaltons and showing a sedimentation coefficient of 22S and, as a whole, occurs as a macromolecule with a molecular weight of about 2 million daltons and a sedimentation coefficient of 26S [Rechsteiner, M., et al., *J. Biol. Chem.*, **268**, 6065-6068 (1993); Yoshimura, T., et al., *J. Struct. Biol.*, **111**, 200-211 (1993); Tanaka, K., et al., *New Biologist*, **4**, 173-187 (1992)].

Despite structural and mechanical analyses thereof, the whole picture of proteasome is not yet fully clear. However, according to studies using yeasts and mice in the main, it reportedly has the functions mentioned below and its functions are becoming more and more elucidated.

The mechanism of energy-dependent proteolysis in cells starts with selection of proteins by ubiquitin binding. It is not 20S proteasome but 26S proteasome that has ubiquitin-conjugated protein decomposing activity which is ATP-dependent [Chu-Ping et al., *J. Biol. Chem.*, **269**, 3539-3547 (1994)]. Hence, human 26S proteasome is considered to be useful in elucidating the mechanism of energy-dependent proteolysis.

Factors involved in the cell cycle regulation are generally short in half-life and in many cases they are subject to strict quantitative control. In fact, it has been made clear that the oncogene products Mos, Myc, Fos and so forth can be decomposed by 26S proteasome in an energy- and ubiquitin-dependent manner [Ishida, N., et al., *FEBS Lett.*, **324**, 345-348 (1993); Hershko, A. and Ciechanover, A., *Annu. Rev. Biochem.*, **61**, 761-807 (1992)] and the importance of proteasome in cell cycle control is being recognized.

Its importance in the immune system has also been pointed out. It is suggested that proteasome is positively involved in class I major histocompatible complex antigen presentation [Michalek, M. T., et al., *Nature*, **363**, 552-554

(1993)] and it is further suggested that proteasome may be involved in Alzheimer disease, since the phenomena of abnormal accumulation of ubiquitin-conjugated proteins in the brain of patients with Alzheimer disease [Kitaguchi, N., et al., *Nature*, **361**, 530-532 (1988)]. Because of its diverse functions such as those mentioned above, proteasome attracts attention from the viewpoint of its utility in the diagnosis and treatment of various diseases.

A main function of 26S proteasome is ubiquitin-conjugated protein decomposing activity. In particular, it is known that cell cycle-related gene products such as oncogene products and cyclins, typically c-Myc, are degraded via ubiquitin-dependent pathways. It has also been observed that the proteasome gene is expressed abnormally in liver cancer cells, renal cancer cells, leukemia cells and the like as compared with normal cells [Kanayama, H., et al., *Cancer Res.*, **51**, 6677-6685 (1991)] and that proteasome is abnormally accumulated in tumor cell nuclei. Hence, constituents of proteasome are expected to be useful in studying the mechanism of such canceration and in the diagnosis or treatment of cancer.

Also, it is known that the expression of proteasome is induced by interferon  $\gamma$  and so on and is deeply involved in antigen presentation in cells [Aki, M., et al., *J. Biochem.*, **115**, 257-269 (1994)]. Hence, constituents of human proteasome are expected to be useful in studying the mechanism of antigen presentation in the immune system and in developing immunoregulating drugs.

Furthermore, proteasome is considered to be deeply associated with ubiquitin abnormally accumulated in the brain of patients with Alzheimer disease. Hence, it is suggested that constituents of human proteasome should be useful in studying the cause of Alzheimer disease and in the treatment of said disease.

In addition to the utilization of expectedly multifunctional proteasome as such in the above manner, it is probably possible to produce antibodies using constituents of proteasome as antigens and use such antibodies in diagnosing various diseases by immunoassay. Its utility in this field of diagnosis is thus also a focus of interest.

Meanwhile, a protein having the characteristics of human 26S proteasome is disclosed, for example in Japanese Unexamined Patent Publication No. 292964/1993 and rat proteasome constituents are disclosed in Japanese Unexamined Patent Publication Nos. 268957/1993 and 317059/1993. However, no human 26S proteasome constituents are known. Therefore, the present inventors made a further search for human 26S proteasome constituents and successfully obtained two novel human 26S proteasome constituents, namely human 26S proteasome constituent P42 protein and human 26S proteasome constituent P27 protein, and performed cloning and DNA sequencing of the corresponding genes in the following manner.

#### (1) Purification of human 26S proteasome constituents P42 protein and P27 protein

Human proteasome was purified using about 100 g of fresh human kidney and following the method of purifying human proteasome as described in Japanese Unexamined Patent Publication No. 292964/1993, namely by column chromatography using BioGel A-1.5 m (5 x 90 cm, Bio-Rad), hydroxyapatite (1.5 x 15 cm, Bio-Rad) and Q-Sepharose (1.5 x 15 cm, Pharmacia) and glycerol density gradient centrifugation.

The thus-obtained human proteasome was subjected to reversed phase high performance liquid chromatography (HPLC) using a Hitachi model L6200 HPLC system. A Shodex RS Pak D4-613 (0.6 x 15 cm, Showa Denko) was used and gradient elution was performed with the following two solutions:

- First solution: 0.06% trifluoroacetic acid;
- Second solution: 0.05% trifluoroacetic acid, 70% acetonitrile.

An aliquot of each eluate fraction was subjected to 8.5% SDS-polyacrylamide electrophoresis under conditions of reduction with dithiothreitol. The P42 protein and P27 protein thus detected were isolated and purified.

The purified P42 and P27 proteins were respectively digested with 1  $\mu$ g of trypsin in 0.1 M Tris buffer (pH 7.8) containing 2 M urea at 37°C for 8 hours and the partial peptide fragments obtained were separated by reversed phase HPLC and their sequences were determined by Edman degradation. The results obtained are as shown below in Table 2.

Table 2

Partial protein		Amino acid sequence
P42	(1)	VLNISLW
	(2)	TLMELLNQMDGFDLHR
	(3)	AVSDFVVSEYXMXA
	(4)	EVDPLVYNX
	(5)	HGEIDYEAIVK
	(6)	LSXGFNGADLRNVXTEAGMFAIXAD
	(7)	MIMATNRPDTLDPALLRPGXL
	(8)	IHIDL PNEQARLDILK
	(9)	ATNGPRYVVVG
	(10)	EIDGRLK
	(11)	ALQSVGQIVGEVLK
	(12)	ILAGPITK
	(13)	XXVIELPLTNPELFQG
	(14)	VVSSSLVDK
	(15)	ALQDYRK
	(16)	EHREQLK
	(17)	KLESKLDYKPVR
P27	(1)	LVPTR
	(2)	AKEEEIEAIIK
	(3)	ANYEVLESQK
	(4)	VEDALHQLHAR
	(5)	DVDLYQVR
	(6)	QSQGLSPAQAFK
	(7)	AGSQSGGSPEASGVTVDVQE
	(8)	GLLGXNIPLQR

#### (2) cDNA library screening, clone isolation and cDNA nucleotide sequence determination

As mentioned in Example 1 (1), the present inventors have a database comprising about 30,000 cDNA data as constructed based on large-scale DNA sequencing using human fetal brain, arterial blood vessel and placenta cDNA libraries.

Based on the amino acid sequences obtained as mentioned above in (1), computer searching was performed with the FASTA program (search for homology between said amino acid sequences and the amino acid sequences estimated from the database). As regards P42, a clone (GEN-331G07) showing identity with regard to two amino acid sequences [(2) and (7) shown in table 2] was screened out and, as regards P27, a clone (GEN-163D09) showing identity with regard to two amino acid sequences [(1) and (8) shown in Table 2] was found.

For each of these clones, the 5' side sequence was determined by 5' RACE and the whole sequence was determined, in the same manner as in Example 2 (2).

As a result, it was revealed that the above-mentioned P42 clone GEN-331G07 comprises a 1,566-nucleotide sequence as shown under SEQ ID NO:15, inclusive of a 1,167-nucleotide open reading frame as shown under SEQ ID NO:14, and that the amino acid sequence encoded thereby is the one shown under SEQ ID NO:13 and comprises 389

amino acid residues.

The results of computer homology search revealed that the P42 protein is significantly homologous to the AAA (ATPase associated with a variety of cellular activities) protein family (e.g. P45, TBP1, TBP7, S4, MSS1, etc.). It was thus suggested that it is a new member of the AAA protein family.

As for the P27 clone GEN-163D09, it was revealed that it comprises a 1,128-nucleotide sequence as shown under SEQ ID NO:18, including a 669-nucleotide open reading frame as shown under SEQ ID NO:17 and that the amino acid sequence encoded thereby is the one shown under SEQ ID NO:16 and comprises 223 amino acid residues.

As regards the P27 protein, homology search using a computer failed to reveal any homologous gene among public databases. Thus, the gene in question is presumably a novel gene having an unknown function.

Originally, the above-mentioned P42 and P27 gene products were both purified as regulatory subunit components of proteasome complex. Therefore, these are expected to play an important role in various biological functions through proteolysis, for example a role in energy supply through decomposition of ATP and, hence, they are presumably useful not only in studying the function of human 26S proteasome but also in the diagnosis and treatment of various diseases caused by lowering of said biological functions, among others.

#### Example 6

#### BNAP gene

##### (1) BNAP gene cloning and DNA sequencing

The nucleosome composed of DNA and histone is a fundamental structure constituting chromosomes in eukaryotic cells and is well conserved over borders among species. This structure is closely associated with the processes of replication and transcription of DNA. However, the nucleosome formation is not fully understood as yet. Only certain specific factors involved in nucleosome assembly (NAPs) have been identified. Thus, two acidic proteins, nucleoplasmin and N1, are already known to facilitate nucleosome construction [Kleinschmidt, J. A., et al., J. Biol. Chem., 260, 1166-1176 (1985); Dilworth, S. M., et al., Cell 51, 1009-1018 (1987)].

A yeast gene, NAP-I, was isolated using a monoclonal antibody and recombinant proteins derived therefrom were tested as to whether they have nucleosome assembling activity *in vivo*.

More recently, a mouse NAP-I gene, which is a mammalian homolog of the yeast NAP-I gene was cloned (Okuda, A.; registered in database under the accession number D12618). Also cloned were a mouse gene, DN38 [Kato, K., Eur. J. Neurosci., 2, 704-711 (1990)] and a human nucleosome assembly protein (hNRP) [Simon, H. U., et al., Biochem. J., 297, 389-397 (1994)]. It was shown that the hNRP gene is expressed in many tissues and is associated with T lymphocyte proliferation.

The present inventors performed sequence analysis of cDNA clones arbitrarily chosen from a human fetal brain cDNA library in the same manner as in Example 1 (1), followed by searches among databases and, as a result, made it clear that a 1,125-nucleotide cDNA clone (free of poly(A)), GEN-078D05, is significantly homologous to the mouse NAP-I gene, which is a gene for a nucleosome assembly protein (NAP) involved in nucleosome construction, a mouse partial cDNA clone, DN38, and hNRP.

Since said clone GEN-078D05 was lacking in the 5' region, 5' RACE was performed in the same manner as in Example 2 (2) to obtain the whole coding region. For this 5' RACE, primers P1 and P2 respectively having the nucleotide sequences shown below in Table 3.

Table 3

Primer	Nucleotide sequence
Primer P1	5'-TTGAAGAATGATGCATTAGGAACCAC-3'
Primer P2	5'-CACTCGAGTGGCTGGATTTC AATTTCTCCAGTAG-3'

After the first 5' RACE, a single band corresponding to a sequence length of 1,300 nucleotides was obtained. This product was inserted into pT7Blue(R) T-Vector and several clones appropriate in insert size were selected.

Ten 5' RACE clones obtained from two independent PCR reactions were sequenced and the longest clone GEN-078D05TA13 (about 1,300 nucleotides long) was further analyzed.

Both strands of the two overlapping cDNA clones GEN-078D05 and GEN-078D05TA13 were sequenced, whereby it was confirmed that the two clones did not yet cover the whole coding region. Therefore, a further second 5' RACE was carried out. For the second 5' RACE, two primers, P3 and P4, respectively having the sequences shown below in

Table 4 were used.

Table 4

Primer	Nucleotide sequence
Primer P3	5'-GTCGAGCTAGCCATCTCCTCTTCG-3'
Primer P4	5'-CATGGGCGACAGGTTCCGAGACC-3'

A clone, GEN-078D0508, obtained by the second 5' RACE was 300 nucleotides long. This clone contained an estimable initiation codon and three preceding in-frame termination codons. From these three overlapping clones, it became clear that the whole coding region comprises 2,636 nucleotides. This gene was named brain-specific nucleosome assembly protein (BNAP) gene.

The BNAP gene contains a 1,518-nucleotide open reading frame shown under SEQ ID NO:20. The amino acid encoded thereby comprises 506 amino acid residues, as shown under SEQ ID NO:19, and the nucleotide sequence of the whole cDNA clone of BNAP is as shown under SEQ ID NO:21.

As shown under SEQ ID NO:21, the 5' noncoding region of said gene was found to be generally rich in GC. Candidate initiation codon sequences were found at nucleotides Nos. 266-268, 287-289 and 329-331. These three sequences all had well conserved sequences in the vicinity of the initiation codons [Kozak, M., J. Biol. Chem., 266, 19867-19870 (1991)].

According to the scanning model, the first ATG (nucleotides Nos. 266-268) of the cDNA clone may be the initiation codon. The termination codon was located at nucleotides Nos. 1784-1786.

The 3' noncoding region was generally rich in AT and two polyadenylation signals (AATAAA) were located at nucleotides Nos. 2606-2611 and 2610-2615, respectively.

The longest open reading frame comprised 1,518 nucleotides coding for 506 amino acid residues and the calculated molecular weight of the BNAP gene product was 57,600 daltons.

Hydrophilic plots indicated that BNAP is very hydrophilic, like other NAPs.

For recombinant BNAP expression and purification and for eliminating the possibility that the BNAP gene sequence might give three chimera clones in the step of 5' RACE, RT-PCR was performed using a sequence comprising nucleotides Nos. 326-356 as a sense primer and a sequence comprising nucleotides Nos. 1758-1786 as an antisense primer.

As a result, a single product of about 1,500 bp was obtained and it was thus confirmed that said sequence is not a chimera but a single transcript.

## (2) Comparison between BNAP and NAPs

The amino acid sequence deduced from BNAP showed 46% identity and 65% similarity to hNRP.

The deduced BNAP gene product had motifs characteristic of the NAPs already reported and of BNAP. In general, half of the C terminus was well conserved in humans and yeasts.

The first motif (domain I) is KGIPDYWLI (corresponding to amino acid residues Nos. 309-317). This was observed also in hNRP (KGIPSFWLT) and in yeast NAP-I (KGIPDFWLT).

The second motif (domain II) is ASFFNFFSPP (corresponding to amino acid residues Nos. 437-446) and this was expressed as DSFFNFFAPP in hNRP and as ESFFNFFSP in yeast NAP-I.

These two motifs were also conserved in the deduced mouse NAP-I and DN38 peptides. Both conserved motifs were each a hydrophilic cluster, and the Cys in position 402 was also found conserved.

Half of the N terminus had no motifs strictly conserved from yeasts to mammalian species, while motifs conserved among mammalian species were found.

For instance, HDLERKYA (corresponding to amino acid residues Nos. 130 to 137) and IINAEYPTEEEECEW (corresponding to amino acid residues Nos. 150-164), which may be associated with mammal-specific functions, were found strictly conserved.

NAPs had acidic stretches, which are believed to be readily capable of binding to histone or other basic proteins. All NAPs had three acidic stretches but the locations thereof were not conserved.

BNAP has no such three acidic stretches but, instead, three repeated sequences (corresponding to amino acid residues Nos. 194-207, 208-221 and 222-235) with a long acidic cluster, inclusive of 41 amino acid residues out of 98 amino acid residues, the consensus sequence being ExxKExPEVKxEEK (each x being a nonconserved, mostly hydrophobic, residue).

Furthermore, it was revealed that the BNAP sequence had several BNAP-specific motifs. Thus, an extremely ser-

ine-rich domain (corresponding to amino acid residues Nos. 24-72) with 33 (67%) of 49 amino acid residues being serine residues was found in the N-terminus portion. On the nucleic acid level, they were reflected as incomplete repetitions of AGC.

Following this serine-rich region, there appeared a basic domain (corresponding to amino acid residues Nos. 71-89) comprising 10 basic amino acid residues among 19 residues.

BNAP is supposed to be localized in the nucleus. Two possible signals localized in the nucleus were observed (NLSs). The first signal was found in the basic domain of BNAP and its sequence YRKRR (corresponding to amino acid residues Nos. 75-79) was similar to NLS (GRKKR) of Tat of HIV-1. The second signal was located in the C terminus and its sequence KKYRK (corresponding to amino acid residues Nos. 502-506) was similar to NLS (KKKRR) of the large T antigen of SV40. The presence of these two presumable NLSs suggested the localization of BNAP in the nucleus. However the possibility that other basic clusters might act as NLSs could not be excluded.

BNAP has several phosphorylation sites and the activity of BNAP may be controlled through phosphorylation thereof.

### (3) Northern blot analysis

Northern blot analysis was performed as described in Example 1 (2). Thus, the clone GEN-078D05TA13 (corresponding to nucleotides Nos. 323 to 1558 in the BNAP gene sequence) was amplified by PCR, the PCR product was purified and labeled with [<sup>32</sup>P]-dCTP (random-primed DNA labeling kit, Boehringer Mannheim), and the expression of BNAP mRNA in normal human tissues was examined using an MTN blot with the labeled product as a probe.

As a result of Northern blot analysis, a 3.0 kb transcript of BNAP was detected (8-hour exposure) in the brain among eight human adult tissues tested, namely heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas and, after longer exposure (24 hours), a dim band of the same size was detected in the heart.

BNAP was found equally expressed in several sites of brain tested whereas, in other tissues, no signal was detected at all even after 72 hours of exposure. hNRP mRNA was found expressed everywhere in the human tissues tested whereas the expression of BNAP mRNA was tissue-specific.

### (4) Radiation hybrid mapping

Chromosomal mapping of the BNAP clone was performed by means of radiation hybrid mapping [Cox, D. R., et al., Science, 250, 245-250 (1990)].

Thus, a total human genome radiation hybrid clone (G3RH) panel was purchased from Research Genetics, Inc., AL, USA and PCR was carried out for chromosomal mapping analysis according to the product manual using two primers, A1 and A2, respectively having the nucleotide sequences shown in Table 5.

Table 5

Primer	Nucleotide sequence
A1 primer	5'-CCTAAAAAGTGTCTAAGTGCCAGTT-3'
A2 primer	5'-TCAGTGAAAGGGAAGGTAGAACAC-3'

The results obtained were analyzed utilizing softwares usable on the Internet [Boehnke, M., et al., Am. J. Hum. Genet., 46, 581-586 (1991)].

As a result, the BNAP gene was found strongly linked to the marker DXS990 (LOD = 1000, cR8000 = -0.00). Since DXS990 is a marker localized on the chromosome Xq21.3-q22, it was established that BNAP is localized to the chromosomal locus Xq21.3-q22 where genes involved in several signs or symptoms of X-chromosome-associated mental retardation are localized.

The nucleosome is not only a fundamental chromosomal structural unit characteristic of eukaryotes but also a gene expression regulating unit. Several results indicate that genes with high transcription activity are sensitive to nuclease treatment, suggesting that the chromosome structure changes with the transcription activity [Elgin, S. C. R., J. Biol. Chem., 263, 19259-19262 (1988)].

NAP-I has been cloned in yeast, mouse and human and is one of the factors capable of promoting nucleosome construction *in vivo*. In a study performed on their sequences, NAPs containing the epitope of the specific antibody 4A8 were detected in human, mouse, frog, *Drosophila* and yeast (*Saccharomyces cerevisiae*) [Ishimi, Y., et al., Eur. J. Biochem., 162, 19-24 (1987)].

In these experiments, NAPs, upon SDS-PAGE analysis, electrophoretically migrated to positions corresponding to

a molecular weight between 50 and 60 kDa, whereas the recombinant BNAP slowly migrated to a position of about 80 kDa. The epitope of 4A8 was shown to be localized in the second, well-conserved, hydrophobic motif. And, it was simultaneously shown that the triplet FNF is important as a part of the epitope [Fujii-Nakata, T., et al., J. Biol. Chem., 267, 20980-20986 (1992)].

BNAP also contained this consensus motif in domain II. The fact that domain II is markedly hydrophobic and the fact that domain II can be recognized by the immune system suggest that it is probably presented on the BNAP surface and is possibly involved in protein-protein interactions.

Domain I, too, may be involved in protein-protein interactions. Considering that these are conserved generally among NAPs, though to a relatively low extent, it is conceivable that they must be essential for nucleosome construction, although the functional meaning of the conserved domains is still unknown.

The hNRP gene is expressed in thyroid gland, stomach, kidney, intestine, leukemia, lung cancer, mammary cancer and so on [Simon, H. U., et al., Biochem. J., 297, 389-397 (1994)]. Like that, NAPs are expressed everywhere and are thought to be playing an important role in fundamental nucleosome formation.

BNAP may be involved in brain-specific nucleosome formation and an insufficiency thereof may cause neurological diseases or mental retardation as a result of deviated functions of neurons.

BNAP was found strongly linked to a marker on the X-chromosome q21.3-q22 where sequences involved in several symptoms of X-chromosome-associated mental retardation are localized. This center-surrounding region of X-chromosome was rich in genes responsible for  $\alpha$ -thalassemia, mental retardation (ATR-X) or some other forms of mental retardation [Gibbons, R. J., et al., Cell, 80, 837-845 (1995)]. Like the analysis of the ATR-X gene which seems to regulate the nucleosome structure, the present inventors suppose that BNAP may be involved in a certain type of X-chromosome-linked mental retardation.

According to this example, the novel BNAP gene is provided and, when said gene is used, it is possible to detect the expression of said gene in various tissues and to produce the BNAP protein by the technology of genetic engineering. Through these, it is possible to study the brain nucleosome formation deeply involved, as mentioned above, in variegated activities essential to cells as well as the functions of cranial nerve cells and to diagnose various neurological diseases or mental retardation in which these are involved and screen out and evaluate drugs for the treatment or prevention of such diseases.

#### Example 7

##### Human skeletal muscle-specific ubiquitin-conjugating enzyme gene (UBE2G gene)

The ubiquitin system is a group of enzymes essential for cellular processes and is conserved from yeast to human. Said system is composed of ubiquitin-activating enzymes (UBAs), ubiquitin-conjugating enzymes (UBCs), ubiquitin protein ligases (UBRs) and 26S proteasome particles.

Ubiquitin is transferred from the above-mentioned UBAs to several UBCs, whereby it is activated. UBCs transfer ubiquitins to target proteins with or without the participation of UBRs. These ubiquitin-conjugated target proteins are said to induce a number of cellular responses, such as protein degradation, protein modification, protein translocation, DNA repair, cell cycle control, transcription control, stress responses, etc. and immunological responses [Jentsch, S., et al., Biochim. Biophys. Acta, 1089, 127-139 (1991); Hershko, A. and Ciechanover, A., Annu. Rev. Biochem., 61, 761-807 (1992); Jentsch, S., Annu. Rev. Genet., 26, 179-207 (1992); Ciechanover, A., Cell 79, 13-21 (1994)].

UBCs are key components of this system and seem to have distinct substrate specificities and modulate different functions. For example, *Saccharomyces cerevisiae* UBC7 is induced by cadmium and involved in resistance to cadmium poisoning [Jungmann, J., et al., Nature, 361, 369-371 (1993)]. Degradation of MAT- $\alpha$ 2 is also executed by UBC7 and UBC6 [Chen, P., et al., Cell, 74, 357-369 (1993)].

The novel gene obtained in this example is UBC7-like gene strongly expressed in human skeletal muscle. In the following, cloning and DNA sequencing thereof are described.

##### (1) Cloning and DNA sequencing of human skeletal muscle-specific ubiquitin-conjugating enzyme gene (UBE2G gene)

Following the same procedure as in Example 1 (1), cDNA clones were arbitrarily selected from a human fetal brain cDNA library and subjected to sequence analysis, and database searches were performed. As a result, a cDNA clone, GEN-423A12, was found to have a significantly high level of homology to the genes coding for ubiquitin-conjugating enzymes (UBCs) in various species.

Since said GEN-423A12 clone was lacking in the 5' side, 5' RACE was performed in the same manner as in Example 2 (2) to obtain an entire coding region.

For said 5' RACE, two primers, P1 and P2, respectively having the nucleotide sequences shown in Table 6 were used.



Table 6

Primer	Nucleotide sequence
P1 primer	5'-TAATGAATTTTCATTTTAGGAGGTCCG-3'
P2 primer	5'-ATCTTTTGGGAAAGTAAGATGAGCC-3'

The 5' RACE product was inserted into pT7Blue(R) T-Vector and clones with an insert proper in size were selected. Four of the 5' RACE clones obtained from two independent PCR reactions contained the same sequence but were different in length.

By sequencing the above clones, the coding sequence and adjacent 5'- and 3'-flanking sequences of the novel gene were determined.

As a result, it was revealed that the novel gene has a total length of 617 nucleotides. This gene was named human skeletal muscle-specific ubiquitin-conjugating enzyme gene (UBE2G gene).

To exclude the conceivable possibility that this sequence was a chimera clone, RT-PCR was performed in the same manner as in Example 6 (1) using the sense primer to amplify said sequence from the human fetal brain cDNA library. As a result, a single PCR product was obtained, whereby it was confirmed that said sequence is not a chimera one.

The UBE2G gene contains an open reading frame of 510 nucleotides, which is shown under SEQ ID NO:23, the amino acid sequence encoded thereby comprises 170 amino acid residues, as shown under SEQ ID NO:22, and the nucleotide sequence of the entire UBE2G cDNA is as shown under SEQ ID NO:24.

As shown under SEQ ID NO:24, the estimable initiation codon was located at nucleotides Nos. 19-21, corresponding to the first ATG triplet of the cDNA clone. Since no preceding in-frame termination codon was found, it was deduced that this clone contains the entire open reading frame on the following grounds.

Thus, (a) the amino acid sequence is highly homologous to *S. cerevisiae* UBC7 and said initiation codon agrees with that of yeast UBC7, supporting said ATG as such. (b) The sequence AGGATGA is similar to the consensus sequence (A/G)CCATGG around the initiation codon [Kozak, M., J. Biol. Chem., 266, 19867-19870 (1991)].

#### (2) Comparison in amino acid sequence between UBE2G and UBCs

Comparison in amino acid sequence between UBE2G and UBCs suggested that the active site cysteine capable of binding to ubiquitin should be the 90th residue cysteine. The peptides encoded by these genes seem to belong to the same family.

#### (3) Northern blot analysis

Northern blot analysis was carried out as described in Example 1 (2). Thus, the entire sequence of UBE2G was amplified by PCR, the PCR product was purified and labeled with [<sup>32</sup>P]-dCTP (random-primed DNA labeling kit, Boehringer Mannheim) and the expression of UBE2G mRNA in normal human tissues using the labeled product as a probe. The membrane used was an MTN blot.

As a result of the Northern blot analysis, 4.4 kb, 2.4 kb and 1.6 kb transcripts could be detected in all 16 human adult tissues, namely heart, brain, placenta, lung, liver, skeletal muscle, kidney, pancreas, spleen, thyroid gland, urinary bladder, testis, ovary, small intestine, large intestine and peripheral blood leukocyte, after 18 hours of exposure. Strong expression of these transcripts was observed in skeletal muscle.

#### (4) Radiation hybrid mapping

Chromosomal mapping of the UBE2G clone was performed by radiation hybrid mapping in the same manner as in Example 6 (4).

The primers C1 and C4 used in PCR for chromosomal mapping analysis respectively correspond to nucleotides Nos. 415-435 and nucleotides Nos. 509-528 in the sequence shown under SEQ ID NO:24 and their nucleotide sequences are as shown below in Table 7.

Table 7

Primer	Nucleotide sequence
C1 primer	5'-GGAGACTCACCTGCTAATGTT-3'
C4 primer	5'-CTCAAAAGCAGTCTCTTGGC-3'

As a result, the UBE2G gene was found linked to the markers D1S446 (LOD = 12.52, cR8000 = 8.60) and D1S235 (LOD = 9.14, cR8000 = 22.46). These markers are localized to the chromosome bands 1q42.13-q42.3.

UBE2G was expressed strongly in skeletal muscle and very weakly in all other tissues examined. All other UBCs are involved in essential cellular functions, such as cell cycle control, and those UBCs are expressed ubiquitously. However, the expression pattern of UBE2G might suggest a muscle-specific role thereof.

While the three transcripts differing in size were detected, attempts failed to identify which corresponds to the cDNA clone. The primary structure of the UBE2G product showed an extreme homology to yeast UBC7. On the other hand, nematode UBC7 showed strong homology to yeast UBC7. It is involved in degradation of the repressor and further confers resistance to cadmium in yeasts. The similarities among these proteins suggest that they belong to the same family.

It is speculated that UBE2G is involved in degradation of muscle-specific proteins and that a defect in said gene could lead to such diseases as muscular dystrophy. Recently, another proteolytic enzyme, calpain 3, was found to be responsible for limb-girdle muscular dystrophy type 2A [Richard, I., et al., Cell, 81, 27-40 (1995)]. At the present, the chromosomal location of UBE2G suggests no significant relationship with any hereditary muscular disease but it is likely that a relation to the gene will be unearthed by linkage analysis in future.

In accordance with this example, the novel UBE2G gene is provided and the use of said gene enables detection of its expression in various tissues and production of the UBE2G protein by the technology of genetic engineering. Through these, it becomes possible to study the degradation of muscle-specific proteins deeply involved in basic activities variegated and essential to cells, as mentioned above, and the functions of skeletal muscle, to diagnose various muscular diseases in which these are involved and further to screen out and evaluate drugs for the treatment and prevention of such diseases.

### Example 8

#### TMP-2 gene

##### (1) TMP-2 gene cloning and DNA sequencing

Following the procedure of Example 1 (1), cDNA clones were arbitrarily selected from a human fetal brain cDNA library and subjected to sequence analysis, and database searches were performed. As a result, a clone (GEN-092E10) having a cDNA sequence highly homologous to a transmembrane protein gene (accession No.: U19878) was found out.

Membrane protein genes have so far been cloned in frog (*Xenopus laevis*) and human. These are considered to be a gene for a transmembrane type protein having a follistatin module and an epidermal growth factor (EGF) domain (accession No.: U19878).

The sequence information of the above protein gene indicated that the GEN-092E10 clone was lacking in the 5' region, so that the  $\lambda$ gt10 cDNA library (human fetal brain 5'-STRETCH PLUS cDNA; Clontech) was screened using the GEN-092E10 clone as a probe, whereby a cDNA clone containing a further 5' upstream region was isolated.

Both strands of this cDNA clone were sequenced, whereby the sequence covering the entire coding region became clear. This gene was named TMP-2 gene.

The TMP-2 gene was found to contain an open reading frame of 1,122 nucleotides, as shown under SEQ ID NO:26, encoding an amino acid sequence of 374 residues, as shown under SEQ ID NO:25. The nucleotide sequence of the entire TMP-2 cDNA clone comprises 1,721 nucleotides, as shown under SEQ ID NO:27.

As shown under SEQ ID NO:27, the 5' noncoding region was generally rich in GC. Several candidates for the initiation codon were found but, according to the scanning model, the 5th ATG of the cDNA clone (bases Nos. 368-370) was estimated as the initiation codon. The termination codon was located at nucleotides Nos. 1490-1492. The polyadenylation signal (AATAAA) was located at nucleotides Nos. 1703-1708. The calculated molecular weight of the TMP-2 gene product was 41,400 daltons.

As mentioned above, the transmembrane genes have a follistatin module and an EGF domain. These motifs were also found conserved in the novel human gene of the present invention.

The TMP-2 gene of the present invention presumably plays an important role in cell proliferation or intercellular communication, since, on the amino acid level, said gene shows homology, across the EGF domain, to TGF- $\alpha$  (transforming growth factor- $\alpha$ ; Derynck, R., et al., *Cell* **38**, 287-297 (1984)), beta-cellulin [Igarashi, K. and Folkman, J., *Science*, **259**, 1604-1607 (1993)], heparin-binding EGF-like growth factor [Higashiyama, S., et al., *Science*, **251**, 936-939 (1991)] and schwannoma-derived growth factor [Kimura, H., et al., *Nature*, **348**, 257-260 (1990)].

## (2) Northern blot analysis

Northern blot analysis was carried out as described in Example 1 (2). Thus, the clone GEN-092E10 was amplified by PCR, the PCR product was purified and labeled with [ $^{32}$ P]-dCTP (random-primed DNA labeling kit, Boehringer Mannheim), and the expression of TMP-2 mRNA in normal human tissues was examined using an MTN blot with the labeled product as a probe.

As a result, high levels of expression were detected in brain and prostate gland. Said TMP-2 gene mRNA was about 2 kb in size.

According to the present invention, the novel human TMP-2 gene is provided and the use of said gene makes it possible to detect the expression of said gene in various tissues or produce the human TMP-2 protein by the technology of genetic engineering and, through these, it becomes possible to study brain tumor and prostatic cancer, which are closely associated with cell proliferation or intercellular communication, as mentioned above, to diagnose these diseases and to screen out and evaluate drugs for the treatment and prevention of such diseases.

## Example 9

### Human NPIK gene

#### (1) Human NPIK gene cloning and DNA sequencing

Following the procedures of Example 1 and Example 2, cDNA clones were arbitrarily selected from a human fetal brain cDNA library and subjected to sequence analysis, and database searches were performed. As a result, two cDNA clones highly homologous to the gene coding for an amino acid sequence conserved in phosphatidylinositol 3 and 4 kinases [Kunz, J., et al., *Cell*, **73**, 585-596 (1993)] were obtained. These were named GEN-428B12c1 and GEN-428B12c2 and the entire sequences of these were determined as in the foregoing examples.

As a result, the GEN-428B12c1 cDNA clone and the GEN-428B12c2 clone were found to have coding sequences differing by 12 amino acid residues at the 5' terminus, the GEN-428B12c1 cDNA clone being longer by 12 amino acid residues.

The GEN-428B12c1 cDNA sequence of the human NPIK gene contained an open reading frame of 2,487 nucleotides, as shown under SEQ ID NO:32, encoding an amino acid sequence comprising 829 amino acid residues, as shown under SEQ ID NO:31. The nucleotide sequence of the full-length cDNA clone comprised 3,324 nucleotides as shown under SEQ ID NO:33.

The estimated initiation codon was located, as shown under SEQ ID NO:33, at nucleotides Nos. 115-117 corresponding to the second ATG triplet of the cDNA clone. The termination codon was located at nucleotides Nos. 2602-2604 and the polyadenylation signal (AATAAA) at Nos. 3305-3310.

On the other hand, the GEN-428B12c2 cDNA sequence of the human NPIK gene contained an open reading frame of 2,451 nucleotides, as shown under SEQ ID NO:29. The amino acid sequence encoded thereby comprised 817 amino acid residues, as shown under SEQ ID NO:28. The nucleotide sequence of the full-length cDNA clone comprised 3,602 nucleotides, as shown under SEQ ID NO:30.

The estimated initiation codon was located, as shown under SEQ ID NO:30, at nucleotides Nos. 429-431 corresponding to the 7th ATG triplet of the cDNA clone. The termination codon was located at nucleotides Nos. 2880-2882 and the polyadenylation signal (AATAAA) at Nos. 3583-3588.

#### (2) Northern blot analysis

Northern blot analysis was carried out as described in Example 1 (2). Thus, the entire sequence of human NPIK was amplified by PCR, the PCR product was purified and labeled with [ $^{32}$ P]-dCTP (random-primed DNA labeling kit, Boehringer Mannheim), and normal human tissues were examined for expression of the human NPIK mRNA using the MTN blot membrane with the labeled product as a probe.

As a result, the expression of the human NPIK gene was observed in 16 various human adult tissues examined and an about 3.8 kb transcript and an about 5 kb one could be detected.

Using primer A having the nucleotide sequence shown below in Table 8 and containing the initiation codon of the GEN-428B12c2 cDNA and primer B shown in table 8 and containing the termination codon, PCR was performed with

Human Fetal Brain Marathon-Ready cDNA (Clontech) as a template, and the nucleotide sequence of the PCR product was determined.

Table 8

Primer	Nucleotide sequence
Primer A	5'-ATGGGAGATACAGTAGTGGAGC-3'
Primer B	5'-TCACATGATGCCGTTGGTGAG-3'

As a result, it was found that the human NPIK mRNA expressed included one lacking in nucleotides Nos. 1060-1104 of the GEN-428B12c1 cDNA sequence (SEQ ID NO:33) (amino acids Nos. 316-330 of the amino acid sequence under SEQ ID NO:31) and one lacking in nucleotides Nos. 1897-1911 of the GEN-428B12c1 cDNA sequence (SEQ ID NO:33) (amino acids Nos. 595-599 of the amino acid sequence under SEQ ID NO:31).

It was further revealed that polymorphism existed in this gene (428B12c1.fasta), as shown below in Table 9, in the region of bases Nos. 1941-1966 of the GEN-428B12c1 cDNA sequence shown under SEQ ID NO:33, whereby a mutant protein was encoded which resulted from the mutation of IQDSCEITT (amino acid residues Nos. 610-618 in the amino acid sequence (SEQ ID NO:31) encoded by GEN-428B12c1) into YKILVISA.

Table 9

```

1930      1940      1950      1959
      TGGATCAAGCCAATACAAGATTCTTGTGAA
      ||||| ||||| ||||| ||||| |||||
TCCATTTGGGAACAGGAGCGAGTGCCCTTTGGATCAAGCC-ATACAAGATTCTTGTG--
1900      1910      1920      1930      1940      1950

1960      1970      1980
ATTACGACTGATAGTGGCATG
||| || ||||| ||||| |||||
ATTTGGGCTGATAGTGGCATGATTGAACCAAGTGGTCAATGCTGTGTCCATCCATCAGGIG
1960      1970      1980      1990      2000      2010

```

### (3) Chromosomal mapping of human NPIK gene by FISH

Chromosomal mapping of the human NPIK gene was carried out by FISH as described in Example 1 (3).

As a result, it was found that the locus of the human NPIK gene is in the chromosomal position 1q21.1-q21.3.

The human NPIK gene, a novel human gene, of the present invention included two cDNAs differing in the 5' region and capable of encoding 829 and 817 amino acid residues, as mentioned above. In view of this and further in view of the findings that the mRNA corresponding to this gene includes two deletable sites and there occurs polymorphism in a specific region corresponding to amino acid residues Nos. 610-618 of the GEN-428B12c1 amino acid sequence (SEQ ID NO:31), whereby a mutant protein is encoded, it is conceivable that human NPIK includes species resulting from a certain number of combinations, namely human NPIK, deletion-containing human NPIK, human NPIK mutant and/or deletion-containing human NPIK mutant.

Recently, several proteins belonging to the family including the above-mentioned PI3 and 4 kinases have protein kinase activity [Dhand, R., et al., EMBO J., 13, 522-533 (1994); Stack, J. H. and Emr, S. D., J. Biol. Chem., 269, 31552-31562 (1994); Hartley, K. O., et al., Cell, 82, 848-856 (1995)].

It was also revealed that a protein belonging to this family is involved in DNA repair [Hartley, K. O., et al., Cell, 82, 849-856 (1995)] and is a causative gene of ataxia [Savitsky, K., et al., Science, 268, 1749-1753 (1995)].

It can be anticipated that the human NPIK gene-encoded protein highly homologous to the family of these PI kinases is a novel enzyme phosphorylating lipids or proteins.

According to this example, the novel human NPIK gene is provided. The use of said gene makes it possible to

detect the expression of said gene in various tissues and manufacture the human NPIK protein by the technology of genetic engineering and, through these, it becomes possible to study lipid- or protein-phosphorylating enzymes such as mentioned above, study DNA repairing, study or diagnose diseases in which these are involved, for example cancer, and screen out and evaluate drugs for the treatment or prevention thereof.

#### (4) Construction of an expression vector for fusion protein

To subclone the coding region for a human NPIK gene (GEN-428B12c2), first of all, two primers, C1 and C2, having the sequences shown below in Table 10 were formed based on the information on the DNA sequences obtained above in (1).

Table 10

Primer	Nucleotide sequence
Primer C1	5'-CTCAGATCTATGGGAGATACAGTAGTGGAGC-3'
Primer C2	5'-TCGAGATCTTCACATGATGCCGTTGGTGAG-3'

Both of the primers C1 and C2 have a BglIII site, and primer C2 is an antisense primer.

Using these two primers, cDNA derived from human fetal brain mRNA was amplified by PCR to provide a product having a length of about 2500 bases. The amplified cDNA was precipitated from ethanol and inserted into pT7BlueT-Vector (product of Novagen) and subcloning was completed. The entire sequence was determined in the same manner as above in Examples. As a result, it was revealed that this gene had polymorphism shown above in Table 9.

The above cDNA was cleaved by BglIII and subjected to agarose gel electrophoresis. The cDNA was then excised from agarose gel and collected using GENECLEAN II KIT (product of Bio 101). The cDNA was inserted into pBlueBacHis2B-Vector (product of Invitrogen) at the BglIII cleavage site and subcloning was completed.

The fusion vector thus obtained had a BglIII cleavage site and was an expression vector for a fusion protein of the contemplated gene product (about 91 kd) and 38 amino acids derived from pBlueBacHis2B-Vector and containing a polyhistidine region and an epitope recognizing Anti-Xpress™ antibody (product of Invitrogen).

#### (5) Transfection into insect cell Sf-9

The human NPIK gene was expressed according to the Baculovirus expression system. Baculovirus is a cyclic double-stranded insect-pathogenic virus and can produce large amounts of inclusion bodies named polyhedrins in the cells of insects. Using Bac-N-Blue™ Transfection Kit utilizing this characteristic of Baculovirus and developed by Invitrogen, the Baculovirus expression was carried out.

Stated more specifically, 4 µg of pBlueBacHis2B containing the region of the human NPIK gene and 1 µg of Bac-N-Blue™ DNA (product of Invitrogen) were co-transfected into Sf-9 cells in the presence of Insectin™ liposomes (product of Invitrogen).

Prior to co-transfection, LacZ gene was incorporated into Bac-N-Blue™ DNA, so that LacZ would be expressed only when homologous recombination took place between the Bac-N-Blue™ DNA and pBlueBacHis2B. Thus when the co-transfected Sf-9 cells were incubated on agar medium, the plaques of the virus expressing the contemplated gene were easily detected as blue plaques.

The blue plaques were excised from each agar and suspended in 400 µl of medium to disperse the virus thereon. The suspension was subjected to centrifugation to give a supernatant containing the virus. Sf-9 cells were infected with the virus again to increase the titre and to obtain a large amount of infective virus solution.

#### (6) Preparation of human NPIK

The expression of the contemplated human NPIK gene was confirmed three days after infection with the virus as follows.

Sf-9 cells were collected and washed with PBS. The cells were boiled with a SDS-PAGE loading buffer for 5 minutes and SDS-PAGE was performed. According to the western blot technique using Anti-Xpress as an antibody, the contemplated protein was detected at the position of its presumed molecular weight. By contrast, in the case of control cells uninfected with the virus, no band corresponding to human NPIK was observed in the same test.

Stated more specifically, three days after the infection of 15 flasks (175-cm<sup>2</sup>, FALCON) of semi-confluent Sf-9 cells, the cells were harvested and washed with PBS, followed by resuspension in a buffer (20 mM Tris/HCl (pH 7.5), 1 mM

EDTA and 1 mM DTT). The suspended cells were lysed by 4 time-sonications for 30 seconds at 4 °C with 30 seconds intervals. The sonicated cells were subjected to centrifugation and the supernatant was collected. The protein in the supernatant was immunoprecipitated using an Anti-Xpress antibody and obtained as a slurry of protein A-Sepharose beads. The slurry was boiled with a SDS-PAGE loading buffer for 5 minutes. SDS-PAGE was performed for identification and quantification of NPIK. The slurry itself was subjected to the following assaying.

#### (7) Confirmation of PI4 Kinase activity

NPIK was expected to have the activity of incorporation phosphoric acid at the 4-position of the inositol ring of phosphatidylinositol (PI), namely, PI4 Kinase activity.

PI4 Kinase activity of NPIK was assayed according to the method of Takenawa, et al. (Yamakawa, A. and Takenawa, T., J. Biol. Chem., 263, 17555-17560 (1988)) as shown below.

First prepared was a mixture of 10 µl of a NPIK slurry (20 mM Tris/HCl (pH 7.5), 1 mM EDTA, 1 mM DTT and 50% protein A beads), 10 µl of a PI solution (prepared by drying 5 mg of a PI-containing commercial chloroform solution in a stream of nitrogen onto a glass tube wall, adding 1 ml of 20 mM Tris/HCl (pH 7.5) buffer and forming micelles by sonication), 10 µl of an applied buffer (210 mM Tris/HCl (pH 7.5), 5 mM EGTA and 100 mM MgCl<sub>2</sub>) and 10 µl of distilled water. Thereto was added 10 µl of an ATP solution (5 µl of 500 µM ATP, 4.9 µl of distilled water and 0.1 µl of γ-<sup>32</sup>P ATP (6000 Ci/mmol, product of NEN Co., Ltd.)). The reaction was started at 30°C and continued for 2, 5, 10 and 20 minutes. The time 10 minutes was set as incubation time because a straight-line increase was observed around 10 minutes in incorporation of phosphoric acid into PI in the assaying process described below.

After completion of the reaction, PI was fractionated by the solvent extraction method and finally re-suspended in chloroform. The suspension was developed by thin layer chromatography (TLC) and the radioactivity of the reaction product at the PI4P-position was assayed using an analyzer (trade name: Bio-Image; product of Fuji Photo Film Co., Ltd.).

Fig. 1 shows the results. Fig. 1 is an analytical diagram of the results of assaying the radioactivity based on TLC as mentioned above. The right lane (2) is the fraction of Sf-9 cell cytoplasm infected with the NPIK-containing virus, whereas the left lane (1) is the fraction of uninfected Sf-9 cell cytoplasm.

Also, predetermined amounts of Triton X-100 and adenosine were added to the above reaction system to check how such addition would affect the PI4 Kinase activity. The PI4 Kinase activity was assayed in the same manner as above.

Fig. 2 shows the results. The results confirmed that NPIK had a typical PI4 Kinase activity accelerated by Triton X-100 and inhibited by adenosine.

#### Example 10

nel-related protein type 1 (NRP1) gene and nel-related protein type 2 (NRP2) gene

##### (1) Cloning and DNA sequencing of NRP1 gene and NRP2 gene

EGF-like repeats have been found in many membrane proteins and in proteins related to growth regulation and differentiation. This motif seems to be involved in protein-protein interactions.

Recently, a gene encoding nel, a novel peptide containing five EGF-like repeats, was cloned from a chick embryonic cDNA library [Matsuhashi, S., et al., Dev. Dynamics, 203, 212-222 (1995)]. This product is considered to be a transmembrane molecule with its EGF-like repeats in the extracellular domain. A 4.5 kb transcript (nel mRNA) is expressed in various tissues at the embryonic stage and exclusively in brain and retina after hatching.

Following the procedure of Example 1 (1), cDNA clones were randomly selected from a human fetal brain cDNA library and subjected to sequence analysis, followed by database searching. As a result, two cDNA clones with significantly high homology to the above-mentioned nel were found and named GEN-073E07 and GEN-093E05, respectively.

Since both clones were lacking in the 5' portion, 5' RACE was performed in the same manner as in Example 2 (2) to obtain the entire coding regions.

As for the primers for 5' RACE, primers having an arbitrary sequence obtained from the cDNA sequences of the above clones were synthesized while the anchor primer attached to a commercial kit was used as such.

5' RACE clones obtained from the PCR were sequenced and the sequences seemingly covering the entire coding regions of both genes were obtained. These genes were respectively named nel-related protein type 1 (NRP1) gene and nel-related protein type 2 (NRP2) gene.

The NRP1 gene contains an open reading frame of 2,430 nucleotides, as shown under SEQ ID NO:35, the amino acid sequence deduced therefrom comprises 810 amino acid residues, as shown under SEQ ID NO:34, and the nucleotide sequence of the entire cDNA clone of said NRP1 gene comprises 2,977 nucleotides, as shown under SEQ ID NO:36.

On the other hand, the NRP2 gene contains an open reading frame of 2,448 nucleotides, as shown under SEQ ID NO:38, the amino acid sequence deduced therefrom comprises 816 amino acid residues, as shown under SEQ ID NO:37, and the nucleotide sequence of the entire cDNA clone of said NRP2 gene comprises 3,198 nucleotides, as shown under SEQ ID NO:39.

Furthermore, the coding regions were amplified by RT-PCR to exclude the possibility that either of the sequences obtained was a chimeric cDNA.

The deduced NRP1 and NRP2 gene products both showed highly hydrophobic N termini capable of functioning as signal peptides for membrane insertion. As compared with chick embryonic nel, they both appeared to have no hydrophobic transmembrane domain. Comparison among NRP1, NRP2 and nel with respect to the deduced peptide sequences revealed that NRP2 has 80% homology on the amino acid level and is more closely related to nel than NRP1 having 50% homology. The cysteine residues in cysteine-rich domains and EGF-like repeats were found completely conserved.

The most remarkable difference between the NRPs and the chick protein was that the human homologs lack the putative transmembrane domain of nel. However, even in this lacking region, the nucleotide sequences of NRPs were very similar to that of nel. Furthermore, the two NRPs each possessed six EGF-like repeats, whereas nel has only five.

Other unique motifs of nel as reported by Matsuhashi et al. [Matsuhashi, S., et al., Dev. Dynamics, 203, 212-222 (1995)] were also found in the NRPs at equivalent positions. Since as mentioned above, it was shown that the two deduced NRP peptides are not transmembrane proteins, the NRPs might be secretory proteins or proteins anchored to membranes as a result of posttranslational modification.

The present inventors speculate that NRPs might function as ligands by stimulating other molecules such as EGF receptors. The present inventors further found that an extra EGF-like repeat could be encoded in nel upon frame shifting of the membrane domain region of nel.

When paralleled and compared with NRP2 and nel, the frame-shifted amino acid sequence showed similarities over the whole range of NRP2 and of nel, suggesting that NRP2 might be a human counterpart of nel. In contrast, NRP1 is considered to be not a human counterpart of nel but a homologous gene.

## (2) Northern blot analysis

Northern blot analysis was carried out as described in Example 1 (2). Thus, the entire sequences of both clones cDNAs were amplified by PCR, the PCR products were purified and labeled with [<sup>32</sup>P]-dCTP (random-primed DNA labeling kit, Boehringer Mannheim) and human normal tissues were examined for NRP mRNA expression using an MTN blot with the labeled products as two probes.

Sixteen adult tissues and four human fetal tissues were examined for the expression pattern of two NRPs.

As a result of the Northern blot analysis, it was found that a 3.5 kb transcript of NRP1 was weakly expressed in fetal and adult brain and kidney. A 3.6 kb transcript of NRP2 was strongly expressed in adult and fetal brain alone, with weak expression thereof in fetal kidney as well.

This suggests that NRPs might play a brain-specific role, for example as signal molecules for growth regulation. In addition, these genes might have a particular function in kidney.

## (3) Chromosomal mapping of NRP1 gene and NRP2 gene by FISH

Chromosomal mapping of the NRP1 gene and NRP2 gene was performed by FISH as described in Example 1 (3).

As a result, it was revealed that the chromosomal locus of the NRP1 gene is localized to 11p15.1-p15.2 and the chromosomal locus of the NRP2 gene to 12q13.11-q13.12.

According to the present invention, the novel human NRP1 gene and NRP2 gene are provided and the use of said genes makes it possible to detect the expression of said genes in various tissues and produce the human NRP1 and NRP2 proteins by the technology of genetic engineering. They can further be used in the study of the brain neurotransmission system, diagnosis of various diseases related to neurotransmission in the brain, and the screening and evaluation of drugs for the treatment and prevention of such diseases. Furthermore, the possibility is suggested that these EGF domain-containing NRPs act as growth factors in brain, hence they may be useful in the diagnosis and treatment of various kinds of intracerebral tumor and effective in nerve regeneration in cases of degenerative nervous diseases.

## Example 11

### GSPT1-related protein (GSPT1-TK) gene

#### (1) GSPT1-TK gene cloning and DNA sequencing

The human GSPT1 gene is one of the human homologous genes of the yeast GST1 gene that encodes the GTP-



binding protein essential for the G1 to S phase transition in the cell cycle. The yeast GST1 gene, first identified as a protein capable of complementing a temperature-sensitive *gst1* (G1-to-S transition) mutant of *Saccharomyces cerevisiae*, was isolated from a yeast genomic library [Kikuchi & Y., Shimatake, H. and Kikuchi, A., EMBO J., 7, 1175-1182 (1988)] and encoded a protein with a target site of cAMP-dependent protein kinases and a GTPase domain.

The human GSPT1 gene was isolated from a KB cell cDNA library by hybridization using the yeast GST1 gene as a probe [Hoshino, S., Miyazawa, H., Enomoto, T., Hanaoka, F., Kikuchi, Y., Kikuchi, A. and Ui, M., EMBO J., 8, 3807-3814 (1989)]. The deduced protein of said GSPT1 gene, like yeast GST1, has a GTP-binding domain and a GTPase activity center, and plays an important role in cell proliferation.

Furthermore, a breakpoint for chromosome re-arrangement has been observed in the GSPT1 gene located in the chromosomal locus 16p13.3 in patients with acute nonlymphocytic leukemia (ANLL) [Ozawa, K., Murakami, Y., Eki, T., Yokoyama, K. Soeda, E., Hoshino, S. Ui, M. and Hanaoka, F., Somatic Cell and Molecular Genet., 18, 189-194 (1992)].

cDNA clones were randomly selected from a human fetal brain cDNA library and subjected to sequence analysis as described in Example 1 (1) and database searching was performed and, as a result, a clone having a 0.3 kb cDNA sequence highly homologous to the above-mentioned GSPT1 gene was found and named GEN-077A09. The GEN-077A09 clone seemed to be lacking in the 5' region, so that 5' RACE was carried out in the same manner as in Example 2 (2) to obtain the entire coding region.

The primers used for the 5' RACE were P1 and P2 primers respectively having the nucleotide sequences shown in Table 11 as designed based on the known cDNA sequence of the above-mentioned cDNA, and the anchor primer used was the one attached to the commercial kit. Thirtyfive cycles of PCR were performed under the following conditions: 94°C for 45 seconds, 58°C for 45 seconds and 72°C for 2 minutes. Finally, elongation reaction was carried out at 72°C for 7 minutes.

Table 11

Primer	Nucleotide sequence
P1 primer	5'-GATTTGTGCTCAATAATCACTATCTGAA-3'
P2 primer	5'-GGTTACTAGGATCACAAAGTATGAATTCTGGAA-3'

Several of the 5' RACE clones obtained from the above PCR were sequenced and the base sequence of that cDNA clone showing overlapping between the 5' RACE clones and the GEN-077A09 clone was determined to thereby reveal the sequence regarded as covering the entire coding region. This was named GSPT1-related protein "GSPT1-TK gene".

The GSPT1-TK gene was found to contain an open reading frame of 1,497 nucleotides, as shown under SEQ ID NO:41. The amino acid sequence deduced therefrom contained 499 amino acid residues, as shown under SEQ ID NO:40.

The nucleotide sequence of the whole cDNA clone of the GSPT1-TK gene was found to comprise 2,057 nucleotides, as shown under SEQ ID NO:42, and the molecular weight was calculated at 55,740 daltons.

The first methionine code (ATG) in the open reading frame had no in-frame termination codon but this ATG was surrounded by a sequence similar to the Kozak consensus sequence for translational initiation. Therefore, it was concluded that this ATG triplet occurring in positions 144-146 of the relevant sequence is the initiation codon.

Furthermore, a polyadenylation signal, AATAAA, was observed 13 nucleotides upstream from the polyadenylation site.

Human GSPT1-TK contains a glutamic acid rich region near the N terminus, and 18 of 20 glutamic acid residues occurring in this region of human GSPT1-TK are conserved and align perfectly with those of the human GSPT1 protein. Several regions (G1, G2, G3, G4 and G5) of GTP-binding proteins that are responsible for guanine nucleotide binding and hydrolysis were found conserved in the GSPT1-TK protein just as in the human GSPT1 protein.

Thus, the DNA sequence of human GSPT1-TK was found 89.4% identical, and the amino acid sequence deduced therefrom 92.4% identical, with the corresponding sequence of human GSPT1 which supposedly plays an important role in the G1 to S phase transition in the cell cycle. Said amino acid sequence showed 50.8% identity with that of yeast GST1.

## (2) Northern blot analysis

Northern blot analysis was carried out as described in Example 1 (2). Thus, the GEN-077A09 cDNA clone was amplified by PCR, the PCR product was purified and labeled with [<sup>32</sup>P]-dCTP (random-primed DNA labeling kit, Boehringer Mannheim), and normal human tissues were examined for the expression of GSPT1-TK mRNA therein using

an MTN blot with the labeled product as a probe.

As a result of the Northern blot analysis, a 2.7 kb major transcript was detected in various tissues. The level of human GSPT1-TK expression seemed highest in brain and in testis.

5 (3) Chromosome mapping of GSPT1-TK gene by FISH

Chromosome mapping of the GSPT1-TK gene was performed by FISH as described in Example 1 (3).

As a result, it was found that the GSPT1-TK gene is localized at the chromosomal locus 19p13.3. In this chromosomal localization site, reciprocal location has been observed very frequently in cases of acute lymphocytic leukemia (ALL) and acute myeloid leukemia (AML). In addition, it is reported that acute non-lymphocytic leukemia (ANLL) is associated with re-arrangements involving the human GSPT1 region [Ozawa, K., Murakami, Y., Eki, T., Yokoyama, K., Soeda, E., Hoshino, S., Ui, M. and Hanaoka, F., Somatic Cell and Molecular Genet., 18, 189-194 (1992)].

In view of the above, it is suggested that this gene is the best candidate gene associated with ALL and AML.

15 In accordance with the present invention, the novel human GSPT1-TK gene is provided and the use of said gene makes it possible to detect the expression of said gene in various tissues and produce the human GSPT1-TK protein by the technology of genetic engineering. These can be used in the studies of cell proliferation, as mentioned above, and further make it possible to diagnose various diseases associated with the chromosomal locus of this gene, for example acute myelocytic leukemia. This is because translocation of this gene may result in decomposition of the GSPT1-TK gene and further some or other fused protein expressed upon said translocation may cause such diseases.

20 Furthermore, it is expected that diagnosis and treatment of said diseases can be made possible by producing antibodies to such fused protein, revealing the intracellular localization of said protein and examining its expression specific to said diseases. Therefore, it is also expected that the use of the gene of the present invention makes it possible to screen out and evaluate drugs for the treatment and prevention of said diseases.

## SEQUENCE LISTING

## (2) INFORMATION FOR SEQ ID NO:1:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 122 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

Met Glu Leu Gly Glu Asp Gly Ser Val Tyr Lys Ser Ile Leu Val Thr  
 1 5 10 15  
 Ser Gln Asp Lys Ala Pro Ser Val Ile Ser Arg Val Leu Lys Lys Asn  
 20 25 30  
 Asn Arg Asp Ser Ala Val Ala Ser Glu Tyr Glu Leu Val Gln Leu Leu  
 35 40 45  
 Pro Gly Glu Arg Glu Leu Thr Ile Pro Ala Ser Ala Asn Val Phe Tyr  
 50 55 60  
 Pro Met Asp Gly Ala Ser His Asp Phe Leu Leu Arg Gln Arg Arg Arg  
 65 70 75 80  
 Ser Ser Thr Ala Thr Pro Gly Val Thr Ser Gly Pro Ser Ala Ser Gly  
 85 90 95  
 Thr Pro Pro Ser Glu Gly Gly Gly Gly Ser Phe Pro Arg Ile Lys Ala  
 100 105 110  
 Thr Gly Arg Lys Ile Ala Arg Ala Leu Phe  
 115 120

## (2) INFORMATION FOR SEQ ID NO:2:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 366 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA(cDNA)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

ATGGAGTTGG GGGAGATGG CAGTGTCTAT AAGAGCATTT TGGTGACAAG CCAGGACAAG 60  
 GCTCCAAGTG TCATCAGTGG TGTCCTTAAG AAAACAATC GTGACTCTGC AGTGGCTTCA 120  
 GAGTATGAGC TGGTACAGCT GCTACCAGGG GAGCGAGAGC TGACTATOCC AGCCTCGGCT 180  
 AATGTATTCT ACCCATGGA TGGAGCTTCA CACGATTTC TOCTGOGGCA GGGGGAAGG 240  
 TOCTCTACTG CTACAACCTGG CGTCAACAGT GGCCCGTCTG CACTAGGAAC TOCTOOGAGT 300  
 GAGGGAGGAG GGGGCTOCTT TOCCAGGATC AAGGCCACAG GGAGGAAGAT TGCAOGGGCA 360  
 CTGTTC 366

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 842 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA(genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vii) IMMEDIATE SOURCE:

- (A) LIBRARY: Human fetal brain cDNA library
- (B) CLONE: GEN-501D08

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 28..393

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

CCACGAGCC GTATCATOOG AGTCCAG ATG GAG TTG GGG GAA GAT GGC AGT 51  
 Met Glu Leu Gly Glu Asp Gly Ser  
 1 5  
 GTC TAT AAG AGC ATT TTG GTG ACA AGC CAG GAC AAG GCT CCA AGT GTC 99

	Val	Tyr	Lys	Ser	Ile	Leu	Val	Thr	Ser	Gln	Asp	Lys	Ala	Pro	Ser	Val	
	10						15					20					
5	ATC	AGT	CGT	GTC	CTT	AAG	AAA	AAC	AAT	CGT	GAC	TCT	GCA	GTG	GCT	TCA	147
	Ile	Ser	Arg	Val	Leu	Lys	Lys	Asn	Asn	Arg	Asp	Ser	Ala	Val	Ala	Ser	
	25					30				35					40		
10	GAG	TAT	GAG	CTG	GTA	CAG	CTG	CTA	CCA	GGG	GAG	CGA	GAG	CTG	ACT	ATC	195
	Glu	Tyr	Glu	Leu	Val	Gln	Leu	Leu	Pro	Gly	Glu	Arg	Glu	Leu	Thr	Ile	
					45					50				55			
15	CCA	GCC	TCG	GCT	AAT	GTA	TTC	TAC	CCC	ATG	GAT	GGA	GCT	TCA	CAC	GAT	243
	Pro	Ala	Ser	Ala	Asn	Val	Phe	Tyr	Pro	Met	Asp	Gly	Ala	Ser	His	Asp	
				60					65					70			
20	TTC	CTC	CTG	CGG	CAG	CGG	CGA	AGG	TCC	TCT	ACT	GCT	ACA	CCT	GGC	GTC	291
	Phe	Leu	Leu	Arg	Gln	Arg	Arg	Arg	Ser	Ser	Thr	Ala	Thr	Pro	Gly	Val	
				75				80						85			
25	ACC	AGT	GGC	CCG	TCT	GCC	TCA	GGA	ACT	OCT	CCG	AGT	GAG	GGA	GGA	GGG	339
	Thr	Ser	Gly	Pro	Ser	Ala	Ser	Gly	Thr	Pro	Pro	Ser	Glu	Gly	Gly	Gly	
		90					95					100					
30	GGC	TOC	TTT	CCC	AGG	ATC	AAG	GCC	ACA	GGG	AGG	AAG	ATT	GCA	CCG	GCA	387
	Gly	Ser	Phe	Pro	Arg	Ile	Lys	Ala	Thr	Gly	Arg	Lys	Ile	Ala	Arg	Ala	
	105					110				115					120		
35	CTG	TTC	TGAGGAGGAA	GOOOCITTTTT	TTACAGAAAGT	CATGGTGTTT	ATAOCAGATG										443
	Leu	Phe															
40	TGGGTAGCCA	TOCTGAATGG	TGGCAATTAT	ATCACATTGA	GACAGAAATT	CAGAAAGGGA											503
	GCCAGCCACC	CTGGGGCAGT	GAAGTGOCAC	TGGTTTACCA	GACAGCTGAG	AAATOCAGOC											563
45	CTGTGGGAAC	TGGTGTCTTA	TAAOCAAAGTT	GGATAOCTGT	GTATAGCTTG	CCAOCTTOCA											623
	TGAGTGCAGC	ACACAGGTAG	TGCTGGAAAA	AOGCATCAGT	TTCTGATTCT	TGGOCATATC											683
50	CTAACATGCA	AGGGCCAAGC	AAAGGCTTCA	AGGCTCTGAG	CCOCAGGGCA	GAGGGGAATG											743
	GCAAAATGTA	GGTOCTGGCA	GGAGCTCTTC	TTCCACTCT	GGGGGTTTCT	ATCACTGTGA											803
55	CAACACTAAG	ATAATAAACC	AAAACACTAC	CTGAATTCT													842

## (2) INFORMATION FOR SEQ ID NO:4:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 193 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Met Glu Leu Glu Leu Tyr Gly Val Asp Asp Lys Phe Tyr Ser Lys Leu  
 1 5 10 15  
 Asp Gln Glu Asp Ala Leu Leu Gly Ser Tyr Pro Val Asp Asp Gly Cys  
 20 25 30  
 Arg Ile His Val Ile Asp His Ser Gly Ala Arg Leu Gly Glu Tyr Glu  
 35 40 45  
 Asp Val Ser Arg Val Glu Lys Tyr Thr Ile Ser Gln Glu Ala Tyr Asp  
 50 55 60  
 Gln Arg Gln Asp Thr Val Arg Ser Phe Leu Lys Arg Ser Lys Leu Gly  
 65 70 75 80  
 Arg Tyr Asn Glu Glu Glu Arg Ala Gln Gln Glu Ala Glu Ala Ala Gln  
 85 90 95  
 Arg Leu Ala Glu Glu Lys Ala Gln Ala Ser Ser Ile Pro Val Gly Ser  
 100 105 110  
 Arg Cys Glu Val Arg Ala Ala Gly Gln Ser Pro Arg Arg Gly Thr Val  
 115 120 125  
 Met Tyr Val Gly Leu Thr Asp Phe Lys Pro Gly Tyr Trp Ile Gly Val  
 130 135 140  
 Arg Tyr Asp Glu Pro Leu Gly Lys Asn Asp Gly Ser Val Asn Gly Lys  
 145 150 155 160  
 Arg Tyr Phe Glu Cys Gln Ala Lys Tyr Gly Ala Phe Val Lys Pro Ala  
 165 170 175  
 Val Val Thr Val Gly Asp Phe Pro Glu Glu Asp Tyr Gly Leu Asp Glu  
 180 185 190  
 Ile

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 579 base pairs

(B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA(cDNA)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

```

10  ATGGAAGTGG AGCTGTATGG AGTTGACGAC AAGTTCTACA GCAAGCTGGA TCAAGAGGAT   60
    GGGCTOCTGG GCTOCTAACC TGTAGATGAC GGCTGCGGCA TOCAAGTCAT TGAACACAGT   120
15  GGGGCGCGGC TTGGTGAGTA TGAGGACGTG TCCCGGGTGG AGAAGTACAC GATCTCACAA   180
    GAAGCTAAG ACCAGAGGCA AGACACGGTC CGCTCTTTTC TGAAGCGCAG CAAGCTGGGC   240
    CGGTACAAAG AGGAGGAGCG GGCTCAGCAG GAGGCGGAGG CCGCGCAGCG OCTGGCGGAG   300
20  GAGAAGGGCC AGGCGAGCTC CATCCCCGTG GGCAGCGGCT GTGAGGTGCG GCGGGGGGA   360
    CAATCCCCTC GCGGGGGCAC CGTCATGTAT GTAGGTCTCA CAGATTTCAA GCGTGGCTAC   420
25  TGGATTGGTG TCGCTATGA TGAGCCACTG GGGAAAAATG ATGGCAGTGT GAATGGGAAA   480
    CGCTACTTGG AATGCCAGGC CAAGTATGGC GCGTTTGTCA AGCCAGCAGT CGTGACGGTG   540
    GGGGACTTCC CGGAGGAGGA CTACGGGTTG GACGAGATA   579
  
```

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1015 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA(genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vii) IMMEDIATE SOURCE:

(A) LIBRARY: Human fetal brain cDNA library  
 (B) CLONE: GEN-080G01

(ix) FEATURE:

(A) NAME/KEY: CDS



(B) LOCATION: 274..852

(x1) SEQUENCE DESCRIPTION: SEQ ID NO:6:

5

TGATTGGTCA GGCACGAGC AGGAGGOGGG CTGATAGOOC AGCAGCAGCA GGGGOGGOGG 60

10

OGGCTGOGGA GGGGTGTGA GGCGGCTGGA OCGOGCTGCA GGCATOOGOG GGOGGGCAA 120

GATGGAGGTG AOGGGGTGT OGGACCAAG GTGACCGTTT TCATCAGCAG CTOOCTCAGC 180

AOCTTOOGCT OOGAGAAGCG ATACAGOOGC AGOCTCAOCC TCGCTGAGTT CAAGTGTAAG 240

15

CTGGAGTTGC TGGTGGGCAG COCTGCTTCC TGC ATG GAA CTG GAG CTG TAT GGA 294  
Met Glu Leu Glu Leu Tyr Gly  
1 5

20

GTT GAC GAC AAG TTC TAC AGC AAG CTG GAT CAA GAG GAT GOG CTC CTG 342  
Val Asp Asp Lys Phe Tyr Ser Lys Leu Asp Gln Glu Asp Ala Leu Leu  
10 15 20

25

GGC TOC TAC OCT GTA GAT GAC GGC TGC OGC ATC CAC GTC ATT GAC CAC 390  
Gly Ser Tyr Pro Val Asp Asp Gly Cys Arg Ile His Val Ile Asp His  
25 30 35

30

AGT GGC GGC OGC CTT GGT GAG TAT GAG GAC GTG TOC OGC GTG GAG AAG 438  
Ser Gly Ala Arg Leu Gly Glu Tyr Glu Asp Val Ser Arg Val Glu Lys  
40 45 50 55TAC ACG ATC TCA CAA GAA GGC TAC GAC CAG AGG CAA GAC ACG GTC OGC 486  
Tyr Thr Ile Ser Gln Glu Ala Tyr Asp Gln Arg Gln Asp Thr Val Arg  
60 65 70

35

TCT TTC CTG AAG OGC AGC AAG CTC GGC OGC TAC AAC GAG GAG GAG OGC 534  
Ser Phe Leu Lys Arg Ser Lys Leu Gly Arg Tyr Asn Glu Glu Glu Arg  
75 80 85

40

GCT CAG CAG GAG GOC GAG GOC GOC CAG OGC CTG GOC GAG GAG AAG GOC 582  
Ala Gln Gln Glu Ala Glu Ala Ala Gln Arg Leu Ala Glu Glu Lys Ala  
90 95 100

45

CAG GOC AGC TOC ATC OOC GTG GGC AGC OGC TGT GAG GTG OGC GOG GOG 630  
Gln Ala Ser Ser Ile Pro Val Gly Ser Arg Cys Glu Val Arg Ala Ala  
105 110 115GGA CAA TOC OCT OGC OGC GGC AOC GTC ATG TAT GTA GGT CTC ACA GAT 678  
Gly Gln Ser Pro Arg Arg Gly Thr Val Met Tyr Val Gly Leu Thr Asp  
120 125 130 135

50

TTC AAG OCT GGC TAC TGG ATT GGT GTC OGC TAT GAT GAG OCA CTG GGG 726

55

Phe Lys Pro Gly Tyr Trp Ile Gly Val Arg Tyr Asp Glu Pro Leu Gly  
 140 145 150

5 AAA AAT GAT GGC AGT GTG AAT GGG AAA CGC TAC TTC GAA TGC CAG GCC 774  
 Lys Asn Asp Gly Ser Val Asn Gly Lys Arg Tyr Phe Glu Cys Gln Ala  
 155 160 165

AAG TAT GGC GGC TTT GTC AAG CCA GCA GTC GTG ACG GTG GGG GAC TTC 822  
 10 Lys Tyr Gly Ala Phe Val Lys Pro Ala Val Val Thr Val Gly Asp Phe  
 170 175 180

CCG GAG GAG GAC TAC GGG TTG GAC GAG ATA TGACACCTAA GGAATTCCCC 872  
 15 Pro Glu Glu Asp Tyr Gly Leu Asp Glu Ile  
 185 190

TGCTTCAGCT OCTAGCTCAG CCACTGACTG CCCCTOCTGT GTGTGCCAT GGCCCTTTTC 932  
 TOCTGACCCC ATTTTAATTT TATTCATTTT TTCTTTGOC ATTGATTTTT GAGACTCATG 992  
 20 CATTAAATTC ACTAGAAOC CAG 1015

## (2) INFORMATION FOR SEQ ID NO:7:

25 (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 128 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

30 (ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

35 Met Thr Glu Ala Asp Val Asn Pro Lys Ala Tyr Pro Leu Ala Asp Ala  
 1 5 10 15

His Leu Thr Lys Lys Leu Leu Asp Leu Val Gln Gln Ser Cys Asn Tyr  
 20 25 30

40 Lys Gln Leu Arg Lys Gly Ala Asn Glu Ala Thr Lys Thr Leu Asn Arg  
 35 40 45

Gly Ile Ser Glu Phe Ile Val Met Ala Ala Asp Ala Glu Pro Leu Glu  
 50 55 60

45 Ile Ile Leu His Leu Pro Leu Leu Cys Glu Asp Lys Asn Val Pro Tyr  
 65 70 75 80

50 Val Phe Val Arg Ser Lys Gln Ala Leu Gly Arg Ala Cys Gly Val Ser

55

85

90

95

Arg Pro Val Ile Ala Cys Ser Val Thr Ile Lys Glu Gly Ser Gln Leu  
 100 105 110

Lys Gln Gln Ile Gln Ser Ile Gln Gln Ser Ile Glu Arg Leu Leu Val  
 115 120 125

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 384 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA(genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

ATGACTGAGG CTGATGTGAA TOCAAAGGOC TATOOOCTTG COGATGOCCA OCTCAOCAAG 60  
 AAGCTACTGG AOCTOGTTCA GCAGTCATGT AACTATAAGC AGCTTOGGAA AGGAGOCAAT 120  
 GAGGOCACCA AAACOOCTCAA CAGGGGCATC TCTGAGTTCA TOGTGATGGC TGCAGAOGOC 180  
 GAGOCACCTGG AGATCATTCT GCAOCTGOOG CTGCTGTGTG AAGACAAGAA TGTGCOOCTAC 240  
 GTGTTTGTGC GCTOCAAGCA GGCOCTGGGG AGAGCOCTGTG GGGTCTOCAG GOCTGTTCATC 300  
 GOCTGTTCCTG TCAOCATCAA AGAAGGCTOG CAGCTGAAAC AGCAGATCCA ATOCATTTCAG 360  
 CAGTGCATTG AAAGGCTCTT AGTC 384

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1493 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA(genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

## (vii) IMMEDIATE SOURCE:

(A) LIBRARY: Human fetal brain cDNA library

(B) CLONE: GEN-025F07

## (ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 95..478

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

	ATCOGTGTCC TTGGGGTGCT GGGCAGCAGA OCGTCCAAAC CGACAAGOGT GGTATCTCTG	60
15	OGGTGTCCGG CAAGAGACTA CCAAGACAGA CGCT ATG ACT GAG GCT GAT GTG Met Thr Glu Ala Asp Val 1 5	112
20	AAT CCA AAG GGC TAT CCC CTT GGC GAT GGC CAC CTC ACC AAG AAG CTA Asn Pro Lys Ala Tyr Pro Leu Ala Asp Ala His Leu Thr Lys Lys Leu 10 15 20	160
25	CTG GAC CTC GTT CAG CAG TCA TGT AAC TAT AAG CAG CTT CGG AAA GGA Leu Asp Leu Val Gln Gln Ser Cys Asn Tyr Lys Gln Leu Arg Lys Gly 25 30 35	208
	GCC AAT GAG GGC ACC AAA ACC CTC AAC AGG GGC ATC TCT GAG TTC ATC Ala Asn Glu Ala Thr Lys Thr Leu Asn Arg Gly Ile Ser Glu Phe Ile 40 45 50	256
30	GTG ATG GCT GCA GAC GGC GAG CCA CTG GAG ATC ATT CTG CAC CTG CCG Val Met Ala Ala Asp Ala Glu Pro Leu Glu Ile Ile Leu His Leu Pro 55 60 65 70	304
35	CTG CTG TGT GAA GAC AAG AAT GTG CCC TAC GTG TTT GTG CGC TOC AAG Leu Leu Cys Glu Asp Lys Asn Val Pro Tyr Val Phe Val Arg Ser Lys 75 80 85	352
40	CAG GGC CTG GGG AGA GGC TGT GGG GTC TOC AGG OCT GTC ATC GGC TGT Gln Ala Leu Gly Arg Ala Cys Gly Val Ser Arg Pro Val Ile Ala Cys 90 95 100	400
45	TCT GTC ACC ATC AAA GAA GGC TCG CAG CTG AAA CAG CAG ATC CAA TOC Ser Val Thr Ile Lys Glu Gly Ser Gln Leu Lys Gln Gln Ile Gln Ser 105 110 115	448
	ATT CAG CAG TOC ATT GAA AGG CTC TTA GTC TAAACCTGTG GOCTCTGACA Ile Gln Gln Ser Ile Glu Arg Leu Leu Val 120 125	498
50	CGTGCTCCCT GGCAGCTTC CCGCTGAGGT TGTGTATCAT ATTATCTGTG TTAGCATGTA	558

5 GTATTTTCAG CTACTCTCTA TTGTTATAAA ATGTAGTACT AAATCTGGTT TCTGGATTTT 618  
 TGATGTGTTT TTGTTCTGTT TTACAGGGTT GCTATCCCCC TTCTTTTCTT CCTTCTCT 678  
 GOCATCTTC ATCTTTTAT CCTTCTTTT TGGAACAAGT GTTCAGAGCA GACAGAAGCA 738  
 GGGTGGTGGC ACOGTTGAAA GGCAGAAAGA GOCAGGAGAA AGCTGATGGA GOCAGGACAG 798  
 10 AGATCTGGTT OCAGCTTTCA GOCAGTAGCT TCTGTGTGTG TGCGGGGTGT GGTGGAATTA 858  
 AACAGCATTG ATTGTGTGTC CCTGTGCTG GCACACAGAA TCATTTCATAC GTGTTCAAGT 918  
 GATCAAGGGG TTTTATTGTC TCTTGGGGGA TTAGGTATCA TTTGGGGAGG AAGCATGTGT 978  
 15 TCTGTGAGGT TGTTOGGCTA TGTOCAAGTG TGTTTTACTA ATGTACCCCT GCTGTTTGCT 1038  
 TTTGGTAATG TGATGTTGAT GTTCTCCCCC TACOCACAAC CATGCCCTTG AGGGTAGCAG 1098  
 20 GGCAGCAGCA TAACAAAGAG ATGTGCTGCA GGACTOOGGA GGCAGCTGG GTGGGTGAGC 1158  
 CATGGGCGAG TTGAOCTGGG TCTTGAAAGA GTGGGGAGTG ACAAGCTCAG AGAGCATGAA 1218  
 CTGATGCTGG CATGAAGGAT TCAGGAAGA TCATGGAGAC CTGGCTGGTA GCTGTAACAG 1278  
 25 AGATGGTGGG GTOCAAGGAA ACAGCTGTC TCTGGTGAAT GGGACTTTCT TTGGTGGACA 1338  
 CTTGGCACCA GCTCTGAGAG CCTTCTTCTT GTGTCTGOC ACCATGTGGG TCAGATGTAC 1398  
 30 TCTCTGTCAC ATGAGGAGAG TGCTAGTTCA TGTGTTCTOC ATTCTTGTGA GCATCTAAT 1458  
 AAATCTGTTT CATTTTGAAA AAAAAAAAAA AAAAA 1493

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 711 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

Met Pro Ala Asp Val Asn Leu Ser Gln Lys Pro Gln Val Leu Gly Pro  
 1 5 10 15

Glu Lys Gln Asp Gly Ser Cys Glu Ala Ser Val Ser Phe Glu Asp Val  
 20 25 30

Thr Val Asp Phe Ser Arg Glu Glu Trp Gln Gln Leu Asp Pro Ala Gln  
 35 40 45  
 5 Arg Cys Leu Tyr Arg Asp Val Met Leu Glu Leu Tyr Ser His Leu Phe  
 50 55 60  
 Ala Val Gly Tyr His Ile Pro Asn Pro Glu Val Ile Phe Arg Met Leu  
 65 70 75 80  
 10 Lys Glu Lys Glu Pro Arg Val Glu Glu Ala Glu Val Ser His Gln Arg  
 85 90 95  
 15 Cys Gln Glu Arg Glu Phe Gly Leu Glu Ile Pro Gln Lys Glu Ile Ser  
 100 105 110  
 Lys Lys Ala Ser Phe Gln Lys Asp Met Val Gly Glu Phe Thr Arg Asp  
 115 120 125  
 20 Gly Ser Trp Cys Ser Ile Leu Glu Glu Leu Arg Leu Asp Ala Asp Arg  
 130 135 140  
 Thr Lys Lys Asp Glu Gln Asn Gln Ile Gln Pro Met Ser His Ser Ala  
 145 150 155 160  
 25 Phe Phe Asn Lys Lys Thr Leu Asn Thr Glu Ser Asn Cys Glu Tyr Lys  
 165 170 175  
 30 Asp Pro Gly Lys Met Ile Arg Thr Arg Pro His Leu Ala Ser Ser Gln  
 180 185 190  
 Lys Gln Pro Gln Lys Cys Cys Leu Phe Thr Glu Ser Leu Lys Leu Asn  
 195 200 205  
 35 Leu Glu Val Asn Gly Gln Asn Glu Ser Asn Asp Thr Glu Gln Leu Asp  
 210 215 220  
 Asp Val Val Gly Ser Gly Gln Leu Phe Ser His Ser Ser Ser Asp Ala  
 225 230 235 240  
 40 Cys Ser Lys Asn Ile His Thr Gly Glu Thr Phe Cys Lys Gly Asn Gln  
 245 250 255  
 45 Cys Arg Lys Val Cys Gly His Lys Gln Ser Leu Lys Gln His Gln Ile  
 260 265 270  
 His Thr Gln Lys Lys Pro Asp Gly Cys Ser Glu Cys Gly Gly Ser Phe  
 275 280 285  
 50 Thr Gln Lys Ser His Leu Phe Ala Gln Gln Arg Ile His Ser Val Gly  
 290 295 300  
 55

Asn Leu His Glu Cys Gly Lys Cys Gly Lys Ala Phe Met Pro Gln Leu  
 305 310 315 320  
 5 Lys Leu Ser Val Tyr Leu Thr Asp His Thr Gly Asp Ile Pro Cys Ile  
 325 330 335  
 Cys Lys Glu Cys Gly Lys Val Phe Ile Gln Arg Ser Glu Leu Leu Thr  
 10 340 345 350  
 His Gln Lys Thr His Thr Arg Lys Lys Pro Tyr Lys Cys His Asp Cys  
 355 360 365  
 15 Gly Lys Ala Phe Phe Gln Met Leu Ser Leu Phe Arg His Gln Arg Thr  
 370 375 380  
 His Ser Arg Glu Lys Leu Tyr Glu Cys Ser Glu Cys Gly Lys Gly Phe  
 385 390 395 400  
 20 Ser Gln Asn Ser Thr Leu Ile Ile His Gln Lys Ile His Thr Gly Glu  
 405 410 415  
 Arg Gln Tyr Ala Cys Ser Glu Cys Gly Lys Ala Phe Thr Gln Lys Ser  
 25 420 425 430  
 Thr Leu Ser Leu His Gln Arg Ile His Ser Gly Gln Lys Ser Tyr Val  
 435 440 445  
 30 Cys Ile Glu Cys Gly Gln Ala Phe Ile Gln Lys Ala His Leu Ile Val  
 450 455 460  
 His Gln Arg Ser His Thr Gly Glu Lys Pro Tyr Gln Cys His Asn Cys  
 465 470 475 480  
 35 Gly Lys Ser Phe Ile Ser Lys Ser Gln Leu Asp Ile His His Arg Ile  
 485 490 495  
 His Thr Gly Glu Lys Pro Tyr Glu Cys Ser Asp Cys Gly Lys Thr Phe  
 40 500 505 510  
 Thr Gln Lys Ser His Leu Asn Ile His Gln Lys Ile His Thr Gly Glu  
 515 520 525  
 45 Arg His His Val Cys Ser Glu Cys Gly Lys Ala Phe Asn Gln Lys Ser  
 530 535 540  
 Ile Leu Ser Met His Gln Arg Ile His Thr Gly Glu Lys Pro Tyr Lys  
 545 550 555 560  
 50 Cys Ser Glu Cys Gly Lys Ala Phe Thr Ser Lys Ser Gln Phe Lys Glu  
 565 570 575  
 55

His Gln Arg Ile His Thr Gly Glu Lys Pro Tyr Val Cys Thr Glu Cys  
580 585 590

5 Gly Lys Ala Phe Asn Gly Arg Ser Asn Phe His Lys His Gln Ile Thr  
595 600 605

His Thr Arg Glu Arg Pro Phe Val Cys Tyr Lys Cys Gly Lys Ala Phe  
610 615 620

10 Val Gln Lys Ser Glu Leu Ile Thr His Gln Arg Thr His Met Gly Glu  
625 630 635 640

15 Lys Pro Tyr Glu Cys Leu Asp Cys Gly Lys Ser Phe Ser Lys Lys Pro  
645 650 655

Gln Leu Lys Val His Gln Arg Ile His Thr Gly Glu Arg Pro Tyr Val  
660 665 670

20 Cys Ser Glu Cys Gly Lys Ala Phe Asn Asn Arg Ser Asn Phe Asn Lys  
675 680 685

His Gln Thr Thr His Thr Arg Asp Lys Ser Tyr Lys Cys Ser Tyr Ser  
690 695 700

25 Val Lys Gly Phe Thr Lys Gln  
705 710

(2) INFORMATION FOR SEQ ID NO:11:

(1) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2133 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(11) MOLECULE TYPE: DNA(genomic)

(x1) SEQUENCE DESCRIPTION: SEQ ID NO:11:

40 ATGCTGCTG ATGGAATTT ATCCAGAAG OCTCAGGTC TGGGTCAGA GAAGCAGGAT 60  
GGATCTTGG AGGCATCAGT GTCAATTTAG GAAGTGACCG TGGACTTCAG CAGGGAGGAG 120  
45 TGGCAGCAAC TGGACCTGC CCAGAGATGC CTGTACGGG ATGTGATGCT GGAGCTCTAT 180  
AGCATCTCT TOGCAGTGGG GTATCACATT OCAACCCAG AGGTCATCTT CAGAATGCTA 240  
50 AAAGAAAAGG AGCCGCTGT GGAGGAGGCT GAAGTCTCAC ATCAGAGGTG TCAAGAAAGG 300



	GAGTTTGGGC TTGAAATOC ACAAAAGGAG ATTTCTAAGA AAGCTTCATT TCAAAAGGAT	360
5	ATGGTAGGTG AGTTCACAAG AGATGGTTCA TGGTGTTOCA TTTTAGAAGA ACTGAGGCTG	420
	GATGCTGAOC GCACAAAGAA AGATGAGCAA AATCAAATTC AAOCCATGAG TCACAGTGCT	480
	TTCTTCAACA AGAAAACATT GAACACAGAA AGCAATTGTG AATATAAGGA COCTGGGAAA	540
10	ATGATTGCA OGAGGOCOC OCTTGCTTCT TCACAGAAAC AAOCTCAGAA ATGTTGCTTA	600
	TTTACAGAAA GTTTGAAGCT GAAOCTAGAA GTGAACGGTC AGAATGAAAG CAATGACACA	660
	GAACAGCTTG ATGACGTTGT TGGGTCTGGT CAGCTATTCA GCCATAGCTC TTCTGATGCC	720
15	TGCAGCAAGA ATATTCATAC AGGAGAGACA TTTTGCAAAG GTAAOCCAGTG TAGAAAAGTC	780
	TGTGGOCATA AACAGTCACT CAAGCAACAT CAAATTCATA CTCAGAAGAA ACCAGATGGA	840
20	TGTTCTGAAT GTGGGGGGAG CTTCACOCAG AAGTCACACC TCTTTGCCCA ACAGAGAATT	900
	CATAGTGTAG GAAAOCTOCA TGAATGTGGC AAATGTGGAA AAGOCCTCAT GOCACAACTA	960
	AAACTCAGTG TATATCTGAC AGATCATACA GGTGATATAC OCTGTATATG CAAGGAATGT	1020
25	GGGAAGGTCT TTATTCAGAG ATCAGAATTG CTTAOGCAOC AGAAAACACA CACTAGAAAG	1080
	AAGOCCTATA AATGOCATGA CTGTGGAAAA GOCCTTTTTOC AGATGTTATC TCTCTTCAGA	1140
30	CATCAGAGAA CTCACAGTAG AGAAAACTC TATGAATGCA GTGAATGTGG CAAAGGCTTC	1200
	TOCAAAACT CAOCCTCAT TATACATCAG AAAATTCATA CTGGTGAGAG ACAGTATGCA	1260
	TGCAGTGAAT GTGGGAAAGC CTTTACOCAG AAGTCAACAC TCAGCTTGCA OCAGAGAATC	1320
35	CACTCAGGGC AGAAGTOCTA TGTGTGTATC GAATGOGGGC AGGOCCTCAT OCAGAAGGCA	1380
	CAOCTGATTG TOCATCAAAG AAGOCACACA GGAGAAAAAC CTTATCAGTG OCACAACTGT	1440
40	GGGAAATOC TCAATTOCAA GTCACAGCTT GATATACATC ATOGAATICA TACAGGGGAG	1500
	AAAOCTTATG AATGCAGTGA CTGTGGAAAA AOCTTCAOC AAAAGTCACA OCTGAATATA	1560
	CAOCAGAAAA TTCATACTGG AGAAAGACAC CATGTATGCA GTGAATGOGG GAAAGOCCTC	1620
45	AACCAGAAGT CAATACTCAG CATGCATCAG AGAATTCACA COGGAGAGAA GOCCTACAAA	1680
	TGCAGTGAAT GTGGGAAAGC CTTCACTTCT AAGTCTCAAT TCAAAGAGCA TCAGCGAATT	1740
50	CACAOGGGTG AGAAAOCTA TGTGTGCACT GAATGTGGGA AGGOCITCAA OGGCAGGTCA	1800
55		

AATTTCATA AACATCAAAT AACTCACACT AGAGAGAGGC CTTTGTCTG TTACAAATGT 1860  
 5 GGAAGGCTT TTGTCCAGAA ATCAGAGTTG ATTAOCCATC AAAGAACTCA CATGGGAGAG 1920  
 AAACCCATG AATGOCITGA CTGTGGGAAA TCGTTCAGTA AGAAACACA ACTCAAGGTG 1980  
 CATCAGCGAA TTCACACGGG AGAAAGAOCT TATGTGTGTT CTGAATGTGG AAAGGOCITC 2040  
 10 AACACAGGT CAAACTTCAA TAAACACCAA ACAACTCATA CCAGAGACAA ATCTTACAAA 2100  
 TGCAGTTATT CTGTGAAAGG CTTTACCAAG CAA 2133

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 3754 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA(genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vii) IMMEDIATE SOURCE:

- (A) LIBRARY: Human fetal brain cDNA library
- (B) CLONE: GEN-076C09

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 346..2478

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

GCTAAGCCTA TGTCGCTTAC TGGACGCTGA AGTGATTGGG AATATTAGCA GTGGGGGTTC 60  
 40 TGTAGGGTCA GGAAGGGGCG GCTGGCTTTG GGGAGTGAT GAGGGGCTTG TTGGGGGTGG 120  
 GGGTGGTGA TAAAGGGATT TCTGGCTGA AGACGAGGCT GTGAGGCTTC TGCAGAACCC 180  
 CCAGGTCAGG CCACATCATT GAGGCTGCAG GATCTCTCTT CATAGCCAG TACGACTCTC 240  
 45 CGCGTGTTC CTGGTTGGAA AATCCAAACA OCTATOCAGC TTCTGGCTOC TGGGAAAAGT 300  
 GGAGTTGTCA GCAAGAGAGA CCGAGAGTAG AAGCCAGAG TGGAG ATG OCT GCT 354  
 Met Pro Ala

5	GAT GTG AAT TTA TOC CAG AAG OCT CAG GTC CTG GGT OCA GAG AAG CAG Asp Val Asn Leu Ser Gln Lys Pro Gln Val Leu Gly Pro Glu Lys Gln 5 10 15	402
10	GAT GGA TCT TGC GAG GCA TCA GTG TCA TTT GAG GAC GTG AOC GTG GAC Asp Gly Ser Cys Glu Ala Ser Val Ser Phe Glu Asp Val Thr Val Asp 20 25 30 35	450
15	TTC AGC AGG GAG GAG TGG CAG CAA CTG GAC OCT GOC CAG AGA TGC CTG Phe Ser Arg Glu Glu Trp Gln Gln Leu Asp Pro Ala Gln Arg Cys Leu 40 45 50	498
20	TAC CGG GAT GTG ATG CTG GAG CTC TAT AGC CAT CTC TTC GCA GTG GGG Tyr Arg Asp Val Met Leu Glu Leu Tyr Ser His Leu Phe Ala Val Gly 55 60 65	546
25	TAT CAC ATT CCC AAC CCA GAG GTC ATC TTC AGA ATG CTA AAA GAA AAG Tyr His Ile Pro Asn Pro Glu Val Ile Phe Arg Met Leu Lys Glu Lys 70 75 80	594
30	GAG CCG CGT GTG GAG GAG GCT GAA GTC TCA CAT CAG AGG TGT CAA GAA Glu Pro Arg Val Glu Glu Ala Glu Val Ser His Gln Arg Cys Gln Glu 85 90 95	642
35	AGG GAG TTT GGG CTT GAA ATC CCA CAA AAG GAG ATT TCT AAG AAA GCT Arg Glu Phe Gly Leu Glu Ile Pro Gln Lys Glu Ile Ser Lys Lys Ala 100 105 110 115	690
40	TCA TTT CAA AAG GAT ATG GTA GGT GAG TTC ACA AGA GAT GGT TCA TGG Ser Phe Gln Lys Asp Met Val Gly Glu Phe Thr Arg Asp Gly Ser Trp 120 125 130	738
45	TGT TOC ATT TTA GAA GAA CTG AGG CTG GAT GCT GAC CGC ACA AAG AAA Cys Ser Ile Leu Glu Glu Leu Arg Leu Asp Ala Asp Arg Thr Lys Lys 135 140 145	786
50	GAT GAG CAA AAT CAA ATT CAA CCC ATG AGT CAC AGT GCT TTC TTC AAC Asp Glu Gln Asn Gln Ile Gln Pro Met Ser His Ser Ala Phe Phe Asn 150 155 160	834
55	AAG AAA ACA TTG AAC ACA GAA AGC AAT TGT GAA TAT AAG GAC OCT GGG Lys Lys Thr Leu Asn Thr Glu Ser Asn Cys Glu Tyr Lys Asp Pro Gly 165 170 175	882
60	AAA ATG ATT CGC ACG AGG CCC CAC CTT GCT TCT TCA CAG AAA CAA OCT Lys Met Ile Arg Thr Arg Pro His Leu Ala Ser Ser Gln Lys Gln Pro 180 185 190 195	930

5	CAG AAA TGT TGC TTA TTT ACA GAA AGT TTG AAG CTG AAC CTA GAA GTG Gln Lys Cys Cys Leu Phe Thr Glu Ser Leu Lys Leu Asn Leu Glu Val 200 205 210	978
10	AAC GGT CAG AAT GAA AGC AAT GAC ACA GAA CAG CTT GAT GAC GTT GTT Asn Gly Gln Asn Glu Ser Asn Asp Thr Glu Gln Leu Asp Asp Val Val 215 220 225	1026
15	GGG TCT GGT CAG CTA TTC AGC CAT AGC TCT TCT GAT GGC TGC AGC AAG Gly Ser Gly Gln Leu Phe Ser His Ser Ser Ser Asp Ala Cys Ser Lys 230 235 240	1074
20	AAT ATT CAT ACA GGA GAG ACA TTT TGC AAA GGT AAC CAG TGT AGA AAA Asn Ile His Thr Gly Glu Thr Phe Cys Lys Gly Asn Gln Cys Arg Lys 245 250 255	1122
25	GTC TGT GGC CAT AAA CAG TCA CTC AAG CAA CAT CAA ATT CAT ACT CAG Val Cys Gly His Lys Gln Ser Leu Lys Gln His Gln Ile His Thr Gln 260 265 270 275	1170
30	AAG AAA OCA GAT GGA TGT TCT GAA TGT GGG GGG AGC TTC AOC CAG AAG Lys Lys Pro Asp Gly Cys Ser Glu Cys Gly Gly Ser Phe Thr Gln Lys 280 285 290	1218
35	TCA CAC CTC TTT GGC CAA CAG AGA ATT CAT AGT GTA GGA AAC CTC CAT Ser His Leu Phe Ala Gln Gln Arg Ile His Ser Val Gly Asn Leu His 295 300 305	1266
40	GAA TGT GGC AAA TGT GGA AAA GGC TTC ATG OCA CAA CTA AAA CTC AGT Glu Cys Gly Lys Cys Gly Lys Ala Phe Met Pro Gln Leu Lys Leu Ser 310 315 320	1314
45	GTA TAT CTG ACA GAT CAT ACA GGT GAT ATA OCC TGT ATA TGC AAG GAA Val Tyr Leu Thr Asp His Thr Gly Asp Ile Pro Cys Ile Cys Lys Glu 325 330 335	1362
50	TGT GGG AAG GTC TTT ATT CAG AGA TCA GAA TTG CTT ACG CAC CAG AAA Cys Gly Lys Val Phe Ile Gln Arg Ser Glu Leu Leu Thr His Gln Lys 340 345 350 355	1410
55	ACA CAC ACT AGA AAG AAG OCC TAT AAA TGC CAT GAC TGT GGA AAA GGC Thr His Thr Arg Lys Lys Pro Tyr Lys Cys His Asp Cys Gly Lys Ala 360 365 370	1458
60	TTT TTC CAG ATG TTA TCT CTC TTC AGA CAT CAG AGA ACT CAC AGT AGA Phe Phe Gln Met Leu Ser Leu Phe Arg His Gln Arg Thr His Ser Arg 375 380 385	1506
65	GAA AAA CTC TAT GAA TGC AGT GAA TGT GGC AAA GGC TTC TOC CAA AAC Glu Lys Leu Tyr Glu Cys Ser Glu Cys Gly Lys Gly Phe Ser Gln Asn	1554

	390	395	400	
5	TCA AOC CTC ATT ATA CAT CAG AAA ATT CAT ACT GGT GAG AGA CAG TAT Ser Thr Leu Ile Ile His Gln Lys Ile His Thr Gly Glu Arg Gln Tyr 405 410 415	1602		
10	GCA TGC AGT GAA TGT GGG AAA GOC TTT AOC CAG AAG TCA ACA CTC AGC Ala Cys Ser Glu Cys Gly Lys Ala Phe Thr Gln Lys Ser Thr Leu Ser 420 425 430 435	1650		
15	TTG CAC CAG AGA ATC CAC TCA GGG CAG AAG TOC TAT GTG TGT ATC GAA Leu His Gln Arg Ile His Ser Gly Gln Lys Ser Tyr Val Cys Ile Glu 440 445 450	1698		
20	TGC GGG CAG GOC TTC ATC CAG AAG GCA CAC CTG ATT GTC CAT CAA AGA Cys Gly Gln Ala Phe Ile Gln Lys Ala His Leu Ile Val His Gln Arg 455 460 465	1746		
25	AGC CAC ACA GGA GAA AAA OCT TAT CAG TGC CAC AAC TGT GGG AAA TOC Ser His Thr Gly Glu Lys Pro Tyr Gln Cys His Asn Cys Gly Lys Ser 470 475 480	1794		
30	TTC ATT TOC AAG TCA CAG CTT GAT ATA CAT CAT OGA ATT CAT ACA GGG Phe Ile Ser Lys Ser Gln Leu Asp Ile His His Arg Ile His Thr Gly 485 490 495	1842		
35	GAG AAA OCT TAT GAA TGC AGT GAC TGT GGA AAA AOC TTC AOC CAA AAG Glu Lys Pro Tyr Glu Cys Ser Asp Cys Gly Lys Thr Phe Thr Gln Lys 500 505 510 515	1890		
40	TCA CAC CTG AAT ATA CAC CAG AAA ATT CAT ACT GGA GAA AGA CAC CAT Ser His Leu Asn Ile His Gln Lys Ile His Thr Gly Glu Arg His His 520 525 530	1938		
45	GTA TGC AGT GAA TGC GGG AAA GOC TTC AAC CAG AAG TCA ATA CTC AGC Val Cys Ser Glu Cys Gly Lys Ala Phe Asn Gln Lys Ser Ile Leu Ser 535 540 545	1986		
50	ATG CAT CAG AGA ATT CAC AOC GGA GAG AAG OCT TAC AAA TGC AGT GAA Met His Gln Arg Ile His Thr Gly Glu Lys Pro Tyr Lys Cys Ser Glu 550 555 560	2034		
55	TGT GGG AAA GOC TTC ACT TCT AAG TCT CAA TTC AAA GAG CAT CAG OGA Cys Gly Lys Ala Phe Thr Ser Lys Ser Gln Phe Lys Glu His Gln Arg 565 570 575	2082		
60	ATT CAC AOC GGT GAG AAA OCC TAT GTG TGC ACT GAA TGT GGG AAG GOC Ile His Thr Gly Glu Lys Pro Tyr Val Cys Thr Glu Cys Gly Lys Ala 580 585 590 595	2130		

5	TTC AAC GGC AGG TCA AAT TTC CAT AAA CAT CAA ATA ACT CAC ACT AGA Phe Asn Gly Arg Ser Asn Phe His Lys His Gln Ile Thr His Thr Arg 600 605 610	2178
10	GAG AGG OCT TTT GTC TGT TAC AAA TGT GGG AAG GCT TTT GTC CAG AAA Glu Arg Pro Phe Val Cys Tyr Lys Cys Gly Lys Ala Phe Val Gln Lys 615 620 625	2226
15	TCA GAG TTG ATT ACC CAT CAA AGA ACT CAC ATG GGA GAG AAA OCC TAT Ser Glu Leu Ile Thr His Gln Arg Thr His Met Gly Glu Lys Pro Tyr 630 635 640	2274
20	GAA TGC CTT GAC TGT GGG AAA TCG TTC AGT AAG AAA OCA CAA CTC AAG Glu Cys Leu Asp Cys Gly Lys Ser Phe Ser Lys Lys Pro Gln Leu Lys 645 650 655	2322
25	GTG CAT CAG CGA ATT CAC ACG GGA GAA AGA OCT TAT GTG TGT TCT GAA Val His Gln Arg Ile His Thr Gly Glu Arg Pro Tyr Val Cys Ser Glu 660 665 670 675	2370
30	TGT GGA AAG GGC TTC AAC AAC AGG TCA AAC TTC AAT AAA CAC CAA ACA Cys Gly Lys Ala Phe Asn Asn Arg Ser Asn Phe Asn Lys His Gln Thr 680 685 690	2418
35	ACT CAT ACC AGA GAC AAA TCT TAC AAA TGC AGT TAT TCT GTG AAA GGC Thr His Thr Arg Asp Lys Ser Tyr Lys Cys Ser Tyr Ser Val Lys Gly 695 700 705	2466
40	TTT ACC AAG CAA TGAATTOCTA GTGCATCAGC ATATTCATAA ATGAAATATA Phe Thr Lys Gln 710	2518
45	CTOOGAGTTT CTTGAAGAAG AGAACATCTT CTCAGAATCA GGTCTAATTA TATGTTATTG AATTCATGCT TCAGAAAAAC TCTAGGGATG CACTGCGTGT GTGAACACAT GATAAAAAAG TCATGCTTTA TTTTAGTGAG GGCAATTACA GAGAAAAGAG TAAGCAGAAA TGTOCTTCTG AGTACTGGOC TCATTAAGGA TTATAAATTT TCTOCCCGGG AAGAAACCT GACTAAGCA TTGAGAAAAG OCTTTCTGTA AAGAATGGTA CAAGACAGGT TGTTACTCGA TTATTTATAG TAAAATATGT GGGAAATTAT ATCAATGATA ACCCTGTTTA TTGTGGGATA TCAATATTTT TAAAGTGOCA ACACAGTCAT GATAGGACAA TATTTTATGT GTGTGTGTGC GOCTTATGTA TATAAGCATA TATATAATAT ATAAGCATAT TATTATATAC AGGTTGAGTA TOOCTTCTOC AAAATGOCTG GGATCAGAAG CATTTTGGAT TTCAGATACT TACAGATTTT GGAATATTTG	2578 2638 2698 2758 2818 2878 2938 2998 3058

CATTATATTT ATTGGTTGAG CATOCTAAT CTGAAAATOC AAGATTAAAT GCTCCAATTA 3118  
 GCATTTTCTT TGAGOGTCAT GTTAGAGTTC AAAAAGTTTC AGATTTTGGG TTTTCAGATT 3178  
 AGGAATACCC AAOCTGTATG TAOGTATATT TCTGTATCTA TGTATGTATA TATATGCATA 3238  
 TGCAGACATA TGTATATGGT CTGGTCAGCA TATGTGTATG TATGCGTATG TATGTATGTA 3298  
 TGTATGCOCT CAGTGCAGTG GGGTTTGCTG CAGAATTCAC TGCATAGCAG GAGATGTAAG 3358  
 CAGATGAGTT ATTTTTTAAAG AGAATCTAAT CTAATTGTTT TTATAAAAAT TATTOOCTAT 3418  
 TGAATATTTA TATAATGAGG TTGTATCAAC AATGATTAAC TOCTTTATTA TACATACACA 3478  
 TGAATGTGCA TTTTGGTAA ATGCATAAAT GAGATTCTAT AATGTTTACT GATCTTTATA 3538  
 TTACAGATTT TCTCTTCTTT TAGGATTAGC TCAGCTTGOC OOOCTTTTC ATCTOCAOCA 3598  
 TCTATAGTGA GOCTCTOCAT AATTAGTGOC AAOCATTAGT CTOGTTTCATA TTTTACACC 3658  
 AGGAGTCAAC AAAGTGTGOC ATTGGOCAAA TATGGOCTOC CAACTGTTTT TTTAAATAA 3718  
 AGTTTTATTG GAACACAAAA AAAAAAAAAA AAAAAA 3754

## (2) INFORMATION FOR SEQ ID NO:13:

## (1) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 389 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

## (11) MOLECULE TYPE: protein

## (x1) SEQUENCE DESCRIPTION: SEQ ID NO:13:

Met Ala Asp Pro Arg Asp Lys Ala Leu Gln Asp Tyr Arg Lys Lys Leu  
 1 5 10 15  
 Leu Glu His Lys Glu Ile Asp Gly Arg Leu Lys Glu Leu Arg Glu Gln  
 20 25 30  
 Leu Lys Glu Leu Thr Lys Gln Tyr Glu Lys Ser Glu Asn Asp Leu Lys  
 35 40 45  
 Ala Leu Gln Ser Val Gly Gln Ile Val Gly Glu Val Leu Lys Gln Leu  
 50 55 60  
 Thr Glu Glu Lys Phe Ile Val Lys Ala Thr Asn Gly Pro Arg Tyr Val

	65		70		75		80
5	Val Gly Cys Arg Arg Gln Leu Asp Lys Ser Lys Leu Lys Pro Gly Thr						
			85		90		95
	Arg Val Ala Leu Asp Met Thr Thr Leu Thr Ile Met Arg Tyr Leu Pro						
		100		105			110
10	Arg Glu Val Asp Pro Leu Val Tyr Asn Met Ser His Glu Asp Pro Gly						
		115		120			125
	Asn Val Ser Tyr Ser Glu Ile Gly Gly Leu Ser Glu Gln Ile Arg Glu						
15		130		135			140
	Leu Arg Glu Val Ile Glu Leu Pro Leu Thr Asn Pro Glu Leu Phe Gln						
		145		150		155	160
20	Arg Val Gly Ile Ile Pro Pro Lys Gly Cys Leu Leu Tyr Gly Pro Pro						
		165		170			175
	Gly Thr Gly Lys Thr Leu Leu Ala Arg Ala Val Ala Ser Gln Leu Asp						
		180		185			190
25	Cys Asn Phe Leu Lys Val Val Ser Ser Ser Ile Val Asp Lys Tyr Ile						
		195		200			205
	Gly Glu Ser Ala Arg Leu Ile Arg Glu Met Phe Asn Tyr Ala Arg Asp						
30		210		215			220
	His Gln Pro Cys Ile Ile Phe Met Asp Glu Ile Asp Ala Ile Gly Gly						
		225		230		235	240
35	Arg Arg Phe Ser Glu Gly Thr Ser Ala Asp Arg Glu Ile Gln Arg Thr						
		245		250			255
	Leu Met Glu Leu Leu Asn Gln Met Asp Gly Phe Asp Thr Leu His Arg						
		260		265			270
40	Val Lys Met Thr Met Ala Thr Asn Arg Pro Asp Thr Leu Asp Pro Ala						
		275		280			285
	Leu Leu Arg Pro Gly Arg Leu Asp Arg Lys Ile His Ile Asp Leu Pro						
45		290		295			300
	Asn Glu Gln Ala Arg Leu Asp Ile Leu Lys Ile His Ala Gly Pro Ile						
		305		310		315	320
50	Thr Lys His Gly Glu Ile Asp Tyr Glu Ala Ile Val Lys Leu Ser Asp						
		325		330			335

55



Gly Phe Asn Gly Ala Asp Leu Arg Asn Val Cys Thr Glu Ala Gly Met  
 340 345 350

Phe Ala Ile Arg Ala Asp His Asp Phe Val Val Gln Glu Asp Phe Met  
 355 360 365

Lys Ala Val Arg Lys Val Ala Asp Ser Lys Lys Leu Glu Ser Lys Leu  
 370 375 380

Asp Tyr Lys Pro Val  
 385

(2) INFORMATION FOR SEQ ID NO:14:

(1) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1167 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(11) MOLECULE TYPE: DNA(genomic)

(x1) SEQUENCE DESCRIPTION: SEQ ID NO:14:

ATGGGGGAOC CTAGAGATAA GGOGCTTCAG GACTACCGCA AGAAGTTGCT TGAACACAAG 60  
 GAGATOGAOC GCOGTCTTAA GGAGTTAAGG GAACAATTAA AAGAACTTAC CAAGCAGTAT 120  
 GAAAAGTCTG AAAATGATCT GAAGGOOCTA CAGAGTGTG GGCAGATCGT GGGTGAAGTG 180  
 CTTAAACAGT TAACTGAAGA AAAATTCAIT GTTAAAGCTA OCAATGGACC AAGATATGTT 240  
 GTGGGTGTG GTGACAGCT TGACAAAAGT AAGCTGAAGC CAGGAACAAG AGTTGCTTTG 300  
 GATATGACTA CACTAACTAT CATGAGATAT TTGCOGAGAG AGGTGGATOC ACTGGTTTAT 360  
 AACATGTCTC ATGAGGAOC TGGGAATGTT TCTTATTCTG AGATTGGAGG GCTATCAGAA 420  
 CAGATCOGGG AATTAAGAGA GGTGATAGAA TTAOCTCTTA CAAACOCAGA GTTATTTCAG 480  
 CGTGTAGGAA TAATAOCTOC AAAAGGCTGT TTGTTATATG GAOCACAGG TACGGGAAAA 540  
 ACACTCTTGG CAOGAGCOGT TGCTAGOCAG CTGGACTGCA ATTTCTTAAA GGTGTATCT 600  
 AGTTCTATTG TAGACAAGTA CATTGGTGAA AGTGCTCGTT TGATCAGAGA AATGTTTAAT 660  
 TATGCTAGAG ATCATCAACC ATGCATCAIT TTTATGGATG AAATAGATGC TATTGGTGGT 720

CGTCGGTTTT CTGAGGGTAC TTCAGCTGAC AGAGAGATTG AGAGAACGTT AATGGAGTTA 780  
 CTGAATCAAA TGGATGGATT TGATACTCTG CATAGAGTTA AAATGAACCAT GGCTACAAAC 840  
 AGACCAGATA CACTGGATOC TGCTTTGCTG CGTCCAGGAA GATTAGATAG AAAAATACAT 900  
 ATTGATTTGC CAAATGAACA AGCAAGATTA GACATACTGA AAATOCATGC AGGTCCCAT 960  
 ACAAAGCATG GTGAAATAGA TTATGAAGCA ATTGTGAAGC TTTCGGATGG CTTTAATGGA 1020  
 GCAGATCTGA GAAATGTTTG TACTGAAGCA GGTATGTTTG CAATTCGTGC TGATCATGAT 1080  
 TTTGTAGTAC AGGAAGACTT CATGAAAGCA GTCAGAAAAG TGGCTGATTC TAAGAAGCTG 1140  
 GAGTCTAAAT TGGACTACAA AACTGTG 1167

## (2) INFORMATION FOR SEQ ID NO:15:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1566 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vii) IMMEDIATE SOURCE:

- (A) LIBRARY: Human fetal brain cDNA library
- (B) CLONE: GEN-331G07

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 17..1183

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

GAGAAGGCTT CTCATC ATG GCG GAC OCT AGA GAT AAG GCG CTT CAG GAC 49  
 Met Ala Asp Pro Arg Asp Lys Ala Leu Gln Asp  
 1 5 10  
 TAC GCG AAG AAG TTG CTT GAA CAC AAG GAG ATC GAC GGC CGT CTT AAG 97  
 Tyr Arg Lys Lys Leu Leu Glu His Lys Glu Ile Asp Gly Arg Leu Lys  
 15 20 25

5	GAG TTA AGG GAA CAA TTA AAA GAA CTT ACC AAG CAG TAT GAA AAG TCT Glu Leu Arg Glu Gln Leu Lys Glu Leu Thr Lys Gln Tyr Glu Lys Ser 30 35 40	145
10	GAA AAT GAT CTG AAG GGC CTA CAG AGT GTT GGG CAG ATC GTG GGT GAA Glu Asn Asp Leu Lys Ala Leu Gln Ser Val Gly Gln Ile Val Gly Glu 45 50 55	193
15	GTG CTT AAA CAG TTA ACT GAA GAA AAA TTC ATT GTT AAA GCT ACC AAT Val Leu Lys Gln Leu Thr Glu Glu Lys Phe Ile Val Lys Ala Thr Asn 60 65 70 75	241
20	GGA CCA AGA TAT GTT GTG GGT TGT CGT CGA CAG CTT GAC AAA AGT AAG Gly Pro Arg Tyr Val Val Gly Cys Arg Arg Gln Leu Asp Lys Ser Lys 80 85 90	289
25	CTG AAG CCA GGA ACA AGA GTT GCT TTG GAT ATG ACT ACA CTA ACT ATC Leu Lys Pro Gly Thr Arg Val Ala Leu Asp Met Thr Thr Leu Thr Ile 95 100 105	337
30	ATG AGA TAT TTG CCG AGA GAG GTG GAT CCA CTG GTT TAT AAC ATG TCT Met Arg Tyr Leu Pro Arg Glu Val Asp Pro Leu Val Tyr Asn Met Ser 110 115 120	385
35	CAT GAG GAC OCT GGG AAT GTT TCT TAT TCT GAG ATT GGA GGG CTA TCA His Glu Asp Pro Gly Asn Val Ser Tyr Ser Glu Ile Gly Gly Leu Ser 125 130 135	433
40	GAA CAG ATC CCG GAA TTA AGA GAG GTG ATA GAA TTA OCT CTT ACA AAC Glu Gln Ile Arg Glu Leu Arg Glu Val Ile Glu Leu Pro Leu Thr Asn 140 145 150 155	481
45	OCA GAG TTA TTT CAG CGT GTA GGA ATA ATA OCT OCA AAA GGC TGT TTG Pro Glu Leu Phe Gln Arg Val Gly Ile Ile Pro Pro Lys Gly Cys Leu 160 165 170	529
50	TTA TAT GGA CCA CCA GGT ACG GGA AAA ACA CTC TTG GCA OGA GGC GTT Leu Tyr Gly Pro Pro Gly Thr Gly Lys Thr Leu Leu Ala Arg Ala Val 175 180 185	577
55	GCT AGC CAG CTG GAC TGC AAT TTC TTA AAG GTT GTA TCT AGT TCT ATT Ala Ser Gln Leu Asp Cys Asn Phe Leu Lys Val Val Ser Ser Ser Ile 190 195 200	625
60	GTA GAC AAG TAC ATT GGT GAA AGT GCT CGT TTG ATC AGA GAA ATG TTT Val Asp Lys Tyr Ile Gly Glu Ser Ala Arg Leu Ile Arg Glu Met Phe 205 210 215	673
65	AAT TAT GCT AGA GAT CAT CAA CCA TGC ATC ATT TTT ATG GAT GAA ATA Asn Tyr Ala Arg Asp His Gln Pro Cys Ile Ile Phe Met Asp Glu Ile	721

	220	225	230	235	
5	GAT GCT ATT GGT GGT CGT CGG TTT TCT GAG GGT ACT TCA GCT GAC AGA Asp Ala Ile Gly Gly Arg Arg Phe Ser Glu Gly Thr Ser Ala Asp Arg 240 245 250				769
10	GAG ATT CAG AGA ACG TTA ATG GAG TTA CTG AAT CAA ATG GAT GGA TTT Glu Ile Gln Arg Thr Leu Met Glu Leu Leu Asn Gln Met Asp Gly Phe 255 260 265				817
15	GAT ACT CTG CAT AGA GTT AAA ATG AOC ATG GCT ACA AAC AGA OCA GAT Asp Thr Leu His Arg Val Lys Met Thr Met Ala Thr Asn Arg Pro Asp 270 275 280				865
	ACA CTG GAT OCT GCT TTG CTG CGT OCA GGA AGA TTA GAT AGA AAA ATA Thr Leu Asp Pro Ala Leu Leu Arg Pro Gly Arg Leu Asp Arg Lys Ile 285 290 295				913
20	CAT ATT GAT TTG OCA AAT GAA CAA GCA AGA TTA GAC ATA CTG AAA ATC His Ile Asp Leu Pro Asn Glu Gln Ala Arg Leu Asp Ile Leu Lys Ile 300 305 310 315				961
25	CAT GCA GGT CCC ATT ACA AAG CAT GGT GAA ATA GAT TAT GAA GCA ATT His Ala Gly Pro Ile Thr Lys His Gly Glu Ile Asp Tyr Glu Ala Ile 320 325 330				1009
30	GTG AAG CTT TCG GAT GGC TTT AAT GGA GCA GAT CTG AGA AAT GTT TGT Val Lys Leu Ser Asp Gly Phe Asn Gly Ala Asp Leu Arg Asn Val Cys 335 340 345				1057
	ACT GAA GCA GGT ATG TTC GCA ATT CGT GCT GAT CAT GAT TTT GTA GTA Thr Glu Ala Gly Met Phe Ala Ile Arg Ala Asp His Asp Phe Val Val 350 355 360				1105
35	CAG GAA GAC TTC ATG AAA GCA GTC AGA AAA GTG GCT GAT TCT AAG AAG Gln Glu Asp Phe Met Lys Ala Val Arg Lys Val Ala Asp Ser Lys Lys 365 370 375				1153
40	CTG GAG TCT AAA TTG GAC TAC AAA OCT GTG TAATTTACTG TAAGATTTTT Leu Glu Ser Lys Leu Asp Tyr Lys Pro Val 380 385				1203
	GATGGCTGCA TGACAGATGT TGGCTTATTG TAAAAATAAA GTTAAAGAAA ATAATGTATG				1263
45	TATTGGCAAT GATGTCATTA AAAGTATATG AATAAAAAATA TGAGTAACAT CATAAAAATT				1323
	AGTAATTCAA CTTTAAAGAT ACAGAAGAAA TTTGTATGTT TGTTAAAGTT GCATTTATTG				1383
50	CAGCAAGTTA CAAAGGGAAA GTGTTGAAGC TTTTCATATT TGCTGOGTGA GCATTTTGTA				1443

55

AAATATTGAA AGTGGTTTGA GATAGTGGTA TAAGAAAGCA TTTCTTATGA CTTATTTTGT 1503  
 ATCATTGTGT TTCTCATCT AAAAAGTTGA ATAAAATCTG TTTGATTCAG TTCTCTAAA 1563  
 AAA 1566

(2) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 223 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

Met Ser Asp Glu Glu Ala Arg Gln Ser Gly Gly Ser Ser Gln Ala Gly  
 1 5 10 15  
 Val Val Thr Val Ser Asp Val Gln Glu Leu Met Arg Arg Lys Glu Glu  
 20 25 30  
 Ile Glu Ala Gln Ile Lys Ala Asn Tyr Asp Val Leu Glu Ser Gln Lys  
 35 40 45  
 Gly Ile Gly Met Asn Glu Pro Leu Val Asp Cys Glu Gly Tyr Pro Arg  
 50 55 60  
 Ser Asp Val Asp Leu Tyr Gln Val Arg Thr Ala Arg His Asn Ile Ile  
 65 70 75 80  
 Cys Leu Gln Asn Asp His Lys Ala Val Met Lys Gln Val Glu Glu Ala  
 85 90 95  
 Leu His Gln Leu His Ala Arg Asp Lys Glu Lys Gln Ala Arg Asp Met  
 100 105 110  
 Ala Glu Ala His Lys Glu Ala Met Ser Arg Lys Leu Gly Gln Ser Glu  
 115 120 125  
 Ser Gln Gly Pro Pro Arg Ala Phe Ala Lys Val Asn Ser Ile Ser Pro  
 130 135 140  
 Gly Ser Pro Ala Ser Ile Ala Gly Leu Gln Val Asp Asp Glu Ile Val  
 145 150 155 160  
 Glu Phe Gly Ser Val Asn Thr Gln Asn Phe Gln Ser Leu His Asn Ile

165                      170                      175  
 5    Gly Ser Val Val Gln His Ser Glu Gly Lys Pro Leu Asn Val Thr Val  
      180                      185                      190  
   Ile Arg Arg Gly Glu Lys His Gln Leu Arg Leu Val Pro Thr Arg Trp  
      195                      200                      205  
 10    Ala Gly Lys Gly Leu Leu Gly Cys Asn Ile Ile Pro Leu Gln Arg  
      210                      215                      220

(2) INFORMATION FOR SEQ ID NO:17:

(1) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 669 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(11) MOLECULE TYPE: DNA(genomic)

(11) SEQUENCE DESCRIPTION: SEQ ID NO:17:

ATGTOOGACG AGGAAGOGAG GCAGAGOGGA GGCTOCTOGC AGGCOGGOGT CGTGACTGTC 60  
 AGCGAAGTOC AGGAGCTGAT GGGGCGCAAG GAGGAGATAG AAGGCGAGAT CAAGGCOAAC 120  
 TATGAAGTGC TGGAAAGCCA AAAAGGCATT GGGATGAAOG AGCOGCTGGT GGACTGTGAG 180  
 GGCTAOCOCOC GGTCAGAOGT GGACCTGTAC CAAGTOOGCA OCGCCAGGCA CAACATCATA 240  
 TGCTGCAGA ATGATCACAA GGCAGTGATG AAGCAGGTGG AGGAGGCOCT GCAOCAGCTG 300  
 CAGCTOGOG ACAAGGAGAA GCAGGCOOGG GACATGGCTG AGGCOOCACAA AGAGGOCATG 360  
 AGCOGCAAAC TGGGTCAGAG TGAGAGOCAG GGOOCTOCAC GGGCTTTOGC CAAAGTGAAC 420  
 AGCATCAGOC COGGCTOOCOC AGCCAGCATC GGGGCTCTGC AAGTGGATGA TGAGATTGTG 480  
 GAGTTGGGCT CTGTGAACAC CCAGAACTTC CAGTCACTGC ATAACATTGG CAGTGTGGTG 540  
 CAGCACAGTG AGGGGAAGOC OCTGAATGTG ACAGTGATOC GCAGGGGGGA AAAACACCAG 600  
 CTTAGACTTG TTCCAACAOG CTGGGCAGGA AAAGGACTGC TGGGCTGCAA CATTATTCT 660  
 CTGCAAAGA 669

## (2) INFORMATION FOR SEQ ID NO:18:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1128 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

## (vii) IMMEDIATE SOURCE:

- (A) LIBRARY: Human fetal brain cDNA library
- (B) CLONE: GEN-163D09

## (ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 125..793

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

25	ACTGTTCTCG CGTTCGGGA CGGCTGTGGT GTTTTGGCGC ATGGGGGAG CGTAGTTACG	60
	GTCGACTGGG GCGTCGTCC TAGCCCGGA GCGGGTCTC TGGAGTCGG GCGGGGGTT	120
30	CAOG ATG TCC GAC GAG GAA GCG AGG CAG AGC GGA GGC TCC TCG CAG GCC	169
	Met Ser Asp Glu Glu Ala Arg Gln Ser Gly Gly Ser Ser Gln Ala	
	1 5 10 15	
35	GGC GTC GTG ACT GTC AGC GAC GTC CAG GAG CTG ATG OGG CGC AAG GAG	217
	Gly Val Val Thr Val Ser Asp Val Gln Glu Leu Met Arg Arg Lys Glu	
	20 25 30	
40	GAG ATA GAA GCG CAG ATC AAG GCC AAC TAT GAC GTG CTG GAA AGC CAA	265
	Glu Ile Glu Ala Gln Ile Lys Ala Asn Tyr Asp Val Leu Glu Ser Gln	
	35 40 45	
45	AAA GGC ATT GGG ATG AAC GAG OGG CTG GTG GAC TGT GAG GGC TAC CCC	313
	Lys Gly Ile Gly Met Asn Glu Pro Leu Val Asp Cys Glu Gly Tyr Pro	
	50 55 60	
50	OGG TCA GAC GTG GAC CTG TAC CAA GTC CGC ACC GCC AGG CAC AAC ATC	361
	Arg Ser Asp Val Asp Leu Tyr Gln Val Arg Thr Ala Arg His Asn Ile	
	65 70 75	
55	ATA TGC CTG CAG AAT GAT CAC AAG GCA GTG ATG AAG CAG GTG GAG GAG	409

	Ile	Cys	Leu	Gln	Asn	Asp	His	Lys	Ala	Val	Met	Lys	Gln	Val	Glu	Glu	
	80					85					90					95	
5	GCC	CTG	CAC	CAG	CTG	CAC	GCT	CGC	GAC	AAG	GAG	AAG	CAG	GCC	CGG	GAC	457
	Ala	Leu	His	Gln	Leu	His	Ala	Arg	Asp	Lys	Glu	Lys	Gln	Ala	Arg	Asp	
				100					105					110			
10	ATG	GCT	GAG	GCC	CAC	AAA	GAG	GCC	ATG	AGC	CGC	AAA	CTG	GGT	CAG	AGT	505
	Met	Ala	Glu	Ala	His	Lys	Glu	Ala	Met	Ser	Arg	Lys	Leu	Gly	Gln	Ser	
				115					120					125			
15	GAG	AGC	CAG	GGC	CGT	CCA	CGG	GCC	TTC	GCC	AAA	GTG	AAC	AGC	ATC	AGC	553
	Glu	Ser	Gln	Gly	Pro	Pro	Arg	Ala	Phe	Ala	Lys	Val	Asn	Ser	Ile	Ser	
				130				135					140				
	CCC	GGC	TCC	CCA	GCC	AGC	ATC	GCG	GGT	CTG	CAA	GTG	GAT	GAT	GAG	ATT	601
	Pro	Gly	Ser	Pro	Ala	Ser	Ile	Ala	Gly	Leu	Gln	Val	Asp	Asp	Glu	Ile	
				145			150					155					
20	GTG	GAG	TTC	GGC	TCT	GTG	AAC	AAC	CAG	AAC	TTC	CAG	TCA	CTG	CAT	AAC	649
	Val	Glu	Phe	Gly	Ser	Val	Asn	Thr	Gln	Asn	Phe	Gln	Ser	Leu	His	Asn	
						165					170					175	
25	ATT	GGC	AGT	GTG	GTG	CAG	CAC	AGT	GAG	GGG	AAG	CCC	CTG	AAT	GTG	ACA	697
	Ile	Gly	Ser	Val	Val	Gln	His	Ser	Glu	Gly	Lys	Pro	Leu	Asn	Val	Thr	
					180					185					190		
30	GTG	ATC	CGC	AGG	GGG	GAA	AAA	CAC	CAG	CTT	AGA	CTT	GTT	CCA	ACA	CGC	745
	Val	Ile	Arg	Arg	Gly	Glu	Lys	His	Gln	Leu	Arg	Leu	Val	Pro	Thr	Arg	
				195					200					205			
35	TGG	GCA	GGA	AAA	GGA	CTG	CTG	GGC	TGC	AAC	ATT	ATT	CGT	CTG	CAA	AGA	793
	Trp	Ala	Gly	Lys	Gly	Leu	Leu	Gly	Cys	Asn	Ile	Ile	Pro	Leu	Gln	Arg	
			210					215					220				
	TGATTGTCCC	TGGGGAACAG	TAACAGGAAA	GCATCTTCCC	TTGCCCTGGA	CTTGGGTCTA											853
	GGGATTTCCA	ACTTGTCTTC	TCTCCCTGAA	GCATAAGGAT	CTGGAAGAGG	CTTGTAACCT											913
40	GAACCTCTGT	GTGGTGGCAG	TACTGTGGOC	CAACAGTGTA	ATCTCCCTGG	ATTAAGGCAT											973
	TCTTAAAAAC	TTAGGCTTGG	CTCTTTTCAC	AAATTAGGOC	AOGGCCCTAA	ATAGGAATTC											1033
	OCTGGATTGT	GGGCAAGTGG	GCGGAAGTTA	TTCTGGCAGG	TACTGGTGTG	ATTATTATTA											1093
45	TTATTTTTTAA	TAAAGAGTTT	TACAGTGCTG	ATATG													1128

(2) INFORMATION FOR SEQ ID NO:19:

50

55



## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 506 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

Met Ala Glu Ala Asp Phe Lys Met Val Ser Glu Pro Val Ala His Gly  
 1 5 10 15  
 Val Ala Glu Glu Glu Met Ala Ser Ser Thr Ser Asp Ser Gly Glu Glu  
 20 25 30  
 Ser Asp Ser Ser Ser Ser Ser Ser Ser Thr Ser Asp Ser Ser Ser Ser  
 35 40 45  
 Ser Ser Thr Ser Gly Ser Ser Ser Gly Ser Gly Ser Ser Ser Ser Ser  
 50 55 60  
 Ser Gly Ser Thr Ser Ser Arg Ser Arg Leu Tyr Arg Lys Lys Arg Val  
 65 70 75 80  
 Pro Glu Pro Ser Arg Arg Ala Arg Arg Ala Pro Leu Gly Thr Asn Phe  
 85 90 95  
 Val Asp Arg Leu Pro Gln Ala Val Arg Asn Arg Val Gln Ala Leu Arg  
 100 105 110  
 Asn Ile Gln Asp Glu Cys Asp Lys Val Asp Thr Leu Phe Leu Lys Ala  
 115 120 125  
 Ile His Asp Leu Glu Arg Lys Tyr Ala Glu Leu Asn Lys Pro Leu Tyr  
 130 135 140  
 Asp Arg Arg Phe Gln Ile Ile Asn Ala Glu Tyr Glu Pro Thr Glu Glu  
 145 150 155 160  
 Glu Cys Glu Trp Asn Ser Glu Asp Glu Glu Phe Ser Ser Asp Glu Glu  
 165 170 175  
 Val Gln Asp Asn Thr Pro Ser Glu Met Pro Pro Leu Glu Gly Glu Glu  
 180 185 190  
 Glu Glu Asn Pro Lys Glu Asn Pro Glu Val Lys Ala Glu Glu Lys Glu  
 195 200 205  
 Val Pro Lys Glu Ile Pro Glu Val Lys Asp Glu Glu Lys Glu Val Ala

	210		215		220
5	Lys Glu Ile Pro Glu Val	Lys Ala Glu Glu	Lys Ala Asp Ser Lys Asp		
	225	230	235	240	
	Cys Met Glu Ala Thr Pro Glu Val	Lys Glu Asp Pro Lys Glu Val Pro			
		245	250	255	
10	Gln Val Lys Ala Asp Asp Lys Glu Gln Pro Lys Ala Thr Glu Ala Lys				
		260	265	270	
	Ala Arg Ala Ala Val Arg Glu Thr His Lys Arg Val Pro Glu Glu Arg				
15		275	280	285	
	Leu Arg Asp Ser Val Asp Leu Lys Arg Ala Arg Lys Gly Lys Pro Lys				
		290	295	300	
20	Arg Glu Asp Pro Lys Gly Ile Pro Asp Tyr Trp Leu Ile Val Leu Lys				
		305	310	315	320
	Asn Val Asp Lys Leu Gly Pro Met Ile Gln Lys Tyr Asp Glu Pro Ile				
		325	330	335	
25	Leu Lys Phe Leu Ser Asp Val Ser Leu Lys Phe Ser Lys Pro Gly Gln				
		340	345	350	
	Pro Val Ser Tyr Thr Phe Glu Phe His Phe Leu Pro Asn Pro Tyr Phe				
30		355	360	365	
	Arg Asn Glu Val Leu Val Lys Thr Tyr Ile Ile Lys Ala Lys Pro Asp				
		370	375	380	
35	His Asn Asp Pro Phe Phe Ser Trp Gly Trp Glu Ile Glu Asp Cys Lys				
		385	390	395	400
	Gly Cys Lys Ile Asp Arg Arg Arg Gly Lys Asp Val Thr Val Thr Thr				
		405	410	415	
40	Thr Gln Ser Arg Thr Thr Ala Thr Gly Glu Ile Glu Ile Gln Pro Arg				
		420	425	430	
	Val Val Pro Asn Ala Ser Phe Phe Asn Phe Phe Ser Pro Pro Glu Ile				
45		435	440	445	
	Pro Met Ile Gly Lys Leu Glu Pro Arg Glu Asp Ala Ile Leu Asp Glu				
		450	455	460	
50	Asp Phe Glu Ile Gly Gln Ile Leu His Asp Asn Val Ile Leu Lys Ser				
		465	470	475	480

55

Lys His Tyr Gly Asn Lys Lys Tyr Arg Lys  
500 505

(1) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 1518 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(x1) SEQUENCE DESCRIPTION: SEQ ID NO:20:

ATGGCAGAAG	CAGATTTTAA	AATGGTCTCG	GAACTGTCTG	CCATGGGGT	TGCGAAGAG	60
GAGATGGCTA	GCTOGACTAG	TGATTCTGGG	GAAGAATCTG	ACAGCAGTAG	CTCTAGCAGC	120
AGCACTAGTG	ACAGCAGCAG	CAGCAGCAGC	ACTAGTGGCA	GCAGCAGGG	CAGCGGCAGC	180
AGCAGCAGCA	GCAGGGGCAG	CACTAGCAGC	CGCAGGCGCT	TGTATAGAAA	GAAGAGGGTA	240
CCTGAGCCTT	CCAGAAGGGC	GCGGGGGGCG	CGTTTGGGAA	CAAATTTGCT	GGATAGGCTG	300
CCTCAGGCAG	TTAGAAATCG	TGTGCAAGCG	CTTAGAAACA	TTCAAGATGA	ATGTGACAAG	360
GTAGATAOC	TGTTCTTAAA	AGCAATTCAT	GATCTTGAAA	GAAAATATGC	TGAACCTAAC	420
AAGCCTCTGT	ATGATAGGCG	GTTTCAAATC	ATCAATGCAG	AATAAGAGCG	TACAGAAGAA	480
GAATGTGAAT	GGAATTCAGA	GGATGAGGAG	TTCAGCAGTG	ATGAGGAGGT	GCAGGATAAC	540
ACCCCTAGTG	AAATGCGCTC	CTTAGAGGGT	GAGGAAGAAG	AAAACCGTAA	AGAAAACCCA	600
GAGGTGAAAG	CTGAAGAGAA	GGAAGTTCTC	AAAGAAATTC	CTGAGGTGAA	GGATGAAGAA	660
AAGGAAGTTG	CTAAAGAAAT	TCCTGAGGTA	AAGGCTGAAG	AAAAAGCAGA	TTCTAAAGAC	720
TGTATGGAGG	CAACCCCTGA	AGTAAAAGAA	GATCTTAAAG	AAGTCCCCCA	GGTAAAGGCA	780
GATGATAAAG	AACAGCGTAA	AGCAACAGAG	GCTAAGGCAA	GGGCTGCAGT	AAGAGAGACT	840
CATAAAAGAG	TTCTGAGGTA	AAGGCTTCGG	GACAGTGTAG	ATCTTAAAAG	AGCTAGGAAG	900

GGAAAGCCTA AAAGAGAAGA COCTAAAGGC ATTCTGACT ATTGGCTGAT TGTTTTAAAG 960  
 5 AATGTTGACA AGCTOGGGCC TATGATTGAG AAGTATGATG AGCCATTCT GAAGTTCTTG 1020  
 TOGGATGTTA GCTGAAGTT CTCAAAOCT GGOCAGCTG TAAGTTACAC CTTTGAATTT 1080  
 CATTTTCTAC OCAACCCATA CTTGAGAAAT GAGGTGCTGG TGAAGACATA TATAATAAAG 1140  
 10 GCAAAACCAG ATCACAATGA TCOCTTCTTT TCTTGGGGAT GGGAAATTGA AGATTGCAAA 1200  
 GGCTGCAAGA TAGACGGAG AAGAGGAAAA GATGTTACTG TGACAACTAC CCAGAGTOGC 1260  
 ACAACTGCTA CTGGAGAAAT TGAAATOCAG CCAAGAGTGG TTCTAATGC ATCATTCTTC 1320  
 15 AACTTCTTTA GTCTCTCTGA GATTCTATG ATTGGGAAGC TGGAACCAAG AGAAGATGCT 1380  
 ATCTGGATG AGGACTTTGA AATTGGGCAG ATTTTACATG ATAATGTCAT CCTGAAATCA 1440  
 20 ATCTATTACT ATACTGGAGA AGTCAATGGT AACTACTATC AATTGGCAA ACATTATGGA 1500  
 AACAGAAAT ACAGAAAA 1518

25 (2) INFORMATION FOR SEQ ID NO:21:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 2636 base pairs  
 (B) TYPE: nucleic acid  
 30 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear  
  
 (ii) MOLECULE TYPE: DNA(genomic)  
  
 (iii) HYPOTHETICAL: NO  
 35 (iv) ANTI-SENSE: NO  
  
 (vii) IMMEDIATE SOURCE:  
 (A) LIBRARY: Human fetal brain cDNA library  
 40 (B) CLONE: GEN-078D05  
  
 (ix) FEATURE:  
 (A) NAME/KEY: CDS  
 (B) LOCATION: 266..1783  
 45 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

GATTGGGCTG CGGTACATCT CGGCACTCTA GCTGCAGGCG GGAGAGGCT TGGGGCCACC 60  
 50 GCTGTGGGCG AAGCTOCAC TGGGCTGCC AACTCAGGCG CGGCTCTGC ATCCAGCT 120

55

	OCAGCTOOGC TCTGCGOOGC TGCTGOCATC GCGCTGOCA OCTOOGCAGC CCGGGCCTOC	180
5	GOOGCOGCOA CCAAGCATC CGTGAGTCAT TTTCTGCOCA TCTCTGGTGG CGGGTCTOC	240
	CTGGTAGAGT TTGTAGGCTT GCAAG ATG GCA GAA GCA GAT TTT AAA ATG GTC Met Ala Glu Ala Asp Phe Lys Met Val 1 5	292
10	TOG GAA OCT GTC GOC CAT GGG GTT GOC GAA GAG GAG ATG GCT AGC TOG Ser Glu Pro Val Ala His Gly Val Ala Glu Glu Glu Met Ala Ser Ser 10 15 20 25	340
15	ACT AGT GAT TCT GGG GAA GAA TCT GAC AGC AGT AGC TCT AGC AGC AGC Thr Ser Asp Ser Gly Glu Glu Ser Asp Ser Ser Ser Ser Ser Ser Ser 30 35 40	388
20	ACT AGT GAC AGC AGC AGC AGC AGC AGC ACT AGT GGC AGC AGC AGC GGC Thr Ser Asp Ser Ser Ser Ser Ser Ser Thr Ser Gly Ser Ser Ser Gly 45 50 55	436
	AGC GGC AGC AGC AGC AGC AGC AGC GGC AGC ACT AGC AGC CGC AGC CGC Ser Gly Ser Ser Ser Ser Ser Ser Gly Ser Thr Ser Ser Arg Ser Arg 60 65 70	484
25	TTG TAT AGA AAG AAG AGG GTA OCT GAG OCT TOC AGA AGG GCG CGG CGG Leu Tyr Arg Lys Lys Arg Val Pro Glu Pro Ser Arg Arg Ala Arg Arg 75 80 85	532
30	GOC OOG TTG GGA ACA AAT TTC GTG GAT AGG CTG OCT CAG GCA GTT AGA Ala Pro Leu Gly Thr Asn Phe Val Asp Arg Leu Pro Gln Ala Val Arg 90 95 100 105	580
35	AAT CGT GTG CAA GCG CTT AGA AAC ATT CAA GAT GAA TGT GAC AAG GTA Asn Arg Val Gln Ala Leu Arg Asn Ile Gln Asp Glu Cys Asp Lys Val 110 115 120	628
40	GAT AOC CTG TTC TTA AAA GCA ATT CAT GAT CTT GAA AGA AAA TAT GCT Asp Thr Leu Phe Leu Lys Ala Ile His Asp Leu Glu Arg Lys Tyr Ala 125 130 135	676
	GAA CTC AAC AAG OCT CTG TAT GAT AGG OGG TTT CAA ATC ATC AAT GCA Glu Leu Asn Lys Pro Leu Tyr Asp Arg Arg Phe Gln Ile Ile Asn Ala 140 145 150	724
45	GAA TAC GAG OCT ACA GAA GAA GAA TGT GAA TGG AAT TCA GAG GAT GAG Glu Tyr Glu Pro Thr Glu Glu Glu Cys Glu Trp Asn Ser Glu Asp Glu 155 160 165	772
50	GAG TTC AGC AGT GAT GAG GAG GTG CAG GAT AAC AOC OCT AGT GAA ATG Glu Phe Ser Ser Asp Glu Glu Val Gln Asp Asn Thr Pro Ser Glu Met	820

55

	170				175					180					185		
5	OCT	CCC	TTA	GAG	GGT	GAG	GAA	GAA	GAA	AAC	OCT	AAA	GAA	AAC	CCA	GAG	868
	Pro	Pro	Leu	Glu	Gly	Glu	Glu	Glu	Glu	Asn	Pro	Lys	Glu	Asn	Pro	Glu	
				190						195					200		
10	GTG	AAA	GCT	GAA	GAG	AAG	GAA	GTT	OCT	AAA	GAA	ATT	OCT	GAG	GTG	AAG	916
	Val	Lys	Ala	Glu	Glu	Lys	Glu	Val	Pro	Lys	Glu	Ile	Pro	Glu	Val	Lys	
			205					210					215				
15	GAT	GAA	GAA	AAG	GAA	GTT	GCT	AAA	GAA	ATT	OCT	GAG	GTA	AAG	GCT	GAA	964
	Asp	Glu	Glu	Lys	Glu	Val	Ala	Lys	Glu	Ile	Pro	Glu	Val	Lys	Ala	Glu	
			220				225					230					
20	GAA	AAA	GCA	GAT	TCT	AAA	GAC	TGT	ATG	GAG	GCA	AOC	OCT	GAA	GTA	AAA	1012
	Glu	Lys	Ala	Asp	Ser	Lys	Asp	Cys	Met	Glu	Ala	Thr	Pro	Glu	Val	Lys	
		235					240					245					
25	GAA	GAT	OCT	AAA	GAA	GTC	CCC	CAG	GTA	AAG	GCA	GAT	GAT	AAA	GAA	CAG	1060
	Glu	Asp	Pro	Lys	Glu	Val	Pro	Gln	Val	Lys	Ala	Asp	Asp	Lys	Glu	Gln	
	250					255					260					265	
30	OCT	AAA	GCA	ACA	GAG	GCT	AAG	GCA	AGG	GCT	GCA	GTA	AGA	GAG	ACT	CAT	1108
	Pro	Lys	Ala	Thr	Glu	Ala	Lys	Ala	Arg	Ala	Ala	Val	Arg	Glu	Thr	His	
				270						275					280		
35	AAA	AGA	GTT	OCT	GAG	GAA	AGG	CTT	CGG	GAC	AGT	GTA	GAT	CTT	AAA	AGA	1156
	Lys	Arg	Val	Pro	Glu	Glu	Arg	Leu	Arg	Asp	Ser	Val	Asp	Leu	Lys	Arg	
			285						290					295			
40	GCT	AGG	AAG	GGA	AAG	OCT	AAA	AGA	GAA	GAC	OCT	AAA	GGC	ATT	OCT	GAC	1204
	Ala	Arg	Lys	Gly	Lys	Pro	Lys	Arg	Glu	Asp	Pro	Lys	Gly	Ile	Pro	Asp	
			300				305						310				
45	TAT	TGG	CTG	ATT	GTT	TTA	AAG	AAT	GTT	GAC	AAG	CTC	GGG	OCT	ATG	ATT	1252
	Tyr	Trp	Leu	Ile	Val	Leu	Lys	Asn	Val	Asp	Lys	Leu	Gly	Pro	Met	Ile	
		315					320					325					
50	CAG	AAG	TAT	GAT	GAG	CCC	ATT	CTG	AAG	TTC	TTG	TOG	GAT	GTT	AGC	CTG	1300
	Gln	Lys	Tyr	Asp	Glu	Pro	Ile	Leu	Lys	Phe	Leu	Ser	Asp	Val	Ser	Leu	
	330					335					340					345	
55	AAG	TTC	TCA	AAA	OCT	GGC	CAG	OCT	GTA	AGT	TAC	AOC	TTT	GAA	TTT	CAT	1348
	Lys	Phe	Ser	Lys	Pro	Gly	Gln	Pro	Val	Ser	Tyr	Thr	Phe	Glu	Phe	His	
				350					355					360			
60	TTT	CTA	CCC	AAC	CCA	TAC	TTC	AGA	AAT	GAG	GTG	CTG	GTG	AAG	ACA	TAT	1396
	Phe	Leu	Pro	Asn	Pro	Tyr	Phe	Arg	Asn	Glu	Val	Leu	Val	Lys	Thr	Tyr	
			365					370						375			

5	ATA ATA AAG GCA AAA OCA GAT CAC AAT GAT CCC TTC TTT TCT TGG GGA Ile Ile Lys Ala Lys Pro Asp His Asn Asp Pro Phe Phe Ser Trp Gly 380 385 390	1444
10	TGG GAA ATT GAA GAT TGC AAA GGC TGC AAG ATA GAC OGG AGA AGA GGA Trp Glu Ile Glu Asp Cys Lys Gly Cys Lys Ile Asp Arg Arg Arg Gly 395 400 405	1492
15	AAA GAT GTT ACT GTG ACA ACT ACC CAG AGT OGC ACA ACT GCT ACT GGA Lys Asp Val Thr Val Thr Thr Thr Gln Ser Arg Thr Thr Ala Thr Gly 410 415 420 425	1540
20	GAA ATT GAA ATC CAG OCA AGA GTG GTT OCT AAT GCA TCA TTC TTC AAC Glu Ile Glu Ile Gln Pro Arg Val Val Pro Asn Ala Ser Phe Phe Asn 430 435 440	1588
25	TTC TTT AGT OCT OCT GAG ATT OCT ATG ATT GGG AAG CTG GAA OCA OGA Phe Phe Ser Pro Pro Glu Ile Pro Met Ile Gly Lys Leu Glu Pro Arg 445 450 455	1636
30	GAA GAT GCT ATC CTG GAT GAG GAC TTT GAA ATT GGG CAG ATT TTA CAT Glu Asp Ala Ile Leu Asp Glu Asp Phe Glu Ile Gly Gln Ile Leu His 460 465 470	1684
35	GAT AAT GTC ATC CTG AAA TCA ATC TAT TAC TAT ACT GGA GAA GTC AAT Asp Asn Val Ile Leu Lys Ser Ile Tyr Tyr Tyr Thr Gly Glu Val Asn 475 480 485	1732
40	GGT AOC TAC TAT CAA TTT GGC AAA CAT TAT GGA AAC AAG AAA TAC AGA Gly Thr Tyr Tyr Gln Phe Gly Lys His Tyr Gly Asn Lys Lys Tyr Arg 490 495 500 505	1780
45	AAA TAAGTCAATC TGAAAGATTT TTCAAGAATC TTAAAATCTC AAGAAGTGAA Lys	1833
50	GCAGATTTCAT ACAGCCTTGA AAAAAGTAAA AOCCTGACCT GTAACCTGAA CACTATTATT OCTTATAGTC AAGTTTTTGT GGTTCCTTGG TAGTCTATAT TTTAAAAATA GTCTAAAAA GTGCTAAGT GCCAGTTTAT TCTATCTAGG CTGTTGTAGT ATAATATTCT TCAAAATATG TAAGCTGTIG TCAATTATCT AAAGCATGTT AGTTTGGTGC TACACAGTGT TGATTTTTGT GATGTCTTTT GGTTCATGTTT CTGTTAGACT GTAGCTGTGA AACTGTCAGA ATTGTTAACT GAAACAAATA TTTGCTTGAA AAAAAAGTT CATGAAGTAC CAATGCAAGT GTTTTATTTT TTTTCTTTTT TOCAGCCCAT AAGACTAAGG GTTTAAATCT GCTTGCACTA GCTGTGCCIT CATTAGTTTG CTATAGAAAT OCAGTACTTA TAGTAAATAA AACAGTGTAT TTTGAAGTTT	1893 1953 2013 2073 2133 2193 2253 2313

GACTGCTTGA AAAAGATTAG CATACTCTA ATGTGAAAAG ACCACATTTG ATTCAACTGA 2373  
 5 GAOCTTGTGT ATGTGACATA TAGTGGCCTA TAAATTTAAT CATAATGATG TTATTGTTTA 2433  
 OCACTGAGGT GTTAATATAA CATAGTATTT TTGAAAAAGT TTCTTCATCT TATATTGTTG 2493  
 AATTGTAAAC TAAAGATAAC GTGTTTTTCTT TGTATTGTTG TCTACCTTCC CTTTCACTGA 2553  
 10 AAATGATCAC TTCATTGAT ACTGTTTTTC ATGTCTTGT ATTGCAAOCT AAAATAAATA 2613  
 AATATTAAAG TGTGTTATAC TAT 2636

(2) INFORMATION FOR SEQ ID NO:22:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 170 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

Met Thr Glu Leu Gln Ser Ala Leu Leu Leu Arg Arg Gln Leu Ala Glu  
 1 5 10 15  
 Leu Asn Lys Asn Pro Val Glu Gly Phe Ser Ala Gly Leu Ile Asp Asp  
 20 25 30  
 Asn Asp Leu Tyr Arg Trp Glu Val Leu Ile Ile Gly Pro Pro Asp Thr  
 35 40 45  
 Leu Tyr Glu Gly Gly Val Phe Lys Ala His Leu Thr Phe Pro Lys Asp  
 50 55 60  
 Tyr Pro Leu Arg Pro Pro Lys Met Lys Phe Ile Thr Glu Ile Trp His  
 65 70 75 80  
 Pro Asn Val Asp Lys Asn Gly Asp Val Cys Ile Ser Ile Leu His Glu  
 85 90 95  
 Pro Gly Glu Asp Lys Tyr Gly Tyr Glu Lys Pro Glu Glu Arg Trp Leu  
 100 105 110  
 Pro Ile His Thr Val Glu Thr Ile Met Ile Ser Val Ile Ser Met Leu  
 115 120 125  
 Ala Asp Pro Asn Gly Asp Ser Pro Ala Asn Val Asp Ala Ala Lys Glu



130

135

140

5 Trp Arg Glu Asp Arg Asn Gly Glu Phe Lys Arg Lys Val Ala Arg Cys  
145 150 155 160

Val Arg Lys Ser Gln Glu Thr Ala Phe Glu  
165 170

10

## (2) INFORMATION FOR SEQ ID NO:23:

## (i) SEQUENCE CHARACTERISTICS:

15

- (A) LENGTH: 510 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

20

## (ii) MOLECULE TYPE: DNA(genomic)

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

25

ATGAOGGAGC TGCAGTOGGC ACTGCTACTG CGAAGACAGC TGGCAGAACT CAACAAAAAT 60

OCAGTGGGAAG GCTTTTCTGC AGGTTTAAATA GATGACAATG ATCTCTACCG ATGGGAAGTC 120

CTTATTATTG GCOCTOCAGA TACACTTTAT GAAGGTGGTG TTTTAAAGGC TCATCTTACT 180

30

TTOCCAAAAG ATTATOOOCT COGACCTOCT AAAATGAAAT TCATTACAGA AATCTGGCAC 240

OCAAATGTTG ATAAAAATGG TGATGTGTGC ATTTCTATTTC TTCATGAGCC TGGGGAAGAT 300

35

AAGTATGGTT ATGAAAAGOC AGAGGAAOGC TGGCTOOCTA TOCACACTGT GGAAACCATC 360

ATGATTAGTG TCATTTCTAT GCTGGCAGAC OCTAATGGAG ACTCAOCTGC TAATGTTGAT 420

GCTGOGAAAG AATGGAGGGA AGATAGAAAT GGAGAATTTA AAAGAAAAGT TGCCCGCTGT 480

40

GTAAGAAAAA GCCAAGAGAC TGCTTTTGAG 510

## (2) INFORMATION FOR SEQ ID NO:24:

## (i) SEQUENCE CHARACTERISTICS:

45

- (A) LENGTH: 617 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

50

## (ii) MOLECULE TYPE: DNA(genomic)

55

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vii) IMMEDIATE SOURCE:

(A) LIBRARY: Human fetal brain cDNA library

(B) CLONE: GEN-423A12

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 19..528

(x1) SEQUENCE DESCRIPTION: SEQ ID NO:24:

51	GGG	OCT	TOGG	CAG	GAG	ATG	ACG	GAG	CTG	CAG	TOG	GCA	CTG	CTA	CTG	OGA	51
						Met	Thr	Glu	Leu	Gln	Ser	Ala	Leu	Leu	Leu	Arg	
						1				5					10		
99	AGA	CAG	CTG	GCA	GAA	CTC	AAC	AAA	AAT	OCA	GTG	GAA	GGC	TTT	TCT	GCA	99
	Arg	Gln	Leu	Ala	Glu	Leu	Asn	Lys	Asn	Pro	Val	Glu	Gly	Phe	Ser	Ala	
				15					20					25			
147	GGT	TTA	ATA	GAT	GAC	AAT	GAT	CTC	TAC	OGA	TGG	GAA	GTC	CTT	ATT	ATT	147
	Gly	Leu	Ile	Asp	Asp	Asn	Asp	Leu	Tyr	Arg	Trp	Glu	Val	Leu	Ile	Ile	
				30				35					40				
195	GGC	OCT	OCA	GAT	ACA	CTT	TAT	GAA	GGT	GGT	GTT	TTT	AAG	GCT	CAT	CTT	195
	Gly	Pro	Pro	Asp	Thr	Leu	Tyr	Glu	Gly	Gly	Val	Phe	Lys	Ala	His	Leu	
		45					50					55					
243	ACT	TTC	OCA	AAA	GAT	TAT	CCC	CTC	OGA	OCT	OCT	AAA	ATG	AAA	TTC	ATT	243
	Thr	Phe	Pro	Lys	Asp	Tyr	Pro	Leu	Arg	Pro	Pro	Lys	Met	Lys	Phe	Ile	
		60				65				70					75		
291	ACA	GAA	ATC	TGG	CAC	OCA	AAT	GTT	GAT	AAA	AAT	GGT	GAT	GTG	TGC	ATT	291
	Thr	Glu	Ile	Trp	His	Pro	Asn	Val	Asp	Lys	Asn	Gly	Asp	Val	Cys	Ile	
				80				85						90			
339	TCT	ATT	CTT	CAT	GAG	OCT	GGG	GAA	GAT	AAG	TAT	GGT	TAT	GAA	AAG	OCA	339
	Ser	Ile	Leu	His	Glu	Pro	Gly	Glu	Asp	Lys	Tyr	Gly	Tyr	Glu	Lys	Pro	
				95				100						105			
387	GAG	GAA	OGC	TGG	CTC	OCT	ATC	CAC	ACT	GTG	GAA	ACC	ATC	ATG	ATT	AGT	387
	Glu	Glu	Arg	Trp	Leu	Pro	Ile	His	Thr	Val	Glu	Thr	Ile	Met	Ile	Ser	
				110				115					120				
435	GTC	ATT	TCT	ATG	CTG	GCA	GAC	OCT	AAT	GGA	GAC	TCA	OCT	GCT	AAT	GTT	435
	Val	Ile	Ser	Met	Leu	Ala	Asp	Pro	Asn	Gly	Asp	Ser	Pro	Ala	Asn	Val	
				125			130				135						

GAT GCT GCG AAA GAA TGG AGG GAA GAT AGA AAT GGA GAA TTT AAA AGA 483  
 Asp Ala Ala Lys Glu Trp Arg Glu Asp Arg Asn Gly Glu Phe Lys Arg  
 140 145 150 155

AAA GTT GCC CGC TGT GTA AGA AAA AGC CAA GAG ACT GCT TTT GAG 528  
 Lys Val Ala Arg Cys Val Arg Lys Ser Gln Glu Thr Ala Phe Glu  
 160 165 170

TGACATTTAT TTAGCAGCTA GTAAC TTCAC TTATTTTCAGG GTCTCCAATT GAGAAACATG 588  
 GCACTGTTTT TCCTGCACTC TACCCACCG 617

(2) INFORMATION FOR SEQ ID NO:25:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 374 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

Met Val Leu Trp Glu Ser Pro Arg Gln Cys Ser Ser Trp Thr Leu Cys  
 1 5 10 15

Glu Gly Phe Cys Trp Leu Leu Leu Leu Pro Val Met Leu Leu Ile Val  
 20 25 30

Ala Arg Pro Val Lys Leu Ala Ala Phe Pro Thr Ser Leu Ser Asp Cys  
 35 40 45

Gln Thr Pro Thr Gly Trp Asn Cys Ser Gly Tyr Asp Asp Arg Glu Asn  
 50 55 60

Asp Leu Phe Leu Cys Asp Thr Asn Thr Cys Lys Phe Asp Gly Glu Cys  
 65 70 75 80

Leu Arg Ile Gly Asp Thr Val Thr Cys Val Cys Gln Phe Lys Cys Asn  
 85 90 95

Asn Asp Tyr Val Pro Val Cys Gly Ser Asn Gly Glu Ser Tyr Gln Asn  
 100 105 110

Glu Cys Tyr Leu Arg Gln Ala Ala Cys Lys Gln Gln Ser Glu Ile Leu  
 115 120 125

Val Val Ser Glu Gly Ser Cys Ala Thr Asp Ala Gly Ser Gly Ser Gly

	130		135		140
5	Asp Gly Val His Glu Gly Ser Gly Glu Thr Ser Gln Lys Glu Thr Ser				
	145		150		155 160
	Thr Cys Asp Ile Cys Gln Phe Gly Ala Glu Cys Asp Glu Asp Ala Glu				
		165		170	175
10	Asp Val Trp Cys Val Cys Asn Ile Asp Cys Ser Gln Thr Asn Phe Asn				
		180		185	190
	Pro Leu Cys Ala Ser Asp Gly Lys Ser Tyr Asp Asn Ala Cys Gln Ile				
15		195		200	205
	Lys Glu Ala Ser Cys Gln Lys Gln Glu Lys Ile Glu Val Met Ser Leu				
		210		215	220
20	Gly Arg Cys Gln Asp Asn Thr Thr Thr Thr Thr Lys Ser Glu Asp Gly				
		225		230	235 240
	His Tyr Ala Arg Thr Asp Tyr Ala Glu Asn Ala Asn Lys Leu Glu Glu				
		245		250	255
25	Ser Ala Arg Glu His His Ile Pro Cys Pro Glu His Tyr Asn Gly Phe				
		260		265	270
	Cys Met His Gly Lys Cys Glu His Ser Ile Asn Met Gln Glu Pro Ser				
30		275		280	285
	Cys Arg Cys Asp Ala Gly Tyr Thr Gly Gln His Cys Glu Lys Lys Asp				
		290		295	300
35	Tyr Ser Val Leu Tyr Val Val Pro Gly Pro Val Arg Phe Gln Tyr Val				
		305		310	315 320
	Leu Ile Ala Ala Val Ile Gly Thr Ile Gln Ile Ala Val Ile Cys Val				
		325		330	335
40	Val Val Leu Cys Ile Thr Arg Lys Cys Pro Arg Ser Asn Arg Ile His				
		340		345	350
	Arg Gln Lys Gln Asn Thr Gly His Tyr Ser Ser Asp Asn Thr Thr Arg				
45		355		360	365
	Ala Ser Thr Arg Leu Ile				
		370			

(2) INFORMATION FOR SEQ ID NO:26:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1122 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA(genomic)

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

	ATGGTGCTGT GGGAGTCCCC GGGCAGTGC AGCAGCTGGA CACTTTGCGA GGGCTTTTGC	60
15	TGGCTGCTGC TGCTGCCCCG CATGCTACTC ATGCTAGGOC GCGCGGTGAA GCTGCTGCT	120
	TTCCCTAOCCT CCTTAAGTGA CTGOCAAAAG CCGACCGGCT GGAATTGCTC TGGTTATGAT	180
	GACAGAGAAA ATGATCTCTT CCTCTGTGAC ACGAACACCT GTAAATTTGA TGGGGAATGT	240
20	TTAAGAATTG GAGACACTGT GACTTGCGTC TGTCAGTTCA AGTGCAACAA TGACTATGTG	300
	CCTGTGTGTG GCTCCAATGG GGAGAGCTAC CAGAATGAGT GTTACCTGCG ACAGGCTGCA	360
25	TGCAACAGC AGAGTGAGAT ACTGTGTGGT TCAGAAGGAT CATGTGOCAC AGATGCAGGA	420
	TCAGGATCTG GAGATGGAGT CCGTGAAGGC TCTGGAGAAA CTAGTCAAAA GGAGACATOC	480
	ACCTGTGATA TTTGOCAGTT TGGTGCAGAA TGTGAAGAAG ATGCGGAGGA TGTCTGGTGT	540
30	GTGTGTAATA TTGACTGTTC TCAAAACCAAC TTCAATCCCC TCTGCGCTTC TGATGGGAAA	600
	TCTTATGATA ATGCATGACA AATCAAAGAA GCATOGTGTG AGAAACAGGA GAAAATTGAA	660
35	GTTCATGTCTT TGGGTGCGATG TCAAGATAAC ACAACTACAA CTAATAAGTC TGAAGATGGG	720
	CATTATGCAA GAACAGATTA TGCAGAGAAT GCTAACAAAT TAGAAGAAAAG TGCCAGAGAA	780
	CAACACATAC CTTGTCCGGA ACATTACAAT GGCTTCTGCA TGCATGGGAA GTGTGAGCAT	840
40	TCTATCAATA TGCAGGAGOC ATCTTGCAGG TGTGATGCTG GTTATACCTG ACAACACTGT	900
	GAAAAAAGG ACTACAGTGT TCTATAAGTT GTTCCCGGTC CTGTACGATT TCAGTATGTC	960
45	TTAATGCGAG CTGTGATTGG AACAAATTCAG ATTGCTGTCA TCTGTGTGGT GGTCCTCTGC	1020
	ATCACAAGGA AATGCCCCAG AAGCAACAGA ATTCACAGAC AGAAGCAAAA TACAGGGCAC	1080
	TACAGTTCAG ACAATACAAC AAGAGCGTCC ACGAGGTAA TC	1122

## (2) INFORMATION FOR SEQ ID NO:27:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1721 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA(genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vii) IMMEDIATE SOURCE:

- (A) LIBRARY: Human fetal brain cDNA library  
 (B) CLONE: GEN-092E10

(ix) FEATURE:

- (A) NAME/KEY: CDS  
 (B) LOCATION: 368..1489

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

```

CTGCGGGGGG OCTTGACTCT OCTTCAOOC TGCTCTCTOG GGCTOCACCTC GTCTGCCCCCT 60
GGACTCCCGT CTCTCTCTGT OCTCCGGCTT CACAGAGCTC OCTCTTTATG GCAGCAGCTT 120
CCCGGCTCTC CCGGCGAGCT TCTCAGGGGA GACCCCTCTC GCTCCGGGGC TGAGCCAGTC 180
OCTGGATGTT GCTGAACTC TOGAGATCAT GCGGGGGTTT GGCTGCTGCT TCCCCGGGG 240
GTGCCACTGC CACCGCGGCC GCTCTGCTG CCGCCGTCCG CCGGATGCTC AGTAGCCCCG 300
TGCCCCGGCC CCGGATCCT GTGTCTCTG GAAGCCGTTT GCTGCTGCAG AGTTGCAAGA 360
ACTAGTC ATG GTG CTG TGG GAG TOC CCG CCG CAG TGC AGC AGC TGG ACA 409
      Met Val Leu Trp Glu Ser Pro Arg Gln Cys Ser Ser Trp Thr
      1             5             10

CTT TGC GAG GGC TTT TGC TGG CTG CTG CTG CTG CCG GTC ATG CTA CTC 457
Leu Cys Glu Gly Phe Cys Trp Leu Leu Leu Leu Pro Val Met Leu Leu
  15             20             25             30

ATC GTA GCC CGC CCG GTG AAG CTC GCT GCT TTC OCT ACC TOC TTA AGT 505
Ile Val Ala Arg Pro Val Lys Leu Ala Ala Phe Pro Thr Ser Leu Ser
      35             40             45

GAC TGC CAA ACG CCC ACC GGC TGG AAT TGC TCT GGT TAT GAT GAC AGA 553

```

	Asp	Cys	Gln	Thr	Pro	Thr	Gly	Trp	Asn	Cys	Ser	Gly	Tyr	Asp	Asp	Arg	
				50					55					60			
5	GAA	AAT	GAT	CTC	TTC	CTC	TGT	GAC	AOC	AAC	AOC	TGT	AAA	TTT	GAT	GGG	601
	Glu	Asn	Asp	Leu	Phe	Leu	Cys	Asp	Thr	Asn	Thr	Cys	Lys	Phe	Asp	Gly	
			65					70					75				
10	GAA	TGT	TTA	AGA	ATT	GGA	GAC	ACT	GTG	ACT	TGC	GTC	TGT	CAG	TTC	AAG	649
	Glu	Cys	Leu	Arg	Ile	Gly	Asp	Thr	Val	Thr	Cys	Val	Cys	Gln	Phe	Lys	
		80					85					90					
15	TGC	AAC	AAT	GAC	TAT	GTG	OCT	GTG	TGT	GGC	TOC	AAT	GGG	GAG	AGC	TAC	697
	Cys	Asn	Asn	Asp	Tyr	Val	Pro	Val	Cys	Gly	Ser	Asn	Gly	Glu	Ser	Tyr	
		95				100					105					110	
20	CAG	AAT	GAG	TGT	TAC	CTG	OGA	CAG	GCT	GCA	TGC	AAA	CAG	CAG	AGT	GAG	745
	Gln	Asn	Glu	Cys	Tyr	Leu	Arg	Gln	Ala	Ala	Cys	Lys	Gln	Gln	Ser	Glu	
					115				120						125		
25	ATA	CTT	GTG	GTG	TCA	GAA	GGA	TCA	TGT	GOC	ACA	GAT	GCA	GGA	TCA	GGA	793
	Ile	Leu	Val	Val	Ser	Glu	Gly	Ser	Cys	Ala	Thr	Asp	Ala	Gly	Ser	Gly	
				130					135					140			
30	TCT	GGA	GAT	GGA	GTC	CAT	GAA	GGC	TCT	GGA	GAA	ACT	AGT	CAA	AAG	GAG	841
	Ser	Gly	Asp	Gly	Val	His	Glu	Gly	Ser	Gly	Glu	Thr	Ser	Gln	Lys	Glu	
			145					150					155				
35	ACA	TOC	AOC	TGT	GAT	ATT	TGC	CAG	TTT	GGT	GCA	GAA	TGT	GAC	GAA	GAT	889
	Thr	Ser	Thr	Cys	Asp	Ile	Cys	Gln	Phe	Gly	Ala	Glu	Cys	Asp	Glu	Asp	
		160					165					170					
40	GOC	GAG	GAT	GTC	TGG	TGT	GTG	TGT	AAT	ATT	GAC	TGT	TCT	CAA	AOC	AAC	937
	Ala	Glu	Asp	Val	Trp	Cys	Val	Cys	Asn	Ile	Asp	Cys	Ser	Gln	Thr	Asn	
		175				180					185					190	
45	TTC	AAT	CCC	CTC	TGC	GCT	TCT	GAT	GGG	AAA	TCT	TAT	GAT	AAT	GCA	TGC	985
	Phe	Asn	Pro	Leu	Cys	Ala	Ser	Asp	Gly	Lys	Ser	Tyr	Asp	Asn	Ala	Cys	
					195					200					205		
50	CAA	ATC	AAA	GAA	GCA	TOG	TGT	CAG	AAA	CAG	GAG	AAA	ATT	GAA	GTC	ATG	1033
	Gln	Ile	Lys	Glu	Ala	Ser	Cys	Gln	Lys	Gln	Glu	Lys	Ile	Glu	Val	Met	
				210					215					220			
55	TCT	TTG	GGT	OGA	TGT	CAA	GAT	AAC	ACA	ACT	ACA	ACT	ACT	AAG	TCT	GAA	1081
	Ser	Leu	Gly	Arg	Cys	Gln	Asp	Asn	Thr	Thr	Thr	Thr	Thr	Lys	Ser	Glu	
			225					230					235				
60	GAT	GGG	CAT	TAT	GCA	AGA	ACA	GAT	TAT	GCA	GAG	AAT	GCT	AAC	AAA	TTA	1129
	Asp	Gly	His	Tyr	Ala	Arg	Thr	Asp	Tyr	Ala	Glu	Asn	Ala	Asn	Lys	Leu	
			240				245					250					

5 GAA GAA AGT GGC AGA GAA CAC CAC ATA OCT TGT CCG GAA CAT TAC AAT 1177  
 Glu Glu Ser Ala Arg Glu His His Ile Pro Cys Pro Glu His Tyr Asn  
 255 260 265 270

GGC TTC TGC ATG CAT GGG AAG TGT GAG CAT TCT ATC AAT ATG CAG GAG 1225  
 Gly Phe Cys Met His Gly Lys Cys Glu His Ser Ile Asn Met Gln Glu  
 275 280 285

10 CCA TCT TGC AGG TGT GAT GCT GGT TAT ACT GGA CAA CAC TGT GAA AAA 1273  
 Pro Ser Cys Arg Cys Asp Ala Gly Tyr Thr Gly Gln His Cys Glu Lys  
 290 295 300

15 AAG GAC TAC AGT GTT CTA TAC GTT GTT CCC GGT OCT GTA OGA TTT CAG 1321  
 Lys Asp Tyr Ser Val Leu Tyr Val Val Pro Gly Pro Val Arg Phe Gln  
 305 310 315

TAT GTC TTA ATC GCA GCT GTG ATT GGA ACA ATT CAG ATT GCT GTC ATC 1369  
 Tyr Val Leu Ile Ala Ala Val Ile Gly Thr Ile Gln Ile Ala Val Ile  
 320 325 330

20 TGT GTG GTG GTC CTC TGC ATC ACA AGG AAA TGC CCC AGA AGC AAC AGA 1417  
 Cys Val Val Val Leu Cys Ile Thr Arg Lys Cys Pro Arg Ser Asn Arg  
 335 340 345 350

25 ATT CAC AGA CAG AAG CAA AAT ACA GGG CAC TAC AGT TCA GAC AAT ACA 1465  
 Ile His Arg Gln Lys Gln Asn Thr Gly His Tyr Ser Ser Asp Asn Thr  
 355 360 365

30 ACA AGA GCG TCC ACG AGG TTA ATC TAA AGGGAGCATG TTTCACAGTG 1512  
 Thr Arg Ala Ser Thr Arg Leu Ile  
 370

GCTGGACTAC CGAGAGCTTG GACTACACAA TACAGTATTA TAGACAAAAG AATAAGACAA 1572

35 GAGATCTACA CATGTTGCGT TGCATTTGTG GTAATCTACA CCAATGAAAA CATGTACTAC 1632

AGCTATATTT GATTATGTAT GGATATATTT GAAATAGTAT ACATTGTCCTT GATGTTTTTTT 1692

40 CTGTAATGTA AATAAACTAT TTATATCAC 1721

## (2) INFORMATION FOR SEQ ID NO:28:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 817 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein



## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

5 Met Gly Asp Thr Val Val Glu Pro Ala Pro Leu Lys Pro Thr Ser Glu  
 1 5 10 15  
 Pro Thr Ser Gly Pro Pro Gly Asn Asn Gly Gly Ser Leu Leu Ser Val  
 10 20 25 30  
 Ile Thr Glu Gly Val Gly Glu Leu Ser Val Ile Asp Pro Glu Val Ala  
 35 40 45  
 15 Gln Lys Ala Cys Gln Glu Val Leu Glu Lys Val Lys Leu Leu His Gly  
 50 55 60  
 Gly Val Ala Val Ser Ser Arg Gly Thr Pro Leu Glu Leu Val Asn Gly  
 65 70 75 80  
 20 Asp Gly Val Asp Ser Glu Ile Arg Cys Leu Asp Asp Pro Pro Ala Gln  
 85 90 95  
 Ile Arg Glu Glu Glu Asp Glu Met Gly Ala Ala Val Ala Ser Gly Thr  
 100 105 110  
 25 Ala Lys Gly Ala Arg Arg Arg Arg Gln Asn Asn Ser Ala Lys Gln Ser  
 115 120 125  
 30 Trp Leu Leu Arg Leu Phe Glu Ser Lys Leu Phe Asp Ile Ser Met Ala  
 130 135 140  
 Ile Ser Tyr Leu Tyr Asn Ser Lys Glu Pro Gly Val Gln Ala Tyr Ile  
 145 150 155 160  
 35 Gly Asn Arg Leu Phe Cys Phe Arg Asn Glu Asp Val Asp Phe Tyr Leu  
 165 170 175  
 Pro Gln Leu Leu Asn Met Tyr Ile His Met Asp Glu Asp Val Gly Asp  
 180 185 190  
 40 Ala Ile Lys Pro Tyr Ile Val His Arg Cys Arg Gln Ser Ile Asn Phe  
 195 200 205  
 45 Ser Leu Gln Cys Ala Leu Leu Leu Gly Ala Tyr Ser Ser Asp Met His  
 210 215 220  
 Ile Ser Thr Gln Arg His Ser Arg Gly Thr Lys Leu Arg Lys Leu Ile  
 225 230 235 240  
 50 Leu Ser Asp Glu Leu Lys Pro Ala His Arg Lys Arg Glu Leu Pro Ser  
 245 250 255  
 55

Leu Ser Pro Ala Pro Asp Thr Gly Leu Ser Pro Ser Lys Arg Thr His  
 260 265 270  
 5 Gln Arg Ser Lys Ser Asp Ala Thr Ala Ser Ile Ser Leu Ser Ser Asn  
 275 280 285  
 10 Leu Lys Arg Thr Ala Ser Asn Pro Lys Val Glu Asn Glu Asp Glu Glu  
 290 295 300  
 Leu Ser Ser Ser Thr Glu Ser Ile Asp Asn Ser Phe Ser Ser Pro Val  
 305 310 315 320  
 15 Arg Leu Ala Pro Glu Arg Glu Phe Ile Lys Ser Leu Met Ala Ile Gly  
 325 330 335  
 Lys Arg Leu Ala Thr Leu Pro Thr Lys Glu Gln Lys Thr Gln Arg Leu  
 340 345 350  
 20 Ile Ser Glu Leu Ser Leu Leu Asn His Lys Leu Pro Ala Arg Val Trp  
 355 360 365  
 Leu Pro Thr Ala Gly Phe Asp His His Val Val Arg Val Pro His Thr  
 370 375 380  
 25 Gln Ala Val Val Leu Asn Ser Lys Asp Lys Ala Pro Tyr Leu Ile Tyr  
 385 390 395 400  
 30 Val Glu Val Leu Glu Cys Glu Asn Phe Asp Thr Thr Ser Val Pro Ala  
 405 410 415  
 Arg Ile Pro Glu Asn Arg Ile Arg Ser Thr Arg Ser Val Glu Asn Leu  
 420 425 430  
 35 Pro Glu Cys Gly Ile Thr His Glu Gln Arg Ala Gly Ser Phe Ser Thr  
 435 440 445  
 Val Pro Asn Tyr Asp Asn Asp Asp Glu Ala Trp Ser Val Asp Asp Ile  
 450 455 460  
 40 Gly Glu Leu Gln Val Glu Leu Pro Glu Val His Thr Asn Ser Cys Asp  
 465 470 475 480  
 45 Asn Ile Ser Gln Phe Ser Val Asp Ser Ile Thr Ser Gln Glu Ser Lys  
 485 490 495  
 Glu Pro Val Phe Ile Ala Ala Gly Asp Ile Arg Arg Arg Leu Ser Glu  
 500 505 510  
 50 Gln Leu Ala His Thr Pro Thr Ala Phe Lys Arg Asp Pro Glu Asp Pro  
 515 520 525  
 55

Ser Ala Val Ala Leu Lys Glu Pro Trp Gln Glu Lys Val Arg Arg Ile  
 530 535 540  
 5 Arg Glu Gly Ser Pro Tyr Gly His Leu Pro Asn Trp Arg Leu Leu Ser  
 545 550 555 560  
 Val Ile Val Lys Cys Gly Asp Asp Leu Arg Gln Glu Leu Leu Ala Phe  
 565 570 575  
 10 Gln Val Leu Lys Gln Leu Gln Ser Ile Trp Glu Gln Glu Arg Val Pro  
 580 585 590  
 15 Leu Trp Ile Lys Pro Ile Gln Asp Ser Cys Glu Ile Thr Thr Asp Ser  
 595 600 605  
 Gly Met Ile Glu Pro Val Val Asn Ala Val Ser Ile His Gln Val Lys  
 610 615 620  
 20 Lys Gln Ser Gln Leu Ser Leu Leu Asp Tyr Phe Leu Gln Glu His Gly  
 625 630 635 640  
 Ser Tyr Thr Thr Glu Ala Phe Leu Ser Ala Gln Arg Asn Phe Val Gln  
 645 650 655  
 25 Ser Cys Ala Gly Tyr Cys Leu Val Cys Tyr Leu Leu Gln Val Lys Asp  
 660 665 670  
 30 Arg His Asn Gly Asn Ile Leu Leu Asp Ala Glu Gly His Ile Ile His  
 675 680 685  
 Ile Asp Phe Gly Phe Ile Leu Ser Ser Ser Pro Arg Asn Leu Gly Phe  
 690 695 700  
 35 Glu Thr Ser Ala Phe Lys Leu Thr Thr Glu Phe Val Asp Val Met Gly  
 705 710 715 720  
 Gly Leu Asp Gly Asp Met Phe Asn Tyr Tyr Lys Met Leu Met Leu Gln  
 725 730 735  
 40 Gly Leu Ile Ala Ala Arg Lys His Met Asp Lys Val Val Gln Ile Val  
 740 745 750  
 45 Glu Ile Met Gln Gln Gly Ser Gln Leu Pro Cys Phe His Gly Ser Ser  
 755 760 765  
 Thr Ile Arg Asn Leu Lys Glu Arg Phe His Met Ser Met Thr Glu Glu  
 770 775 780  
 50 Gln Leu Gln Leu Leu Val Glu Gln Met Val Asp Gly Ser Met Arg Ser  
 785 790 795 800  
 55

Ile Thr Thr Lys Leu Tyr Asp Gly Phe Gln Tyr Leu Thr Asn Gly Ile  
                   805                                  810                                  815

Met

(2) INFORMATION FOR SEQ ID NO:29:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 2451 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA(genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

20	ATGGGAGATA CAGTAGTGA GOCTGCCCCC TTGAAGCCAA CTTCTGAGCC CACTTCTGGC	60
	OCAACCAGGA ATAATGGGGG GTOOCTGCTA AGTGTCATCA CGGAGGGGGT CGGGGAACATA	120
25	TCAGTGATTTG ACOCTGAGGT GGCCAGAAAG GOCTGOCAGG AGGTGTTTGA GAAAGTCAAG	180
	CTTTTGTCATG GAGGCGTGGC AGTCTCTAGC AGAGCCACCC CACTGGAGTT GGTCAATGGG	240
	GATGGTGTGG ACAGTGAGAT COGTTGCTTA GATGATCCAC CTGCCCAGAT CAGGGAGGAG	300
30	GAAGATGAGA TGGGGGCOGC TGTGGCTCA GGCACAGCCA AAGGAGCAAG AAGACGGGGG	360
	CAGAACAACCT CAGCTAAACA GTCTTGGCTG CTGAGGCTGT TTGAGTCAAA ACTGTTTGAC	420
35	ATCTOCATGG CCATTTCATA OCTGTATAAC TOCAAGGAGC CTGGAGTACA AGOCTACATT	480
	GGCAACCGGC TCTTCTGCTT TOGCAACGAG GACGTGGACT TCTATCTGOC CCAGTTGCTT	540
	AACATGTACA TOCATATGA TGAGGAAGTG GGTGATGCCA TTAAGCCCTA CATAGTCCAC	600
40	CGTTGCOGCC AGAGCATTA CTTTTOCTC CAGTGTGCCC TGTGCTTGG GGCTATTCTT	660
	TCAGACATGC ACATTTCCAC TCAACGACAC TCCCGTGGGA CCAAGCTAAG GAAGCTGATC	720
45	CTCTCAGATG AGCTAAAGCC AGCTCACAGG AAGAGGGAGC TGCCCTOCTT GAGCCCGGCC	780
	OCTGATACAG GGCTGTCTCC CTCAAAAGG ACTCAACCAGC GCTCTAAGTC AGATGOCAC	840
	GCCAGCATAA GTCTCAGCAG CAAOCTGAAA CGAACAGCCA GCAACCOCTA AGTGGAGAAT	900
50	GAGGATGAGG AGCTCTCCTC CAGCACCGAG AGTATTGATA ATTCATTGAG TTCCOCTGTT	960

OGACTGGCTC CTGAGAGAGA ATTCATCAAG TOOCTGATGG OGATCGGCAA GGGGCTGGCC 1020  
 5 AOGCTOOCCA CCAAAGAGCA GAAAACACAG AGGCTGATCT CAGAGCTCTC CCTGCTCAAC 1080  
 CATAAGCTOC CTGCCCAGT CTGGCTGCCC ACTGCTGGCT TTGAACACCA CGTGGTCCGT 1140  
 GTACCCACCA CACAGGCTGT TGTCTCAAC TOCAAGGACA AGGCTCCCTA CCTGATTTAT 1200  
 10 GTGGAAGTCC TTGAATGTGA AAACCTTGAC ACCACCAAGTG TOOCTGCCC GATCCCGAG 1260  
 AACCGAATTC GGAGTACGAG GTCCTAGAA AACTTGCCG AATGTGGTAT TACCATGAG 1320  
 CAGGAGCTG GCAGCTTCAG CACTGTGCC AACTATGACA ACGATGATGA GGCTGGTGG 1380  
 15 GTGGATGACA TAGGCGAGCT GCAAGTGGAG CTCGCCAAG TGCATACCA CAGCTGTGAC 1440  
 AACATCTCC AGTTCTCTGT GGACAGCATC ACCAGCCAGG AGAGCAAGGA GCTGTGTTC 1500  
 20 ATTGCAGCAG GGGACATCC CGGGGCTTT TOGGAACAGC TGGCTCATAC CCGACAGCC 1560  
 TTCAAAGAG ACCAGAAGA TCTTCTGCA GTTCTCTCA AAGAGCCCTG GCAGGAGAAA 1620  
 GTAAGGGGGA TCAGAGAGGG CTCCTCTAC GGCCATCTCC CCAATTGGCG GCTCTGTCA 1680  
 25 GTCAATTGCA AGTGTGGGA TGAOCTTGG CAAGAGCTTC TGGCTTTCA GGTGTGAAG 1740  
 CAACTGCAGT CCATTTGGGA ACAGGAGCGA GTGCCCTTT GGATCAAGCC AATACAAGAT 1800  
 30 TCTTGTGAAA TTACGACTGA TAGTGGCATG ATTGAACCAG TGGTCAATGC TGTGTCCATC 1860  
 CATCAGGTGA AGAAACAGTC ACAGCTCTCC TTGCTGATT ACTTCTACA GGAGCACGGC 1920  
 AGTTACAACA CTGAGGCATT CCTCAGTCA CAGGCAATT TTGTGCAAAG TTGTGCTGGG 1980  
 35 TACTGCTTGG TCTGCTAACC GCTGCAAGTC AAGGACAGAC ACAATGGGAA TATCTTTTG 2040  
 GACGAGAAG GGCACATCAT CCACATGAC TTTGGCTTCA TCTCTCCAG CTCACCCGA 2100  
 40 AATCTGGGCT TTGAGACGTC AGCCTTTAAG CTGACCACAG AGTTTGTGGA TGTGATGGGC 2160  
 GGCTGGATG GGCACATGTT CAACTACTAT AAGATGCTGA TGCTGCAAGG GCTGATTGCC 2220  
 GCTCGAAAC ACATGGACAA GGTGGTGCAG ATGTGGAGA TCATGCAGCA AGGTTCTCAG 2280  
 45 CTTCCTTGCT TCCATGGCTC CAGCAOATT CGAAOCTCA AAGAGAGGTT CCACATGAGC 2340  
 ATGACTGAGG AGCAGCTGCA GCTGCTGGTG GAGCAGATGG TGGATGGCAG TATGCGGTCT 2400  
 50 ATCAACACCA AACTCTATGA CGGCTTCCAG TAOCTACCA ACGGCATCAT G 2451

55

## (2) INFORMATION FOR SEQ ID NO:30:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 3602 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA(genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

## (vii) IMMEDIATE SOURCE:

- (A) LIBRARY: Human fetal brain cDNA library  
 (B) CLONE: GEN-428B12c2

## (ix) FEATURE:

- (A) NAME/KEY: CDS  
 (B) LOCATION: 429..2879

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

```

GGTGGCTCAC GCGTGTAAATC CCAGCACTTT GGGAGGACAA GGCAGATCCC TTGAGCCCAG      60
GAGGTAGAGG CTGCAGTGAG CTGTGATGGT GCGACTGCAC TCAGGCGTGG GCAATGAAGC      120
AAGACCCATAT CTGAAAAAAA AAATTTTAA AAAAGGCAAA GATGGGCGCTG GGGCAACAAA      180
TATTCAGAG GAAAGGGAAC GTGTGTACTC CTTGAGGTGG GGAACATGAC CCACTTGAGG      240
TGCAGAAAGA AGACTTGTAT GGGGCTGGTG CAGGCTCCGC GCGCGCTGTC AGGGAAGGCG      300
AGGGGGCCAA TGGAAACCGG GAGCGGTGCG TGCTGCTGAG GCGGCAGTGT CGGCAGTCCA      360
ACCGGACTG CCGCAACCCC CTCGGGGGGG TCCCCAGAG CTTGGAAGCT CGAAGTCTGG      420
CTGTGGCC ATG GGA GAT ACA GTA GTG GAG OCT GOC CCG TTG AAG CCA ACT      470
      Met Gly Asp Thr Val Val Glu Pro Ala Pro Leu Lys Pro Thr
          1              5              10

TCT GAG CCG ACT TCT GGC CCA CCA GGG AAT AAT GGG GGG TCC CTG CTA      518
Ser Glu Pro Thr Ser Gly Pro Pro Gly Asn Asn Gly Gly Ser Leu Leu
      15              20              25              30

AGT GTC ATC ACG GAG GGG GTC GGG GAA CTA TCA GTG ATT GAC OCT GAG      566
Ser Val Ile Thr Glu Gly Val Gly Glu Leu Ser Val Ile Asp Pro Glu

```

	35								40				45					
5	GTG	GCC	CAG	AAG	GCC	TGC	CAG	GAG	GTG	TTG	GAG	AAA	GTC	AAG	CTT	TTG	614	
	Val	Ala	Gln	Lys	Ala	Cys	Gln	Glu	Val	Leu	Glu	Lys	Val	Lys	Leu	Leu		
				50					55					60				
10	CAT	GGA	GGC	GTG	GCA	GTC	TCT	AGC	AGA	GGC	ACC	CCA	CTG	GAG	TTG	GTC	662	
	His	Gly	Gly	Val	Ala	Val	Ser	Ser	Arg	Gly	Thr	Pro	Leu	Glu	Leu	Val		
			65				70					75						
15	AAT	GGG	GAT	GGT	GTG	GAC	AGT	GAG	ATC	OGT	TGC	CTA	GAT	GAT	OCA	CCT	710	
	Asn	Gly	Asp	Gly	Val	Asp	Ser	Glu	Ile	Arg	Cys	Leu	Asp	Asp	Pro	Pro		
		80					85					90						
20	GCC	CAG	ATC	AGG	GAG	GAG	GAA	GAT	GAG	ATG	GGG	GCC	GCT	GTG	GCC	TCA	758	
	Ala	Gln	Ile	Arg	Glu	Glu	Glu	Asp	Glu	Met	Gly	Ala	Ala	Val	Ala	Ser		
		95				100					105					110		
25	GGC	ACA	GCC	AAA	GGA	GCA	AGA	AGA	CGG	CGG	CAG	AAC	AAC	TCA	GCT	AAA	806	
	Gly	Thr	Ala	Lys	Gly	Ala	Arg	Arg	Arg	Arg	Gln	Asn	Asn	Ser	Ala	Lys		
					115					120					125			
30	CAG	TCT	TGG	CTG	CTG	AGG	CTG	TTT	GAG	TCA	AAA	CTG	TTT	GAC	ATC	TCC	854	
	Gln	Ser	Trp	Leu	Leu	Arg	Leu	Phe	Glu	Ser	Lys	Leu	Phe	Asp	Ile	Ser		
				130					135					140				
35	ATG	GCC	ATT	TCA	TAC	CTG	TAT	AAC	TCC	AAG	GAG	OCT	GGA	GTA	CAA	GCC	902	
	Met	Ala	Ile	Ser	Tyr	Leu	Tyr	Asn	Ser	Lys	Glu	Pro	Gly	Val	Gln	Ala		
			145					150					155					
40	TAC	ATT	GGC	AAC	CGG	CTC	TTC	TGC	TTT	CGC	AAC	GAG	GAC	GTG	GAC	TTC	950	
	Tyr	Ile	Gly	Asn	Arg	Leu	Phe	Cys	Phe	Arg	Asn	Glu	Asp	Val	Asp	Phe		
		160					165					170						
45	TAT	CTG	CCC	CAG	TTG	CTT	AAC	ATG	TAC	ATC	CAC	ATG	GAT	GAG	GAC	GTG	998	
	Tyr	Leu	Pro	Gln	Leu	Leu	Asn	Met	Tyr	Ile	His	Met	Asp	Glu	Asp	Val		
		175				180					185					190		
50	GGT	GAT	GCC	ATT	AAG	CCC	TAC	ATA	GTC	CAC	OGT	TGC	CGC	CAG	AGC	ATT	1046	
	Gly	Asp	Ala	Ile	Lys	Pro	Tyr	Ile	Val	His	Arg	Cys	Arg	Gln	Ser	Ile		
					195					200					205			
55	AAC	TTT	TCC	CTC	CAG	TGT	GCC	CTG	TTG	CTT	GGG	GCC	TAT	TCT	TCA	GAC	1094	
	Asn	Phe	Ser	Leu	Gln	Cys	Ala	Leu	Leu	Leu	Gly	Ala	Tyr	Ser	Ser	Asp		
				210				215						220				
60	ATG	CAC	ATT	TCC	ACT	CAA	CGA	CAC	TCC	OGT	GGG	ACC	AAG	CTA	CGG	AAG	1142	
	Met	His	Ile	Ser	Thr	Gln	Arg	His	Ser	Arg	Gly	Thr	Lys	Leu	Arg	Lys		
			225					230					235					

5	CTG ATC CTC TCA GAT GAG CTA AAG OCA GCT CAC AGG AAG AGG GAG CTG Leu Ile Leu Ser Asp Glu Leu Lys Pro Ala His Arg Lys Arg Glu Leu 240 245 250	1190
10	OCC TOC TTG AGC CCG GOC OCT GAT ACA GGG CTG TCT OCC TOC AAA AGG Pro Ser Leu Ser Pro Ala Pro Asp Thr Gly Leu Ser Pro Ser Lys Arg 255 260 265 270	1238
15	ACT CAC CAG CGC TCT AAG TCA GAT GOC ACT GOC AGC ATA AGT CTC AGC Thr His Gln Arg Ser Lys Ser Asp Ala Thr Ala Ser Ile Ser Leu Ser 275 280 285	1286
20	AGC AAC CTG AAA CGA ACA GOC AGC AAC CCT AAA GTG GAG AAT GAG GAT Ser Asn Leu Lys Arg Thr Ala Ser Asn Pro Lys Val Glu Asn Glu Asp 290 295 300	1334
25	GAG GAG CTC TOC TOC AGC ACC GAG AGT ATT GAT AAT TCA TTC AGT TOC Glu Glu Leu Ser Ser Ser Thr Glu Ser Ile Asp Asn Ser Phe Ser Ser 305 310 315	1382
30	OCT GTT CGA CTG GCT OCT GAG AGA GAA TTC ATC AAG TOC CTG ATG GCG Pro Val Arg Leu Ala Pro Glu Arg Glu Phe Ile Lys Ser Leu Met Ala 320 325 330	1430
35	ATC GGC AAG CGG CTG GOC ACG CTC OCC ACC AAA GAG CAG AAA ACA CAG Ile Gly Lys Arg Leu Ala Thr Leu Pro Thr Lys Glu Gln Lys Thr Gln 335 340 345 350	1478
40	AGG CTG ATC TCA GAG CTC TOC CTG CTC AAC CAT AAG CTC OCT GCC CGA Arg Leu Ile Ser Glu Leu Ser Leu Leu Asn His Lys Leu Pro Ala Arg 355 360 365	1526
45	GTC TGG CTG CCC ACT GCT GGC TTT GAC CAC CAC GTG GTC CGT GTA CCC Val Trp Leu Pro Thr Ala Gly Phe Asp His His Val Val Arg Val Pro 370 375 380	1574
50	CAC ACA CAG GCT GTT GTC CTC AAC TOC AAG GAC AAG GCT CCC TAC CTG His Thr Gln Ala Val Val Leu Asn Ser Lys Asp Lys Ala Pro Tyr Leu 385 390 395	1622
55	ATT TAT GTG GAA GTC CTT GAA TGT GAA AAC TTT GAC ACC ACC AGT GTC Ile Tyr Val Glu Val Leu Glu Cys Glu Asn Phe Asp Thr Thr Ser Val 400 405 410	1670
60	OCT GOC CGG ATC OCC GAG AAC CGA ATT CGG AGT ACG AGG TOC GTA GAA Pro Ala Arg Ile Pro Glu Asn Arg Ile Arg Ser Thr Arg Ser Val Glu 415 420 425 430	1718
65	AAC TTG CCC GAA TGT GGT ATT AOC CAT GAG CAG CGA GCT GGC AGC TTC Asn Leu Pro Glu Cys Gly Ile Thr His Glu Gln Arg Ala Gly Ser Phe 435 440 445 450	1766



	435							440					445					
5	AGC Ser	ACT Thr	GTG Val	CCC Pro	AAC Asn	TAT Tyr	GAC Asp	AAC Asn	GAT Asp	GAT Asp	GAG Glu	GCC Ala	TGG Trp	TCG Ser	GTG Val	GAT Asp	1814	
				450					455					460				
10	GAC Asp	ATA Ile	GGC Gly	GAG Glu	CTG Leu	CAA Gln	GTG Val	GAG Glu	CTC Leu	CCC Pro	GAA Glu	GTG Val	CAT His	AOC Thr	AAC Asn	AGC Ser	1862	
			465					470					475					
15	TGT Cys	GAC Asp	AAC Asn	ATC Ile	TOC Ser	CAG Gln	TTC Phe	TCT Ser	GTG Val	GAC Asp	AGC Ser	ATC Ile	AOC Thr	AGC Ser	CAG Gln	GAG Glu	1910	
		480					485					490						
20	AGC Ser	AAG Lys	GAG Glu	OCT Pro	GTG Val	TTC Phe	ATT Ile	GCA Ala	GCA Ala	GGG Gly	GAC Asp	ATC Ile	OGC Arg	OGG Arg	OGC Arg	CTT Leu	1958	
	495					500					505					510		
25	TOG Ser	GAA Glu	CAG Gln	CTG Leu	GCT Ala	CAT His	AOC Thr	COG Pro	ACA Thr	GOC Ala	TTC Phe	AAA Lys	OGA Arg	GAC Asp	CCA Pro	GAA Glu	2006	
					515					520					525			
30	GAT Asp	OCT Pro	TCT Ser	GCA Ala	GTT Val	GCT Ala	CTC Leu	AAA Lys	GAG Glu	CCC Pro	TGG Trp	CAG Gln	GAG Glu	AAA Lys	GTA Val	CGG Arg	2054	
				530					535					540				
35	OGG Arg	ATC Ile	AGA Arg	GAG Glu	GGC Gly	TOC Ser	CCC Pro	TAC Tyr	GGC Gly	CAT His	CTC Leu	CCC Pro	AAT Asn	TGG Trp	OGG Arg	CTC Leu	2102	
			545					550					555					
40	CTG Leu	TCA Ser	GTC Val	ATT Ile	GTC Val	AAG Lys	TGT Cys	GGG Gly	GAT Asp	GAC Asp	CTT Leu	OGG Arg	CAA Gln	GAG Glu	CTT Leu	CTG Leu	2150	
		560					565					570						
45	GOC Ala	TTT Phe	CAG Gln	GTG Val	TTG Leu	AAG Lys	CAA Gln	CTG Leu	CAG Gln	TOC Ser	ATT Ile	TGG Trp	GAA Glu	CAG Gln	GAG Glu	OGA Arg	2198	
	575					580					585					590		
50	GTG Val	CCC Pro	CTT Leu	TGG Trp	ATC Ile	AAG Lys	CCA Pro	ATA Ile	CAA Gln	GAT Asp	TCT Ser	TGT Cys	GAA Glu	ATT Ile	ACG Thr	ACT Thr	2246	
					595					600					605			
55	GAT Asp	AGT Ser	GGC Gly	ATG Met	ATT Ile	GAA Glu	CCA Pro	GTG Val	GTC Val	AAT Asn	GCT Ala	GTG Val	TOC Ser	ATC Ile	CAT His	CAG Gln	2294	
				610					615					620				
60	GTG Val	AAG Lys	AAA Lys	CAG Gln	TCA Ser	CAG Gln	CTC Leu	TOC Ser	TTG Leu	CTC Leu	GAT Asp	TAC Tyr	TTC Phe	CTA Leu	CAG Gln	GAG Glu	2342	
			625					630					635					

5	CAC GGC AGT TAC AOC ACT GAG GCA TTC CTC AGT GCA CAG CGC AAT TTT His Gly Ser Tyr Thr Thr Glu Ala Phe Leu Ser Ala Gln Arg Asn Phe 640 645 650	2390
10	GTG CAA AGT TGT GCT GGG TAC TGC TTG GTC TGC TAC CTG CTG CAA GTC Val Gln Ser Cys Ala Gly Tyr Cys Leu Val Cys Tyr Leu Leu Gln Val 655 660 665 670	2438
15	AAG GAC AGA CAC AAT GGG AAT ATC CTT TTG GAC GCA GAA GGC CAC ATC Lys Asp Arg His Asn Gly Asn Ile Leu Leu Asp Ala Glu Gly His Ile 675 680 685	2486
20	ATC CAC ATC GAC TTT GGC TTC ATC CTC TOC AGC TCA OCC CGA AAT CTG Ile His Ile Asp Phe Gly Phe Ile Leu Ser Ser Ser Pro Arg Asn Leu 690 695 700	2534
25	GGC TTT GAG ACG TCA GOC TTT AAG CTG ACC ACA GAG TTT GTG GAT GTG Gly Phe Glu Thr Ser Ala Phe Lys Leu Thr Thr Glu Phe Val Asp Val 705 710 715	2582
30	ATG GGC GGC CTG GAT GGC GAC ATG TTC AAC TAC TAT AAG ATG CTG ATG Met Gly Gly Leu Asp Gly Asp Met Phe Asn Tyr Tyr Lys Met Leu Met 720 725 730	2630
35	CTG CAA GGG CTG ATT GOC GCT OGG AAA CAC ATG GAC AAG GTG GTG CAG Leu Gln Gly Leu Ile Ala Ala Arg Lys His Met Asp Lys Val Val Gln 735 740 745 750	2678
40	ATC GTG GAG ATC ATG CAG CAA GGT TCT CAG CTT OCT TGC TTC CAT GGC Ile Val Glu Ile Met Gln Gln Gly Ser Gln Leu Pro Cys Phe His Gly 755 760 765	2726
45	TOC AGC ACC ATT CGA AAC CTC AAA GAG AGG TTC CAC ATG AGC ATG ACT Ser Ser Thr Ile Arg Asn Leu Lys Glu Arg Phe His Met Ser Met Thr 770 775 780	2774
50	GAG GAG CAG CTG CAG CTG CTG GTG GAG CAG ATG GTG GAT GGC AGT ATG Glu Glu Gln Leu Gln Leu Leu Val Glu Gln Met Val Asp Gly Ser Met 785 790 795	2822
55	OGG TCT ATC ACC ACC AAA CTC TAT GAC GGC TTC CAG TAC CTC ACC AAC Arg Ser Ile Thr Thr Lys Leu Tyr Asp Gly Phe Gln Tyr Leu Thr Asn 800 805 810	2870
60	GGC ATC ATG TGA CACGCTCTC AGCCCAGGAG TGGTGGGGGG TCAGGGCAC Gly Ile Met * 815	2922
65	OCTCOCTAGA GGGCOCTTGT CTGAGAAACC OCAAACAGG AAACCCACC TACCCAACCA	2982

TOCAOCCAAG GGAAATGGAA GGCAAGAAAC ACGAAGGATC ATGTGGTAAC TCGAGAGCT 3042  
 5 TGCTGAGGGG TGGGAGAGCC AGCTGTGGGG TOCAGACTTG TTGGGGCTTC OCTGCCCCCTC 3102  
 CTGGTCTGTG TCAGTATTAC CAOCAGACTG ACTCCAGGAC TCACTGCCCT CCAGAAAACA 3162  
 GAGGTGACAA ATGTGAGGGA CACTGGGGOC TTTCTTCTOC TTGTAGGGGT CTCTCAGAGG 3222  
 10 TTCTTTTCAC AGGOCATCCT CTTATTTCGT TCTGGGGOC AGGAAGTGGG GAAGAGTAGG 3282  
 TTCTCGGTAC TTAGGACTTG ATCCTGTGGT TGCCACTGGC CATGCTGCTG OCCAGCTCTA 3342  
 15 CCOCTOCCAG GGAOCTACCC CTOOCAGGA CCGACCOCTG GOCCAAGCTC COCTTGCTGG 3402  
 OGGGGCTGC GTGGGCCCTG CACTTGCTGA GGTTCOCCAT CATGGGCAAG GCAAGGGAAT 3462  
 TOCCACAGCC CTCAGTGTA CTGAGGTAC TGGCCTAGOC ATGTGGAATT COCTACCOCTG 3522  
 20 ACTOCTTOCC CAAOCCAGG GAAAAGAGCT CTCAATTTTT TATTTTAAAT TTTTGTGGA 3582  
 AATAAAGTCC TTAGTTAGCC 3602

25 (2) INFORMATION FOR SEQ ID NO:31:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 829 amino acids  
 (B) TYPE: amino acid  
 30 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

35 Met Arg Phe Leu Glu Ala Arg Ser Leu Ala Val Ala Met Gly Asp Thr  
 1 5 10 15  
 40 Val Val Glu Pro Ala Pro Leu Lys Pro Thr Ser Glu Pro Thr Ser Gly  
 20 25 30  
 Pro Pro Gly Asn Asn Gly Gly Ser Leu Leu Ser Val Ile Thr Glu Gly  
 35 40 45  
 45 Val Gly Glu Leu Ser Val Ile Asp Pro Glu Val Ala Gln Lys Ala Cys  
 50 55 60  
 Gln Glu Val Leu Glu Lys Val Lys Leu Leu His Gly Gly Val Ala Val  
 50 65 70 75 80

55

Ser Ser Arg Gly Thr Pro Leu Glu Leu Val Asn Gly Asp Gly Val Asp  
 85 90 95  
 5 Ser Glu Ile Arg Cys Leu Asp Asp Pro Pro Ala Gln Ile Arg Glu Glu  
 100 105 110  
 Glu Asp Glu Met Gly Ala Ala Val Ala Ser Gly Thr Ala Lys Gly Ala  
 115 120 125  
 10 Arg Arg Arg Arg Gln Asn Asn Ser Ala Lys Gln Ser Trp Leu Leu Arg  
 130 135 140  
 15 Leu Phe Glu Ser Lys Leu Phe Asp Ile Ser Met Ala Ile Ser Tyr Leu  
 145 150 155 160  
 Tyr Asn Ser Lys Glu Pro Gly Val Gln Ala Tyr Ile Gly Asn Arg Leu  
 165 170 175  
 20 Phe Cys Phe Arg Asn Glu Asp Val Asp Phe Tyr Leu Pro Gln Leu Leu  
 180 185 190  
 Asn Met Tyr Ile His Met Asp Glu Asp Val Gly Asp Ala Ile Lys Pro  
 195 200 205  
 25 Tyr Ile Val His Arg Cys Arg Gln Ser Ile Asn Phe Ser Leu Gln Cys  
 210 215 220  
 30 Ala Leu Leu Leu Gly Ala Tyr Ser Ser Asp Met His Ile Ser Thr Gln  
 225 230 235 240  
 Arg His Ser Arg Gly Thr Lys Leu Arg Lys Leu Ile Leu Ser Asp Glu  
 245 250 255  
 35 Leu Lys Pro Ala His Arg Lys Arg Glu Leu Pro Ser Leu Ser Pro Ala  
 260 265 270  
 Pro Asp Thr Gly Leu Ser Pro Ser Lys Arg Thr His Gln Arg Ser Lys  
 275 280 285  
 40 Ser Asp Ala Thr Ala Ser Ile Ser Leu Ser Ser Asn Leu Lys Arg Thr  
 290 295 300  
 45 Ala Ser Asn Pro Lys Val Glu Asn Glu Asp Glu Glu Leu Ser Ser Ser  
 305 310 315 320  
 Thr Glu Ser Ile Asp Asn Ser Phe Ser Ser Pro Val Arg Leu Ala Pro  
 325 330 335  
 50 Glu Arg Glu Phe Ile Lys Ser Leu Met Ala Ile Gly Lys Arg Leu Ala  
 340 345 350  
 55

Thr Leu Pro Thr Lys Glu Gln Lys Thr Gln Arg Leu Ile Ser Glu Leu  
 355 360 365  
 5 Ser Leu Leu Asn His Lys Leu Pro Ala Arg Val Trp Leu Pro Thr Ala  
 370 375 380  
 Gly Phe Asp His His Val Val Arg Val Pro His Thr Gln Ala Val Val  
 385 390 395 400  
 10 Leu Asn Ser Lys Asp Lys Ala Pro Tyr Leu Ile Tyr Val Glu Val Leu  
 405 410 415  
 Glu Cys Glu Asn Phe Asp Thr Thr Ser Val Pro Ala Arg Ile Pro Glu  
 420 425 430  
 15 Asn Arg Ile Arg Ser Thr Arg Ser Val Glu Asn Leu Pro Glu Cys Gly  
 435 440 445  
 Ile Thr His Glu Gln Arg Ala Gly Ser Phe Ser Thr Val Pro Asn Tyr  
 450 455 460  
 Asp Asn Asp Asp Glu Ala Trp Ser Val Asp Asp Ile Gly Glu Leu Gln  
 465 470 475 480  
 25 Val Glu Leu Pro Glu Val His Thr Asn Ser Cys Asp Asn Ile Ser Gln  
 485 490 495  
 Phe Ser Val Asp Ser Ile Thr Ser Gln Glu Ser Lys Glu Pro Val Phe  
 500 505 510  
 Ile Ala Ala Gly Asp Ile Arg Arg Arg Leu Ser Glu Gln Leu Ala His  
 515 520 525  
 35 Thr Pro Thr Ala Phe Lys Arg Asp Pro Glu Asp Pro Ser Ala Val Ala  
 530 535 540  
 Leu Lys Glu Pro Trp Gln Glu Lys Val Arg Arg Ile Arg Glu Gly Ser  
 545 550 555 560  
 40 Pro Tyr Gly His Leu Pro Asn Trp Arg Leu Leu Ser Val Ile Val Lys  
 565 570 575  
 Cys Gly Asp Asp Leu Arg Gln Glu Leu Leu Ala Phe Gln Val Leu Lys  
 580 585 590  
 Gln Leu Gln Ser Ile Trp Glu Gln Glu Arg Val Pro Leu Trp Ile Lys  
 595 600 605  
 50 Pro Ile Gln Asp Ser Cys Glu Ile Thr Thr Asp Ser Gly Met Ile Glu  
 610 615 620  
 55

5      Pro Val Val Asn Ala Val Ser Ile His Gln Val Lys Lys Gln Ser Gln  
       625                                  630                                  635                                  640  
 Leu Ser Leu Leu Asp Tyr Phe Leu Gln Glu His Gly Ser Tyr Thr Thr  
                                 645                                  650                                  655  
 10      Glu Ala Phe Leu Ser Ala Gln Arg Asn Phe Val Gln Ser Cys Ala Gly  
                                 660                                  665                                  670  
 Tyr Cys Leu Val Cys Tyr Leu Leu Gln Val Lys Asp Arg His Asn Gly  
                                 675                                  680                                  685  
 15      Asn Ile Leu Leu Asp Ala Glu Gly His Ile Ile His Ile Asp Phe Gly  
                                 690                                  695                                  700  
 Phe Ile Leu Ser Ser Ser Pro Arg Asn Leu Gly Phe Glu Thr Ser Ala  
       705                                  710                                  715                                  720  
 20      Phe Lys Leu Thr Thr Glu Phe Val Asp Val Met Gly Gly Leu Asp Gly  
                                 725                                  730                                  735  
 Asp Met Phe Asn Tyr Tyr Lys Met Leu Met Leu Gln Gly Leu Ile Ala  
                                 740                                  745                                  750  
 25      Ala Arg Lys His Met Asp Lys Val Val Gln Ile Val Glu Ile Met Gln  
                                 755                                  760                                  765  
 30      Gln Gly Ser Gln Leu Pro Cys Phe His Gly Ser Ser Thr Ile Arg Asn  
                                 770                                  775                                  780  
 Leu Lys Glu Arg Phe His Met Ser Met Thr Glu Glu Gln Leu Gln Leu  
       785                                  790                                  795                                  800  
 35      Leu Val Glu Gln Met Val Asp Gly Ser Met Arg Ser Ile Thr Thr Lys  
                                 805                                  810                                  815  
 40      Leu Tyr Asp Gly Phe Gln Tyr Leu Thr Asn Gly Ile Met  
                                 820                                  825

## (2) INFORMATION FOR SEQ ID NO:32:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2487 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA(genomic)

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

5	ATGAGATTCT TGGAAGCTOG AAGTCTGGCT GTGGOCATGG GAGATACAGT AGTGGAGOCT	60
	GCCCCCTTGA AGCCAACTTC TGAGOOCACT TCTGGCCCAC CAGGGAATAA TGGGGGGTCC	120
	CTGCTAAGTG TCATCAOCCA GGGGGTGGG GAACTATCAG TGATTGACCC TGAGGTGGCC	180
10	CAGAAGGOCT GOCAGGAGGT GTTGAGAGAA GTCAAGCTTT TGCATGGAGG CGTGGCAGTC	240
	TCTAGCAGAG GCACCCCACT GGAGTTGGTC AATGGGGATG GTGTGGACAG TGAGATCOGT	300
15	TGCTTAGATG ATCAOCTGC CCAGATCAGG GAGGAGGAAG ATGAGATGGG GGCGCTGTG	360
	GOCTCAGGCA CAGOCAGG AGCAAGAAGA CGGOGGCAGA ACAACTCAGC TAAACAGTCT	420
	TGGCTGCTGA GGCTGTTTGA GTCAAACTG TTTGACATCT CCATGGOCAT TTCATAOCTG	480
20	TATAACTOCA AGGAGCTGG AGTACAAGOC TACATTGGCA ACOGGCTCTT CTGCTTTGCG	540
	AAOGAGGACG TGGACTTCTA TCTGCCCCAG TTGCTTAACA TGTACATCCA CATGGATGAG	600
25	GACGTGGGTG ATGCCATTAA GCOCTACATA GTCCACCGTT GCGCCAGAG CATTAACTTT	660
	TOOCTOCAGT GTGCCCTGTT GCTTGGGGCC TATTCTTCAG ACATGCACAT TTCACTCAA	720
	CGACACTOCC GTGGGAOCAA GCTACGGAAG CTGATCTCTT CAGATGAGCT AAAGCCAGCT	780
30	CACAGGAAGA GGGAGCTGCC CTCTTGAGC CCGGCCCCCTG ATACAGGGCT GTCTOCTOC	840
	AAAAGGACTC ACCAGGCTC TAAGTCAGAT GOCCTGCCA GCATAAGTCT CAGCAGCAAC	900
35	CTGAAACGAA CAGCCAGCAA CCTAAAGTG GAGAATGAGG ATGAGGAGCT CTCTOCAGC	960
	ACCGAGAGTA TTGATAATTC ATTCAAGTTC CCTGTTGAC TGGCTOCTGA GAGAGAATTC	1020
	ATCAAGTOCC TGATGGOGAT CGCAAGGG CTGGCCAGC TCCCCACCA AGAGCAGAAA	1080
40	ACACAGAGGC TGATCTCAGA GCTCTOCTG CTCAACATA AGCTOCTGC CCGAGTCTGG	1140
	CTGCOCACTG CTGGCTTTGA CCAOCCAGTG GTCOGTGTAC CACACACA GGCTGTTGTC	1200
	CTCAACTCCA AGGACAAGGC TOCTAOCCTG ATTTATGTGG AAGTCTTGA ATGTGAAAAC	1260
45	TTTGACACCA CCAGTGTGCC TGCCCGGATC CCGAGAAOC GAATTGGAG TACGAGGTCC	1320
	GTAAGAACT TGCCGAATG TGGTATTACC CATGAGCAGC GAGCTGGCAG CTTCACTACT	1380
50	GTGCOCACT ATGACAAOGA TGATGAGGCC TGGTOGGTG ATGACATAGG CGAGCTGCAA	1440

55

GTGGAGCTOC CCGAAGTGCA TACCAACAGC TGTGACAACA TCTOCCAGTT CTCTGTGGAC 1500  
 5 AGCATCACCA GOCAGGAGAG CAAGGAGOCT GTGTTTCATTG CAGCAGGGGA CATOOGCOGG 1560  
 CGOCTTTTOGG AACAGCTGGC TCATAOCCOOG ACAGOCTTCA AACGAGACOC AGAAGATOCT 1620  
 TCTGCAGTTG CTCTCAAAGA GOCCTGGCAG GAGAAAGTAC GGGGGATCAG AGAGGGCTOC 1680  
 10 COCTAOGGOC ATCTOCCCAA TTGGGGGCTC CTGTCAGTCA TTGTCAAGTG TGGGGATGAC 1740  
 CTTOGGCAAG AGCTTCTGGC CTTTCAGGTG TTGAAGCAAC TGCAGTOCAT TTGGGAACAG 1800  
 GAGCGAGTGC COCTTTGGAT CAAGCCAATA CAAGATTCTT GTGAAATTAC GACTGATAGT 1860  
 15 GGCATGATTG AACCAGTGGT CAATGCTGTG TOCATOCATC AGGTGAAGAA ACAGTCACAG 1920  
 CTCTOCTTGC TOGATTACTT OCTACAGGAG CACGGCAGTT ACACCACTGA GGCATTCTCT 1980  
 20 AGTGCACAGC GCAATTTTGT GCAAAGTTGT GCTGGGTACT GCTTGGTCTG CTACCTGCTG 2040  
 CAAGTCAAGG ACAGACACAA TGGGAATATC CTTTGTGAGC CAGAAGGCCA CATCATOCAC 2100  
 ATOGACTTTG GCTTCATOCT CTCCAGCTCA CCCCAGAAATC TGGGCTTTGA GACGTCAGOC 2160  
 25 TTTAAGCTGA OCACAGAGTT TGTGGATGTG ATGGGGGGOC TGGATGGCGA CATGTTCAAC 2220  
 TACTATAAGA TGCTGATGCT GCAAGGGCTG ATTGCOGCTC GGAAACACAT GGACAAGGTG 2280  
 30 GTGCAGATOG TGGAGATCAT GCAGCAAGGT TCTCAGCTTC CTTGCTTCCA TGGCTOCAGC 2340  
 ACCATTOGAA AACTCAAAGA GAGGTTCCAC ATGAGCATGA CTGAGGAGCA GCTGCAGCTG 2400  
 CTGGTGGAGC AGATGGTGGG TGGCAGTATG OGGTCTATCA OCAOCAAAC CTATGAOGGC 2460  
 35 TTCCAGTAOC TCACCAACGG CATCATG 2487

## (2) INFORMATION FOR SEQ ID NO:33:

- 40 (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 3324 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 45 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA(genomic)

(iii) HYPOTHETICAL: NO

50 (iv) ANTI-SENSE: NO

55



## (vii) IMMEDIATE SOURCE:

(A) LIBRARY: Human fetal brain cDNA library

(B) CLONE: GEN-428B12c1

## (ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 115..2601

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

	COGGAATTCC GGGGAAGGCOG GAGCAAGTTT TGAAGAAGTC CCTATCAGAT TACACTTGGT	60
15	TGACTACTOC GGAGCAGOCA CTAAGAGGGA TGAACAGGOC TGCGTGGAAA TTGA ATG	117
	Met	
	1	
20	AGA TTC TTG GAA GCT CGA AGT CTG GCT GTG GOC ATG GGA GAT ACA GTA	165
	Arg Phe Leu Glu Ala Arg Ser Leu Ala Val Ala Met Gly Asp Thr Val	
	5 10 15	
25	GTG GAG OCT GOC OOC TTG AAG OCA ACT TCT GAG OOC ACT TCT GGC OCA	213
	Val Glu Pro Ala Pro Leu Lys Pro Thr Ser Glu Pro Thr Ser Gly Pro	
	20 25 30	
30	OCA GGG AAT AAT GGG GGG TOC CTG CTA AGT GTC ATC ACG GAG GGG GTC	261
	Pro Gly Asn Asn Gly Gly Ser Leu Leu Ser Val Ile Thr Glu Gly Val	
	35 40 45	
35	GGG GAA CTA TCA GTG ATT GAC OCT GAG GTG GOC CAG AAG GOC TGC CAG	309
	Gly Glu Leu Ser Val Ile Asp Pro Glu Val Ala Gln Lys Ala Cys Gln	
	50 55 60 65	
40	GAG GTG TTG GAG AAA GTC AAG CTT TTG CAT GGA GGC GTG GCA GTC TCT	357
	Glu Val Leu Glu Lys Val Lys Leu Leu His Gly Gly Val Ala Val Ser	
	70 75 80	
45	AGC AGA GGC ACC OCA CTG GAG TTG GTC AAT GGG GAT GGT GTG GAC AGT	405
	Ser Arg Gly Thr Pro Leu Glu Leu Val Asn Gly Asp Gly Val Asp Ser	
	85 90 95	
50	GAG ATC OGT TGC CTA GAT GAT OCA OCT GOC CAG ATC AGG GAG GAG GAA	453
	Glu Ile Arg Cys Leu Asp Asp Pro Pro Ala Gln Ile Arg Glu Glu Glu	
	100 105 110	
55	GAT GAG ATG GGG GOC GCT GTG GOC TCA GGC ACA GOC AAA GGA GCA AGA	501
	Asp Glu Met Gly Ala Ala Val Ala Ser Gly Thr Ala Lys Gly Ala Arg	
	115 120 125	
60	AGA CGG CGG CAG AAC AAC TCA GCT AAA CAG TCT TGG CTG CTG AGG CTG	549

	Arg	Arg	Arg	Gln	Asn	Asn	Ser	Ala	Lys	Gln	Ser	Trp	Leu	Leu	Arg	Leu	
	130					135					140					145	
5	TTT	GAG	TCA	AAA	CTG	TTT	GAC	ATC	TOC	ATG	GCC	ATT	TCA	TAC	CTG	TAT	597
	Phe	Glu	Ser	Lys	Leu	Phe	Asp	Ile	Ser	Met	Ala	Ile	Ser	Tyr	Leu	Tyr	
					150					155					160		
10	AAC	TOC	AAG	GAG	OCT	GGA	GTA	CAA	GCC	TAC	ATT	GGC	AAC	CGG	CTC	TTC	645
	Asn	Ser	Lys	Glu	Pro	Gly	Val	Gln	Ala	Tyr	Ile	Gly	Asn	Arg	Leu	Phe	
					165				170					175			
15	TGC	TTT	CGC	AAC	GAG	GAC	GTG	GAC	TTC	TAT	CTG	CCC	CAG	TTG	CTT	AAC	693
	Cys	Phe	Arg	Asn	Glu	Asp	Val	Asp	Phe	Tyr	Leu	Pro	Gln	Leu	Leu	Asn	
				180				185					190				
20	ATG	TAC	ATC	CAC	ATG	GAT	GAG	GAC	GTG	GGT	GAT	GCC	ATT	AAG	CCC	TAC	741
	Met	Tyr	Ile	His	Met	Asp	Glu	Asp	Val	Gly	Asp	Ala	Ile	Lys	Pro	Tyr	
							200					205					
25	ATA	GTC	CAC	OGT	TGC	OGC	CAG	AGC	ATT	AAC	TTT	TOC	CTC	CAG	TGT	GCC	789
	Ile	Val	His	Arg	Cys	Arg	Gln	Ser	Ile	Asn	Phe	Ser	Leu	Gln	Cys	Ala	
	210					215					220					225	
30	CTG	TTG	CTT	GGG	GCC	TAT	TCT	TCA	GAC	ATG	CAC	ATT	TOC	ACT	CAA	OGA	837
	Leu	Leu	Leu	Gly	Ala	Tyr	Ser	Ser	Asp	Met	His	Ile	Ser	Thr	Gln	Arg	
					230					235					240		
35	CAC	TOC	OGT	GGG	AOC	AAG	CTA	OGG	AAG	CTG	ATC	CTC	TCA	GAT	GAG	CTA	885
	His	Ser	Arg	Gly	Thr	Lys	Leu	Arg	Lys	Leu	Ile	Leu	Ser	Asp	Glu	Leu	
				245					250					255			
40	AAG	OCA	GCT	CAC	AGG	AAG	AGG	GAG	CTG	CCC	TOC	TTG	AGC	CCG	GCC	OCT	933
	Lys	Pro	Ala	His	Arg	Lys	Arg	Glu	Leu	Pro	Ser	Leu	Ser	Pro	Ala	Pro	
				260				265					270				
45	GAT	ACA	GGG	CTG	TCT	CCC	TOC	AAA	AGG	ACT	CAC	CAG	OGC	TCT	AAG	TCA	981
	Asp	Thr	Gly	Leu	Ser	Pro	Ser	Lys	Arg	Thr	His	Gln	Arg	Ser	Lys	Ser	
		275					280					285					
50	GAT	GCC	ACT	GCC	AGC	ATA	AGT	CTC	AGC	AGC	AAC	CTG	AAA	CGA	ACA	GCC	1029
	Asp	Ala	Thr	Ala	Ser	Ile	Ser	Leu	Ser	Ser	Asn	Leu	Lys	Arg	Thr	Ala	
						295					300					305	
55	AGC	AAC	OCT	AAA	GTG	GAG	AAT	GAG	GAT	GAG	GAG	CTC	TOC	TOC	AGC	AOC	1077
	Ser	Asn	Pro	Lys	Val	Glu	Asn	Glu	Asp	Glu	Glu	Leu	Ser	Ser	Ser	Thr	
					310					315					320		
60	GAG	AGT	ATT	GAT	AAT	TCA	TTC	AGT	TOC	OCT	GTT	OGA	CTG	GCT	OCT	GAG	1125
	Glu	Ser	Ile	Asp	Asn	Ser	Phe	Ser	Ser	Pro	Val	Arg	Leu	Ala	Pro	Glu	
				325					330					335			

	AGA	GAA	TTC	ATC	AAG	TCC	CTG	ATG	GCG	ATC	GGC	AAG	CGG	CTG	GCC	ACG	1173
	Arg	Glu	Phe	Ile	Lys	Ser	Leu	Met	Ala	Ile	Gly	Lys	Arg	Leu	Ala	Thr	
			340					345					350				
5																	
	CTC	CCC	ACC	AAA	GAG	CAG	AAA	ACA	CAG	AGG	CTG	ATC	TCA	GAG	CTC	TCC	1221
	Leu	Pro	Thr	Lys	Glu	Gln	Lys	Thr	Gln	Arg	Leu	Ile	Ser	Glu	Leu	Ser	
		355					360					365					
10																	
	CTG	CTC	AAC	CAT	AAG	CTC	CCT	GCC	CGA	GTC	TGG	CTG	CCC	ACT	GCT	GGC	1269
	Leu	Leu	Asn	His	Lys	Leu	Pro	Ala	Arg	Val	Trp	Leu	Pro	Thr	Ala	Gly	
		370				375					380					385	
15																	
	TTT	GAC	CAC	CAC	GTG	GTC	CGT	GTA	CCC	CAC	ACA	CAG	GCT	GTT	GTC	CTC	1317
	Phe	Asp	His	His	Val	Val	Arg	Val	Pro	His	Thr	Gln	Ala	Val	Val	Leu	
					390					395					400		
20																	
	AAC	TOC	AAG	GAC	AAG	GCT	CCC	TAC	CTG	ATT	TAT	GTG	GAA	GTC	CTT	GAA	1365
	Asn	Ser	Lys	Asp	Lys	Ala	Pro	Tyr	Leu	Ile	Tyr	Val	Glu	Val	Leu	Glu	
				405					410					415			
25																	
	TGT	GAA	AAC	TTT	GAC	ACC	ACC	AGT	GTC	CCT	GCC	CGG	ATC	CCC	GAG	AAC	1413
	Cys	Glu	Asn	Phe	Asp	Thr	Thr	Ser	Val	Pro	Ala	Arg	Ile	Pro	Glu	Asn	
			420					425					430				
30																	
	OGA	ATT	CGG	AGT	ACG	AGG	TCC	GTA	GAA	AAC	TTG	CCC	GAA	TGT	GGT	ATT	1461
	Arg	Ile	Arg	Ser	Thr	Arg	Ser	Val	Glu	Asn	Leu	Pro	Glu	Cys	Gly	Ile	
		435					440					445					
35																	
	ACC	CAT	GAG	CAG	OGA	GCT	GGC	AGC	TTC	AGC	ACT	GTG	CCC	AAC	TAT	GAC	1509
	Thr	His	Glu	Gln	Arg		Gly	Ser	Phe	Ser	Thr	Val	Pro	Asn	Tyr	Asp	
		450				455					460					465	
40																	
	AAC	GAT	GAT	GAG	GCC	TGG	TOG	GTG	GAT	GAC	ATA	GGC	GAG	CTG	CAA	GTG	1557
	Asn	Asp	Asp	Glu	Ala	Trp	Ser	Val	Asp	Asp	Ile	Gly	Glu	Leu	Gln	Val	
					470					475					480		
45																	
	GAG	CTC	CCC	GAA	GTG	CAT	ACC	AAC	AGC	TGT	GAC	AAC	ATC	TCC	CAG	TTC	1605
	Glu	Leu	Pro	Glu	Val	His	Thr	Asn	Ser	Cys	Asp	Asn	Ile	Ser	Gln	Phe	
				485					490					495			
50																	
	TCT	GTG	GAC	AGC	ATC	ACC	AGC	CAG	GAG	AGC	AAG	GAG	CCT	GTG	TTC	ATT	1653
	Ser	Val	Asp	Ser	Ile	Thr	Ser	Gln	Glu	Ser	Lys	Glu	Pro	Val	Phe	Ile	
			500					505					510				
55																	
	GCA	GCA	GGG	GAC	ATC	CGC	CGG	CGC	CTT	TOG	GAA	CAG	CTG	GCT	CAT	ACC	1701
	Ala	Ala	Gly	Asp	Ile	Arg	Arg	Arg	Leu	Ser	Glu	Gln	Leu	Ala	His	Thr	
		515					520					525					
60																	
	CCG	ACA	GCC	TTC	AAA	CGA	GAC	CCA	GAA	GAT	CCT	TCT	GCA	GTT	GCT	CTC	1749
	Pro	Thr	Ala	Phe	Lys	Arg	Asp	Pro	Glu	Asp	Pro	Ser	Ala	Val	Ala	Leu	

	530		535		540		545	
5	AAA GAG CCC TGG CAG GAG AAA GTA CCG CCG ATC AGA GAG GGC TCC CCC							1797
	Lys Glu Pro Trp Gln Glu Lys Val Arg Arg Ile Arg Glu Gly Ser Pro							
			550			555		560
10	TAC GGC CAT CTC CCC AAT TGG CCG CTC CTG TCA GTC ATT GTC AAG TGT							1845
	Tyr Gly His Leu Pro Asn Trp Arg Leu Leu Ser Val Ile Val Lys Cys							
			565			570		575
15	GGG GAT GAC CTT CCG CAA GAG CTT CTG GGC TTT CAG GTG TTG AAG CAA							1893
	Gly Asp Asp Leu Arg Gln Glu Leu Leu Ala Phe Gln Val Leu Lys Gln							
			580			585		590
	CTG CAG TCC ATT TGG GAA CAG GAG CGA GTG CCC CTT TGG ATC AAG CCA							1941
	Leu Gln Ser Ile Trp Glu Gln Glu Arg Val Pro Leu Trp Ile Lys Pro							
			595			600		605
20	ATA CAA GAT TCT TGT GAA ATT ACG ACT GAT AGT GGC ATG ATT GAA CCA							1989
	Ile Gln Asp Ser Cys Glu Ile Thr Thr Asp Ser Gly Met Ile Glu Pro							
			610			615		620
25	GTG GTC AAT GCT GTG TCC ATC CAT CAG GTG AAG AAA CAG TCA CAG CTC							2037
	Val Val Asn Ala Val Ser Ile His Gln Val Lys Lys Gln Ser Gln Leu							
			630			635		640
30	TCC TTG CTC GAT TAC TTC CTA CAG GAG CAC GGC AGT TAC ACC ACT GAG							2085
	Ser Leu Leu Asp Tyr Phe Leu Gln Glu His Gly Ser Tyr Thr Thr Glu							
			645			650		655
35	GCA TTC CTC AGT GCA CAG CCG AAT TTT GTG CAA AGT TGT GCT GGC TAC							2133
	Ala Phe Leu Ser Ala Gln Arg Asn Phe Val Gln Ser Cys Ala Gly Tyr							
			660			665		670
	TGC TTG GTC TGC TAC CTG CTG CAA GTC AAG GAC AGA CAC AAT GGC AAT							2181
	Cys Leu Val Cys Tyr Leu Leu Gln Val Lys Asp Arg His Asn Gly Asn							
			675			680		685
40	ATC CTT TTG GAC GCA GAA GGC CAC ATC ATC CAC ATC GAC TTT GGC TTC							2229
	Ile Leu Leu Asp Ala Glu Gly His Ile Ile His Ile Asp Phe Gly Phe							
			690			695		700
45	ATC CTC TCC AGC TCA CCC CGA AAT CTG GGC TTT GAG ACG TCA GGC TTT							2277
	Ile Leu Ser Ser Ser Pro Arg Asn Leu Glu Phe Glu Thr Ser Ala Phe							
			710			715		720
50	AAG CTG ACC ACA GAG TTT GTG GAT GTG ATG GGC GGC CTG GAT GGC GAC							2325
	Lys Leu Thr Thr Glu Phe Val Asp Val Met Gly Gly Leu Asp Gly Asp							
			725			730		735
55								

5 ATG TTC AAC TAC TAT AAG ATG CTG ATG CTG CAA GGG CTG ATT GOC GCT 2373  
 Met Phe Asn Tyr Tyr Lys Met Leu Met Leu Gln Gly Leu Ile Ala Ala  
 740 745 750  
 OGG AAA CAC ATG GAC AAG GTG GTG CAG ATC GTG GAG ATC ATG CAG CAA 2421  
 Arg Lys His Met Asp Lys Val Val Gln Ile Val Glu Ile Met Gln Gln  
 755 760 765  
 10 GGT TCT CAG CTT OCT TGC TTC CAT GGC TOC AGC ACC ATT OGA AAC CTC 2469  
 Gly Ser Gln Leu Pro Cys Phe His Gly Ser Ser Thr Ile Arg Asn Leu  
 770 775 780 785  
 15 AAA GAG AGG TTC CAC ATG AGC ATG ACT GAG GAG CAG CTG CAG CTG CTG 2517  
 Lys Glu Arg Phe His Met Ser Met Thr Glu Glu Gln Leu Gln Leu Leu  
 790 795 800  
 GTG GAG CAG ATG GTG GAT GGC AGT ATG OGG TCT ATC ACC ACC AAA CTC 2565  
 Val Glu Gln Met Val Asp Gly Ser Met Arg Ser Ile Thr Thr Lys Leu  
 805 810 815  
 20 TAT GAC GGC TTC CAG TAC CTC ACC AAC GGC ATC ATG TGA CACGCTOCTC 2614  
 Tyr Asp Gly Phe Gln Tyr Leu Thr Asn Gly Ile Met \*  
 820 825 830  
 25 AGOCCAGGAG TGGTGGGGGG TOCAGGGCAC OCTOCTAGA GGGCOCTTGT CTGAGAAACC 2674  
 CCAAACCAGG AAAOCCCAOC TAOCCAAOCA TOCACCAAG GGAAATGGAA GGCAAGAAAC 2734  
 30 ACGAAGGATC ATGTGGTAAC TGOAGAGAGCT TGCTGAGGGG TGGGAGAGCC AGCTGTGGGG 2794  
 TOCAGACTTG TTGGGGCTTC OCTGCOOCTC CTGGTCTGTG TCAGTATTAC CAOCAGACTG 2854  
 ACTOCAGGAC TCACTGCOCT OCAGAAAACA GAGGTGACAA ATGTGAGGGA CACTGGGGOC 2914  
 35 TTCTCTCTOC TTGTAGGGGT CTCTCAGAGG TTCTTTCCAC AGGOCATCCT CTTATTCCGT 2974  
 TCTGGGGOCC AGGAAGTGGG GAAGAGTAGG TTCTCGGTAC TTAGGACTTG ATCCTGTGGT 3034  
 TGCCACTGGC CATGCTGCTG CCAGCTCTA CCOCTOCCAG GGAOCTACCC CTOCCAGGGA 3094  
 40 CCGACCOCTG GCOCAAGCTC CCTTGCTG GGGGOGCTGC GTGGGCOCTG CACTTGCTGA 3154  
 GGTTOOCCAT CATGGGCAAG GCAAGGGAAT TOCCACAGOC CTCAGTGTA CTGAGGGTAC 3214  
 45 TGGOCTAGCC ATGTGGAATT COCTACCOCTG ACTCCTTOOC CAAACCCAGG GAAAAGAGCT 3274  
 CTCAATTTTT TATTTTTAAT TTTTGTTGA AATAAAGTCC TTAGTTAGCC 3324

50 (2) INFORMATION FOR SEQ ID NO:34:

55

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 810 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:34:

Met Pro Met Asp Leu Ile Leu Val Val Trp Phe Cys Val Cys Thr Ala  
 1 5 10 15  
 Arg Thr Val Val Gly Phe Gly Met Asp Pro Asp Leu Gln Met Asp Ile  
 20 25 30  
 Val Thr Glu Leu Asp Leu Val Asn Thr Thr Leu Gly Val Ala Gln Val  
 35 40 45  
 Ser Gly Met His Asn Ala Ser Lys Ala Phe Leu Phe Gln Asp Ile Glu  
 50 55 60  
 Arg Glu Ile His Ala Ala Pro His Val Ser Glu Lys Leu Ile Gln Leu  
 65 70 75 80  
 Phe Gln Asn Lys Ser Glu Phe Thr Ile Leu Ala Thr Val Gln Gln Lys  
 85 90 95  
 Pro Ser Thr Ser Gly Val Ile Leu Ser Ile Arg Glu Leu Glu His Ser  
 100 105 110  
 Tyr Phe Glu Leu Glu Ser Ser Gly Leu Arg Asp Glu Ile Arg Tyr His  
 115 120 125  
 Tyr Ile His Asn Gly Lys Pro Arg Thr Glu Ala Leu Pro Tyr Arg Met  
 130 135 140  
 Ala Asp Gly Gln Trp His Lys Val Ala Leu Ser Val Ser Ala Ser His  
 145 150 155 160  
 Leu Leu Leu His Val Asp Cys Asn Arg Ile Tyr Glu Arg Val Ile Asp  
 165 170 175  
 Pro Pro Asp Thr Asn Leu Pro Pro Gly Ile Asn Leu Trp Leu Gly Gln  
 180 185 190  
 Arg Asn Gln Lys His Gly Leu Phe Lys Gly Ile Ile Gln Asp Gly Lys  
 195 200 205  
 Ile Ile Phe Met Pro Asn Gly Tyr Ile Thr Gln Cys Pro Asn Leu Asn

55

Gly Ser Gly Gln His Asn Cys Asp Glu Asn Ala Ile Cys Thr Asn Thr  
 485 490 495  
 5 Val Gln Gly His Ser Cys Thr Cys Lys Pro Gly Tyr Val Gly Asn Gly  
 500 505 510  
 Thr Ile Cys Arg Ala Phe Cys Glu Gly Cys Arg Tyr Gly Gly Thr  
 515 520 525  
 10 Cys Val Ala Pro Asn Lys Cys Val Cys Pro Ser Gly Phe Thr Gly Ser  
 530 535 540  
 His Cys Glu Lys Asp Ile Asp Glu Cys Ser Glu Gly Ile Ile Glu Cys  
 545 550 555 560  
 His Asn His Ser Arg Cys Val Asn Leu Pro Gly Trp Tyr His Cys Glu  
 565 570 575  
 20 Cys Arg Ser Gly Phe His Asp Asp Gly Thr Tyr Ser Leu Ser Gly Glu  
 580 585 590  
 Ser Cys Ile Asp Ile Asp Glu Cys Ala Leu Arg Thr His Thr Cys Trp  
 595 600 605  
 25 Asn Asp Ser Ala Cys Ile Asn Leu Ala Gly Gly Phe Asp Cys Leu Cys  
 610 615 620  
 Pro Ser Gly Pro Ser Cys Ser Gly Asp Cys Pro His Glu Gly Gly Leu  
 625 630 635 640  
 Lys His Asn Gly Gln Val Trp Thr Leu Lys Glu Asp Arg Cys Ser Val  
 645 650 655  
 35 Cys Ser Cys Lys Asp Gly Lys Ile Phe Cys Arg Arg Thr Ala Cys Asp  
 660 665 670  
 Cys Gln Asn Pro Ser Ala Asp Leu Phe Cys Cys Pro Glu Cys Asp Thr  
 675 680 685  
 40 Arg Val Thr Ser Gln Cys Leu Asp Gln Asn Gly His Lys Leu Tyr Arg  
 690 695 700  
 Ser Gly Asp Asn Trp Thr His Ser Cys Gln Gln Cys Arg Cys Leu Glu  
 705 710 715 720  
 Gly Glu Val Asp Cys Trp Pro Leu Thr Cys Pro Asn Leu Ser Cys Glu  
 725 730 735  
 50 Tyr Thr Ala Ile Leu Glu Gly Glu Cys Cys Pro Arg Cys Val Ser Asp  
 740 745 750  
 55



Pro Cys Leu Ala Asp Asn Ile Thr Tyr Asp Ile Arg Lys Thr Cys Leu  
755 760 765

5 Asp Ser Tyr Gly Val Ser Arg Leu Ser Gly Ser Val Trp Thr Met Ala  
770 775 780

Gly Ser Pro Cys Thr Thr Cys Lys Cys Lys Asn Gly Arg Val Cys Cys  
785 790 795 800

10 Ser Val Asp Phe Glu Cys Leu Gln Asn Asn  
805 810

15 (2) INFORMATION FOR SEQ ID NO:35:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 2430 base pairs

(B) TYPE: nucleic acid

20 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA(genomic)

25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:35:

ATGOOGATGG ATTTGATTTT AGTTGTGTGG TTCTGTGTGT GCACTGOCAG GACAGTGGTG 60  
30 GGCTTTGGGA TGGACCTGA OCTTCAGATG GATATCGTCA CCGAGCTTGA OCTTGTGAAC 120  
AOCACCTTG GAGTTGCTCA GGTGTCTGGA ATGCACAATG CCAGCAAAGC ATTTTATTTT 180  
CAAGACATAG AAAGAGAGAT OCATGCAGCT OCTCATGTGA GTGAGAAATT AATTCAGCTG 240  
35 TTOCAGAACA AGAGTGAATT CACCATTTTG GCCACTGTAC AGCAGAAGOC ATOCACCTCA 300  
GGAGTGATAC TGTOCATTCG AGAACTGGAG CACAGCTATT TTGAACTGGA GAGCAGTGGC 360  
40 CTGAGGGATG AGATTGGTA TCACTACATA CACAATGGGA AGCCAAGGAC AGAGGCACTT 420  
OCTTAOCGA TGGCAGATGG ACAATGGCAC AAGGTTGCAC TGTCAGTTAG CGCTCTCAT 480  
CTCCTGCTCC ATGTGACTG TAACAGGATT TATGAGCGTG TGATAGACC TOCAGATAOC 540  
45 AAOCTTCCC CAGGAATCAA TTTATGGCTT GGCCAGCGCA ACCAAAAGCA TGGCTTATTC 600  
AAAGGGATCA TOCAAGATGG GAAGATCATC TTTATGCOGA ATGGATATAT AACACAGTGT 660  
50 CCAAATCTAA ATCACACTTG CCAACCTGC AGTGATTTCT TAAGCCTGGT GCAAGGAATA 720

	ATGGATTTAC AAGAGCTTTT GGOCAAGATG ACTGCAAAAC TAAATTATGC AGAGACAAGA	780
5	CTTAGTCAAT TGGAAACTG TCATTGTGAG AAGACTTGTC AAGTGAGTGG ACTGCTCTAT	840
	CGAGATCAAG ACTCTTGGGT AGATGGTGAC CATTGCAGGA ACTGCACTTG CAAAAGTGGT	900
	GCGGTGGAAT GCGGAAGGAT GTCTGTGCC CCTCTCAATT GCTCCCCAGA CTCCCTCCCA	960
10	GTACACATTG CTGGCCAGTG CTGTAAGGTC TGOOGACCAA AATGTATCTA TGGAGGAAAA	1020
	GTCTTGCAG AAGGCCAGCG GATTTTAAOC AAGAGCTGTC GGAATGCOG AGGTGGAGTT	1080
15	TTAGTAAAAA TTACAGAAAT GTGTCTCCT TTGAACTGCT CAGAAAAGGA TCACATTCTT	1140
	CCTGAGAATC AGTGCTGCOG TGTCTGTAGA GGTCACTAAT TTTGTGCAGA AGGACCTAAA	1200
	TGTGGTGAAA ACTCAGAGTG CAAAACTGG AATACAAAAG CTACTTGTGA GTGCAAGAGT	1260
20	GGTTACATCT CTGTCCAGG AGACTCTGCC TACTGTGAAG ATATTGATGA GTGTGCAGCT	1320
	AAGATGCATT ACTGTCTATC CAATACTGTG TGTGTCAAOC TTCTGGGTT ATATGCTGT	1380
	GAATGTGTC CAGGATACAT TCGTGTGGAT GACTTCTCTT GTACAGAACA CGATGAATGT	1440
25	GGCAGCGGC AGCACAACCTG TGATGAGAAT GOCATCTGCA CCAACACTGT CCAGGGACAC	1500
	AGCTGCACT GCAAACCGG CTACGTGGGG AACGGGAOCA TCTGCAGAGC TTTCTGTGAA	1560
30	GAGGGCTGCA GATACGGTGG AACGTGTGTG GCTCCCAACA AATGTGTCTG TOCATCTGGA	1620
	TTACAGGAA GCACTGCGA GAAAGATATT GATGAATGTT CAGAGGGAAT CATTGAGTGC	1680
	CACAAOCATT CCGCTGGT TAACTGCA GGGTGGTAOC ACTGTGAGTG CAGAAGCGT	1740
35	TTCCATGACG ATGGGAOCTA TTCACTGTCC GGGGAGTCT GTATTGACAT TGATGAATGT	1800
	GOCTTAAGAA CTCACACTG TTGGAACGAT TCTGCCTGCA TCAACCTGGC AGGGGGTTTT	1860
40	GACTGTCTCT GCOCTCTGG GCOCTCTGC TCTGGTGAAT GTCTCATGA AGGGGGCTG	1920
	AAGCACAATG GCCAGGTGTG GACCTTGAAA GAAGACAGGT GTCTGTCTG CTCTGCAAG	1980
	GATGGCAAGA TATTCTGCOG ACGGACAGCT TGTGATTGOC AGAATOCAAG TGCTGACCTA	2040
45	TTCTGTGOC CAGAATGTGA CACCAGAGTC ACAAGTCAAT GTTTAGACCA AAATGGTCAC	2100
	AAGCTGTATC GAAGTGGAGA CAATTGGACC CATAGCTGTC AGCAGTGTG GTGTCTGGAA	2160
50	GGAGAGGTAG ATTGCTGGOC ACTCACTTGC CCAACTTGA GCTGTGAGTA TACAGCTATC	2220

55

TTAGAAGGGG AATGTTGTCC CCGCTGTGTC AGTGACCOCT GOCTAGCTGA TAACATCAACC 2280  
 TATGACATCA GAAAACTTG CCTGGACAGC TATGGTGTTC CACGGCTTAG TGGCTCAGTG 2340  
 TGGACGATGG CTGGATCTOC CTGCACAAOC TGTAAATGCA AGAATGGAAG AGTCTGTGTG 2400  
 TCTGTGGATT TTGAGTGTCT TCAAAATAAT 2430

(2) INFORMATION FOR SEQ ID NO:36:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2977 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA(genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vii) IMMEDIATE SOURCE:

- (A) LIBRARY: Human fetal brain cDNA library  
 (B) CLONE: GEN-073E07

(ix) FEATURE:

- (A) NAME/KEY: CDS  
 (B) LOCATION: 103..2532

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:36:

TAGCAAGTTT GGCGGCTOCA AGCCAGGGGC GOCTCAGGAT CCAGGCTCAT TTGCTTCAC 60  
 CTAGCTTOGG TGCCCCCTGC TAGGCGGGGA CCTOGAGAG CG ATG CCG ATG GAT 114  
 Met Pro Met Asp  
 1  
 TTG ATT TTA GTT GTG TGG TTC TGT GTG TGC ACT GGC AGG ACA GTG GTG 162  
 Leu Ile Leu Val Val Trp Phe Cys Val Cys Thr Ala Arg Thr Val Val  
 5 10 15 20  
 GGC TTT GGG ATG GAC CCT GAC CTT CAG ATG GAT ATC GTC ACC GAG CTT 210  
 Gly Phe Gly Met Asp Pro Asp Leu Gln Met Asp Ile Val Thr Glu Leu  
 25 30 35  
 GAC CTT GTG AAC ACC ACC CTT GGA GTT GCT CAG GTG TCT GGA ATG CAC 258  
 Asp Leu Val Asn Thr Thr Leu Gly Val Ala Gln Val Ser Gly Met His

	40	45	50	
5	AAT GOC AGC AAA GCA TTT TTA TTT CAA GAC ATA GAA AGA GAG ATC CAT Asn Ala Ser Lys Ala Phe Leu Phe Gln Asp Ile Glu Arg Glu Ile His 55 60 65			306
10	GCA GCT OCT CAT GTG AGT GAG AAA TTA ATT CAG CTG TTC CAG AAC AAG Ala Ala Pro His Val Ser Glu Lys Leu Ile Gln Leu Phe Gln Asn Lys 70 75 80			354
15	AGT GAA TTC ACC ATT TTG GOC ACT GTA CAG CAG AAG CCA TCC ACT TCA Ser Glu Phe Thr Ile Leu Ala Thr Val Gln Gln Lys Pro Ser Thr Ser 85 90 95 100			402
	GGA GTG ATA CTG TCC ATT CGA GAA CTG GAG CAC AGC TAT TTT GAA CTG Gly Val Ile Leu Ser Ile Arg Glu Leu Glu His Ser Tyr Phe Glu Leu 105 110 115			450
20	GAG AGC AGT GGC CTG AGG GAT GAG ATT CGG TAT CAC TAC ATA CAC AAT Glu Ser Ser Gly Leu Arg Asp Glu Ile Arg Tyr His Tyr Ile His Asn 120 125 130			498
25	GGG AAG CCA AGG ACA GAG GCA CTT OCT TAC CGC ATG GCA GAT GGA CAA Gly Lys Pro Arg Thr Glu Ala Leu Pro Tyr Arg Met Ala Asp Gly Gln 135 140 145			546
30	TGG CAC AAG GTT GCA CTG TCA GTT AGC GOC TCT CAT CTC CTG CTC CAT Trp His Lys Val Ala Leu Ser Val Ser Ala Ser His Leu Leu Leu His 150 155 160			594
	GTC GAC TGT AAC AGG ATT TAT GAG CGT GTG ATA GAC OCT CCA GAT ACC Val Asp Cys Asn Arg Ile Tyr Glu Arg Val Ile Asp Pro Pro Asp Thr 165 170 175 180			642
35	AAC CTT CCC CCA GGA ATC AAT TTA TGG CTT GGC CAG CGC AAC CAA AAG Asn Leu Pro Pro Gly Ile Asn Leu Trp Leu Gly Gln Arg Asn Gln Lys 185 190 195			690
40	CAT GGC TTA TTC AAA GGG ATC ATC CAA GAT GGG AAG ATC ATC TTT ATG His Gly Leu Phe Lys Gly Ile Ile Gln Asp Gly Lys Ile Ile Phe Met 200 205 210			738
45	COG AAT GGA TAT ATA ACA CAG TGT OCA AAT CTA AAT CAC ACT TGC OCA Pro Asn Gly Tyr Ile Thr Gln Cys Pro Asn Leu Asn His Thr Cys Pro 215 220 225			786
50	ACC TGC AGT GAT TTC TTA AGC CTG GTG CAA GGA ATA ATG GAT TTA CAA Thr Cys Ser Asp Phe Leu Ser Leu Val Gln Gly Ile Met Asp Leu Gln 230 235 240			834

5	GAG CTT TTG GCC AAG ATG ACT GCA AAA CTA AAT TAT GCA GAG ACA AGA Glu Leu Leu Ala Lys Met Thr Ala Lys Leu Asn Tyr Ala Glu Thr Arg 245 250 255 260	882
10	CTT AGT CAA TTG GAA AAC TGT CAT TGT GAG AAG ACT TGT CAA GTG AGT Leu Ser Gln Leu Glu Asn Cys His Cys Glu Lys Thr Cys Gln Val Ser 265 270 275	930
15	GGA CTG CTC TAT CGA GAT CAA GAC TCT TGG GTA GAT GGT GAC CAT TGC Gly Leu Leu Tyr Arg Asp Gln Asp Ser Trp Val Asp Gly Asp His Cys 280 285 290	978
20	AGG AAC TGC ACT TGC AAA AGT GGT GCC GTG GAA TGC CGA AGG ATG TCC Arg Asn Cys Thr Cys Lys Ser Gly Ala Val Glu Cys Arg Arg Met Ser 295 300 305	1026
25	TGT CCC OCT CTC AAT TGC TCC CCA GAC TCC CTC CCA GTA CAC ATT GCT Cys Pro Pro Leu Asn Cys Ser Pro Asp Ser Leu Pro Val His Ile Ala 310 315 320	1074
30	GGC CAG TGC TGT AAG GTC TGC CGA CCA AAA TGT ATC TAT GGA GGA AAA Gly Gln Cys Cys Lys Val Cys Arg Pro Lys Cys Ile Tyr Gly Gly Lys 325 330 335 340	1122
35	GTT CTT GCA GAA GGC CAG CGG ATT TTA ACC AAG AGC TGT CGG GAA TGC Val Leu Ala Glu Gly Gln Arg Ile Leu Thr Lys Ser Cys Arg Glu Cys 345 350 355	1170
40	CGA GGT GGA GTT TTA GTA AAA ATT ACA GAA ATG TGT OCT OCT TTG AAC Arg Gly Gly Val Leu Val Lys Ile Thr Glu Met Cys Pro Pro Leu Asn 360 365 370	1218
45	TGC TCA GAA AAG GAT CAC ATT CTT OCT GAG AAT CAG TGC TGC CGT GTC Cys Ser Glu Lys Asp His Ile Leu Pro Glu Asn Gln Cys Cys Arg Val 375 380 385	1266
50	TGT AGA GGT CAT AAC TTT TGT GCA GAA GGA CCT AAA TGT GGT GAA AAC Cys Arg Gly His Asn Phe Cys Ala Glu Gly Pro Lys Cys Gly Glu Asn 390 395 400	1314
55	TCA GAG TGC AAA AAC TGG AAT ACA AAA GCT ACT TGT GAG TGC AAG AGT Ser Glu Cys Lys Asn Trp Asn Thr Lys Ala Thr Cys Glu Cys Lys Ser 405 410 415 420	1362
60	GGT TAC ATC TCT GTC CAG GGA GAC TCT GGC TAC TGT GAA GAT ATT GAT Gly Tyr Ile Ser Val Gln Gly Asp Ser Ala Tyr Cys Glu Asp Ile Asp 425 430 435	1410
65	GAG TGT GCA GCT AAG ATG CAT TAC TGT CAT GCC AAT ACT GTG TGT GTC Glu Cys Ala Ala Lys Met His Tyr Cys His Ala Asn Thr Val Cys Val 440 445 450 455	1458

	440	445	450	
5	AAC CTT OCT GGG TTA TAT CGC TGT GAC TGT GTC OCA GGA TAC ATT CGT Asn Leu Pro Gly Leu Tyr Arg Cys Asp Cys Val Pro Gly Tyr Ile Arg 455 460 465			1506
10	GTG GAT GAC TTC TCT TGT ACA GAA CAC GAT GAA TGT GGC AGC GGC CAG Val Asp Asp Phe Ser Cys Thr Glu His Asp Glu Cys Gly Ser Gly Gln 470 475 480			1554
15	CAC AAC TGT GAT GAG AAT GGC ATC TGC ACC AAC ACT GTC CAG GGA CAC His Asn Cys Asp Glu Asn Ala Ile Cys Thr Asn Thr Val Gln Gly His 485 490 495 500			1602
	AGC TGC ACC TGC AAA CCG GGC TAC GTG GGG AAC GGG ACC ATC TGC AGA Ser Cys Thr Cys Lys Pro Gly Tyr Val Gly Asn Gly Thr Ile Cys Arg 505 510 515			1650
20	GCT TTC TGT GAA GAG GGC TGC AGA TAC GGT GGA ACG TGT GTG GCT CCC Ala Phe Cys Glu Glu Gly Cys Arg Tyr Gly Gly Thr Cys Val Ala Pro 520 525 530			1698
25	AAC AAA TGT GTC TGT CCA TCT GGA TTC ACA GGA AGC CAC TGC GAG AAA Asn Lys Cys Val Cys Pro Ser Gly Phe Thr Gly Ser His Cys Glu Lys 535 540 545			1746
30	GAT ATT GAT GAA TGT TCA GAG GGA ATC ATT GAG TGC CAC AAC CAT TOC Asp Ile Asp Glu Cys Ser Glu Gly Ile Ile Glu Cys His Asn His Ser 550 555 560			1794
35	CGC TGC GTT AAC CTG CCA GGG TGG TAC CAC TGT GAG TGC AGA AGC GGT Arg Cys Val Asn Leu Pro Gly Trp Tyr His Cys Glu Cys Arg Ser Gly 565 570 575 580			1842
	TTC CAT GAC GAT GGG AOC TAT TCA CTG TOC GGG GAG TOC TGT ATT GAC Phe His Asp Asp Gly Thr Tyr Ser Leu Ser Gly Glu Ser Cys Ile Asp 585 590 595			1890
40	ATT GAT GAA TGT GCC TTA AGA ACT CAC ACC TGT TGG AAC GAT TCT GCC Ile Asp Glu Cys Ala Leu Arg Thr His Thr Cys Trp Asn Asp Ser Ala 600 605 610			1938
45	TGC ATC AAC CTG GCA GGG GGT TTT GAC TGT CTC TGC OCC TCT GGG OCC Cys Ile Asn Leu Ala Gly Gly Phe Asp Cys Leu Cys Pro Ser Gly Pro 615 620 625			1986
50	TOC TGC TCT GGT GAC TGT OCT CAT GAA GGG GGG CTG AAG CAC AAT GGC Ser Cys Ser Gly Asp Cys Pro His Glu Gly Gly Leu Lys His Asn Gly 630 635 640			2034

55

5	CAG GTG TGG ACC TTG AAA GAA GAC AGG TGT TCT GTC TGC TCC TGC AAG Gln Val Trp Thr Leu Lys Glu Asp Arg Cys Ser Val Cys Ser Cys Lys 645 650 655 660	2082
	GAT GGC AAG ATA TTC TGC CGA CGG ACA GCT TGT GAT TGC CAG AAT CCA Asp Gly Lys Ile Phe Cys Arg Arg Thr Ala Cys Asp Cys Gln Asn Pro 665 670 675	2130
10	AGT GCT GAC CTA TTC TGT TGC CCA GAA TGT GAC ACC AGA GTC ACA AGT Ser Ala Asp Leu Phe Cys Cys Pro Glu Cys Asp Thr Arg Val Thr Ser 680 685 690	2178
15	CAA TGT TTA GAC CAA AAT GGT CAC AAG CTG TAT CGA AGT GGA GAC AAT Gln Cys Leu Asp Gln Asn Gly His Lys Leu Tyr Arg Ser Gly Asp Asn 695 700 705	2226
20	TGG ACC CAT AGC TGT CAG CAG TGT OGG TGT CTG GAA GGA GAG GTA GAT Trp Thr His Ser Cys Gln Gln Cys Arg Cys Leu Glu Gly Glu Val Asp 710 715 720	2274
25	TGC TGG CCA CTC ACT TGC CCC AAC TTG AGC TGT GAG TAT ACA GCT ATC Cys Trp Pro Leu Thr Cys Pro Asn Leu Ser Cys Glu Tyr Thr Ala Ile 725 730 735 740	2322
	TTA GAA GGG GAA TGT TGT CCC CGC TGT GTC AGT GAC CCC TGC CTA GCT Leu Glu Gly Glu Cys Cys Pro Arg Cys Val Ser Asp Pro Cys Leu Ala 745 750 755	2370
30	GAT AAC ATC ACC TAT GAC ATC AGA AAA ACT TGC CTG GAC AGC TAT GGT Asp Asn Ile Thr Tyr Asp Ile Arg Lys Thr Cys Leu Asp Ser Tyr Gly 760 765 770	2418
35	GTT TCA CCG CTT AGT GGC TCA GTG TGG ACG ATG GCT GGA TCT CCC TGC Val Ser Arg Leu Ser Gly Ser Val Trp Thr Met Ala Gly Ser Pro Cys 775 780 785	2466
40	ACA AOC TGT AAA TGC AAG AAT GGA AGA GTC TGT TGT TCT GTG GAT TTT Thr Thr Cys Lys Cys Lys Asn Gly Arg Val Cys Cys Ser Val Asp Phe 790 795 800	2514
	GAG TGT CTT CAA AAT AAT TGAAGTATTT ACAGTGGACT CAACGCAGAA Glu Cys Leu Gln Asn Asn 805 810	2562
45	GAATGGAAGA AATGAACATC CAACGTGATT AAGGATAGGA ATGGGTAGTT TGGTTTTTTTTT	2622
	GTTTGTTTTG TTTTTTTAAC CACAGATAAT TGCCAAAGTT TCCACCTGAG GACGGTGTTT	2682
50	CGGAGGTGTC CTTTTGGAOC TACCACTTTG CTCATTCTTG CTAACCTAGT CTAGGTGACC	2742
55		

TACAGTGOOG TGCATTTAAG TCAATGGTTG TTAAAAGAAG TTTCCCGTGT TGTAATCAT 2802  
 GTTTCOCTTA TCAGATCATT TGCAAATACA TTAAATGAT CTCATGGTAA ATGGTTGATG 2862  
 TATTTTTTGG GTTATTTTG TGTACTAACC ATAATAGAGA GAGACTCAGC TCCTTTTATT 2922  
 TATTTTGTG ATTTATGGAT CAAATTCTAA AATAAAGTTG OCTGTTGTA CTTTT 2977

(2) INFORMATION FOR SEQ ID NO:37:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 816 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:37:

Met Glu Ser Arg Val Leu Leu Arg Thr Phe Cys Leu Ile Phe Gly Leu  
 1 5 10 15  
 Gly Ala Val Trp Gly Leu Gly Val Asp Pro Ser Leu Gln Ile Asp Val  
 20 25 30  
 Leu Thr Glu Leu Glu Leu Gly Glu Ser Thr Thr Gly Val Arg Gln Val  
 35 40 45  
 Pro Gly Leu His Asn Gly Thr Lys Ala Phe Leu Phe Gln Asp Thr Pro  
 50 55 60  
 Arg Ser Ile Lys Ala Ser Thr Ala Thr Ala Glu Gln Phe Phe Gln Lys  
 65 70 75 80  
 Leu Arg Asn Lys His Glu Phe Thr Ile Leu Val Thr Leu Lys Gln Thr  
 85 90 95  
 His Leu Asn Ser Gly Val Ile Leu Ser Ile His His Leu Asp His Arg  
 100 105 110  
 Tyr Leu Glu Leu Glu Ser Ser Gly His Arg Asn Glu Val Arg Leu His  
 115 120 125  
 Tyr Arg Ser Gly Ser His Arg Pro His Thr Glu Val Phe Pro Tyr Ile  
 130 135 140  
 Leu Ala Asp Asp Lys Trp His Lys Leu Ser Leu Ala Ile Ser Ala Ser



	145		150		155		160
5	His Leu Ile Leu	His Ile Asp Cys Asn Lys	Ile Tyr Glu Arg Val Val				
		165		170		175	
	Glu Lys Pro Ser	Thr Asp Leu Pro Leu Gly Thr Thr Phe Trp Leu Gly					
		180		185		190	
10	Gln Arg Asn Asn Ala His Gly Tyr Phe Lys Gly Ile Met Gln Asp Val						
		195		200		205	
	Gln Leu Leu Val Met Pro Gln Gly Phe Ile Ala Gln Cys Pro Asp Leu						
15		210		215		220	
	Asn Arg Thr Cys Pro Thr Cys Asn Asp Phe His Gly Leu Val Gln Lys						
		225		230		235	
20	Ile Met Glu Leu Gln Asp Ile Leu Ala Lys Thr Ser Ala Lys Leu Ser						
		245		250		255	
	Arg Ala Glu Gln Arg Met Asn Arg Leu Asp Gln Cys Tyr Cys Glu Arg						
		260		265		270	
25	Thr Cys Thr Met Lys Gly Thr Thr Tyr Arg Glu Phe Glu Ser Trp Ile						
		275		280		285	
	Asp Gly Cys Lys Asn Cys Thr Cys Leu Asn Gly Thr Ile Gln Cys Glu						
30		290		295		300	
	Thr Leu Ile Cys Pro Asn Pro Asp Cys Pro Leu Lys Ser Ala Leu Ala						
		305		310		315	
35	Tyr Val Asp Gly Lys Cys Cys Lys Glu Cys Lys Ser Ile Cys Gln Phe						
		325		330		335	
	Gln Gly Arg Thr Tyr Phe Glu Gly Glu Arg Asn Thr Val Tyr Ser Ser						
		340		345		350	
40	Ser Gly Val Cys Val Leu Tyr Glu Cys Lys Asp Gln Thr Met Lys Leu						
		355		360		365	
	Val Glu Ser Ser Gly Cys Pro Ala Leu Asp Cys Pro Glu Ser His Gln						
45		370		375		380	
	Ile Thr Leu Ser His Ser Cys Cys Lys Val Cys Lys Gly Tyr Asp Phe						
		385		390		395	
50	Cys Ser Glu Arg His Asn Cys Met Glu Asn Ser Ile Cys Arg Asn Leu						
		405		410		415	

55

Asn Asp Arg Ala Val Cys Ser Cys Arg Asp Gly Phe Arg Ala Leu Arg  
 420 425 430  
 5 Glu Asp Asn Ala Tyr Cys Glu Asp Ile Asp Glu Cys Ala Glu Gly Arg  
 435 440 445  
 His Tyr Cys Arg Glu Asn Thr Met Cys Val Asn Thr Pro Gly Ser Phe  
 450 455 460  
 10 Met Cys Ile Cys Lys Thr Gly Tyr Ile Arg Ile Asp Asp Tyr Ser Cys  
 465 470 475 480  
 Thr Glu His Asp Glu Cys Ile Thr Asn Gln His Asn Cys Asp Glu Asn  
 485 490 495  
 15 Ala Leu Cys Phe Asn Thr Val Gly Gly His Asn Cys Val Cys Lys Pro  
 500 505 510  
 Gly Tyr Thr Gly Asn Gly Thr Thr Cys Lys Ala Phe Cys Lys Asp Gly  
 515 520 525  
 Cys Arg Asn Gly Gly Ala Cys Ile Ala Ala Asn Val Cys Ala Cys Pro  
 530 535 540  
 25 Gln Gly Phe Thr Gly Pro Ser Cys Glu Thr Asp Ile Asp Glu Cys Ser  
 545 550 555 560  
 Asp Gly Phe Val Gln Cys Asp Ser Arg Ala Asn Cys Ile Asn Leu Pro  
 565 570 575  
 Gly Trp Tyr His Cys Glu Cys Arg Asp Gly Tyr His Asp Asn Gly Met  
 580 585 590  
 35 Phe Ser Pro Ser Gly Glu Ser Cys Glu Asp Ile Asp Glu Cys Gly Thr  
 595 600 605  
 Gly Arg His Ser Cys Ala Asn Asp Thr Ile Cys Phe Asn Leu Asp Gly  
 610 615 620  
 40 Gly Tyr Asp Cys Arg Cys Pro His Gly Lys Asn Cys Thr Gly Asp Cys  
 625 630 635 640  
 Ile His Asp Gly Lys Val Lys His Asn Gly Gln Ile Trp Val Leu Glu  
 645 650 655  
 Asn Asp Arg Cys Ser Val Cys Ser Cys Gln Asn Gly Phe Val Met Cys  
 660 665 670  
 50 Arg Arg Met Val Cys Asp Cys Glu Asn Pro Thr Val Asp Leu Phe Cys  
 675 680 685  
 55

Cys Pro Glu Cys Asp Pro Arg Leu Ser Ser Gln Cys Leu His Gln Asn  
 690 695 700  
 5 Gly Glu Thr Leu Tyr Asn Ser Gly Asp Thr Trp Val Gln Asn Cys Gln  
 705 710 715 720  
 Gln Cys Arg Cys Leu Gln Gly Glu Val Asp Cys Trp Pro Leu Pro Cys  
 725 730 735  
 10 Pro Asp Val Glu Cys Glu Phe Ser Ile Leu Pro Glu Asn Glu Cys Cys  
 740 745 750  
 15 Pro Arg Cys Val Thr Asp Pro Cys Gln Ala Asp Thr Ile Arg Asn Asp  
 755 760 765  
 Ile Thr Lys Thr Cys Leu Asp Glu Met Asn Val Val Arg Phe Thr Gly  
 770 775 780  
 20 Ser Ser Trp Ile Lys His Gly Thr Glu Cys Thr Leu Cys Gln Cys Lys  
 785 790 795 800  
 Asn Gly His Ile Cys Cys Ser Val Asp Pro Gln Cys Leu Gln Glu Leu  
 805 810 815  
 25

## (2) INFORMATION FOR SEQ ID NO:38:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2448 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA(genomic)

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:38:

40 ATGGAGTCTC GGGTCTTACT GAGAACATTC TGTTTGATCT TGGTCTCGG AGCAGTTTGG 60  
 GGGCTTGGTG TGGACCTTC OCTACAGATT GAAGTCTTAA CAGAGTTAGA ACTTGGGGAG 120  
 TOACGACCG GAGTGGGTCA GGTCCCGGG CTGCATAATG GGACGAAAGC CTTTCTCTTT 180  
 45 CAAGATACTC OCAGAAGCAT AAAAGCATCC ACTGCTACAG CTGAACAGTT TTTTCAGAAG 240  
 CTGAGAAATA AACATGAATT TACTATTTTG GTGACCTTAA AACAGACCA CTTAAATTCA 300  
 50 GGAGTTATTC TCTCAATTCA CCACTTGGAT CACAGGTACC TGGAACTGGA AAGTAGTGGC 360

	CATCGGAATG AAGTCAGACT GCATTACCGC TCAGGCAGTC ACGGCOCTCA CACAGAAGTG	420
5	TTTCTTACA TTTTGGCTGA TGACAAGTGG CACAAGCTCT OCTTAGCCAT CAGTGCTTCC	480
	CATTTGATTT TACACATTGA CTGCAATAAA ATTTATGAAA GGGTAGTAGA AAAGCOCTCC	540
	ACAGACTTGC CTCTAGGCAC AACATTTTGG CTAGGACAGA GAAATAATGC GCATGGATAT	600
10	TTTAAGGGTA TAATGCAAGA TGTCOAATTA CTTGTTCATGC CCGAGGGATT TATTGCTCAG	660
	TGCCCAGATC TTAATCGCAC CTGTCCAATC TGCAATGACT TOCATGGACT TGTGCAGAAA	720
15	ATCATGGAGC TACAGGATAT TTTAGCCAAA ACATCAGCCA AGCTGTCTCG AGCTGAACAG	780
	CGAATGAATA GATTGGATCA GTGCTATTGT GAAAGGACTT GCAOCATGAA GGGAAOCACC	840
	TACCGAGAAT TTGAGTCTCG GATAGACGGC TGTAAGAACT GCACATGCCT GAATGGAACC	900
20	ATOCAGTGTG AAACCTCTAAT CTGCCCCAAT OCTGACTGCC CACTTAAGTC GGCTCTTGCG	960
	TATGTGGATG GCAAATGCTG TAAGGAATGC AAATCGATAT GCCAATTTCA AGGAOGAACC	1020
25	TACTTTGAAG GAGAAAGAAA TACAGTCTAT TOCTCTTCTG GAGTATGTGT TCTCTATGAG	1080
	TGCAAGGACC AGACCATGAA ACTTGTGAG AGTTCAGGCT GTCCAGCTTT GGATTGTOCA	1140
	GAGTCTCATC AGATAAOCCT GTCTCACAGC TGTTGCAAAG TTTGTAAAGG TTATGACTTT	1200
30	TGTTCTGAAA GGCATAACTG CATGGAGAAT TOCATCTGCA GAAATCTGAA TGACAGGGCT	1260
	GTTTGTAGCT GTCGAGATGG TTTTAGGGCT CTTGAGAGG ATAATGCCTA CTGTGAAGAC	1320
35	ATOGATGAGT GTGCTGAAGG GCGCCATTAC TGTCGTGAAA ATACAATGTG TGTCAACACC	1380
	CCGGGTCTTT TTATGTGCAT CTGCAAAACT GGATACATCA GAATTGATGA TTATTCATGT	1440
	ACAGAACATG ATGAGTGTAT CACAAATCAG CACAACGTG ATGAAAATGC TTTATGCTTC	1500
40	AACACTGTTG GAGGACACAA CTGTGTTTGC AAGCCGGGCT ATACAGGGAA TGGAAOGACA	1560
	TGCAAAGCAT TTTGCAAAGA TGGCTGTAGG AATGGAGGAG CCTGTATTGC CGCTAATGTG	1620
	TGTGCTGCC CACAAGGCTT CACTGGACCC AGCTGTGAAA CGGACATTGA TGAATGCTCT	1680
45	GATGGTTTTG TTCAATGTGA CAGTCGTGCT AATTGCATTA AACTGCTGG ATGGTAOCAC	1740
	TGTGAGTGCA GAGATGGCTA CCATGACAAT GGGATGTTTT CACCAAGTGG AGAATCGTGT	1800
50	GAAGATATTG ATGAGTGTGG GACCGGGAGG CACAGCTGTG CCAATGATAC CATTTGCTTC	1860

55

AATTTGGATG GGGGATATGA TTGTGATGT OCTCATGGAA AGAATTGCAC AGGGGACTGC 1920  
 5 ATCCATGATG GAAAAGTTAA GCACAATGGT CAGATTTGGG TGTGGGAAA TGACAGGTGC 1980  
 TCTGTGTGCT CATGTCAGAA TGGATTGGTT ATGTGTGAC GGATGGTCTG TGAATGTGAG 2040  
 AATCCACAG TTGATCTTTT TTGCTGCOCT GAATGTGAOC CAAGGCTTAG TAGTCAGTGC 2100  
 10 CTCCATCAA ATGGGGAAAC TTGTATATAAC AGTGGTGACA CCTGGGTCCA GAATTGTCAA 2160  
 CAGTGCCGCT GCTTGCAAGG GGAAGTTGAT TGTGGGCOOC TGCOCTGCOOC AGATGTGGAG 2220  
 TGTGAATTCA GCATTCTOOC AGAGAATGAG TGCTGCOOGC GCTGTGTAC AGACCOCTGC 2280  
 15 CAGGCTGACA CCATCCGCAA TGACATCAOC AAGACTTGOC TGGACGAAAT GAATGTGGTT 2340  
 CGCTTCACCG GGTCTCTTG GATCAAACAT GGCATGAGT GTACTCTCTG CCAGTGCAAG 2400  
 20 AATGGCCACA TCTGTGCTC AGTGGATCCA CAGTGCOCTC AGGAAGT 2448

## (2) INFORMATION FOR SEQ ID NO:39:

- 25 (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 3198 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear  
 30 (ii) MOLECULE TYPE: DNA(genomic)  
 (iii) HYPOTHETICAL: NO  
 35 (iv) ANTI-SENSE: NO  
 (vii) IMMEDIATE SOURCE:  
 (A) LIBRARY: Human fetal brain cDNA library  
 (B) CLONE: GEN-093E05  
 40 (ix) FEATURE:  
 (A) NAME/KEY: CDS  
 (B) LOCATION: 97..2544  
 45 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:39:

TTGGGAGGAG CAGTCTCTCC GCTGTCTCC CGGAGCTTTC TCATTTGTCT CTGCOCTTAC 60  
 50 AACAGAGGGA GACGATGGAC TGAGCTGATC CGCAAC ATG GAG TCT CGG GTC TTA 114  
 Met Glu Ser Arg Val Leu

55

5

5	CTG	AGA	ACA	TTC	TGT	TTG	ATC	TTC	GGT	CTC	GGA	GCA	GTT	TGG	GGG	CTT	162
	Leu	Arg	Thr	Phe	Cys	Leu	Ile	Phe	Gly	Leu	Gly	Ala	Val	Trp	Gly	Leu	
				10					15					20			
	GGT	GTG	GAC	OCT	TOC	CTA	CAG	ATT	GAC	GTC	TTA	ACA	GAG	TTA	GAA	CTT	210
10	Gly	Val	Asp	Pro	Ser	Leu	Gln	Ile	Asp	Val	Leu	Thr	Glu	Leu	Glu	Leu	
			25					30					35				
	GGG	GAG	TOC	ACG	ACC	GGA	GTG	CGT	CAG	GTC	CCG	GGG	CTG	CAT	AAT	GGG	258
	Gly	Glu	Ser	Thr	Thr	Gly	Val	Arg	Gln	Val	Pro	Gly	Leu	His	Asn	Gly	
		40					45					50					
15	ACG	AAA	GOC	TTT	CTC	TTT	CAA	GAT	ACT	CCC	AGA	AGC	ATA	AAA	GCA	TOC	306
	Thr	Lys	Ala	Phe	Leu	Phe	Gln	Asp	Thr	Pro	Arg	Ser	Ile	Lys	Ala	Ser	
	55				60					65						70	
20	ACT	GCT	ACA	GCT	GAA	CAG	TTT	TTT	CAG	AAG	CTG	AGA	AAT	AAA	CAT	GAA	354
	Thr	Ala	Thr	Ala	Glu	Gln	Phe	Phe	Gln	Lys	Leu	Arg	Asn	Lys	His	Glu	
				75					80						85		
	TTT	ACT	ATT	TTG	GTG	AOC	CTA	AAA	CAG	AOC	CAC	TTA	AAT	TCA	GGA	GTT	402
25	Phe	Thr	Ile	Leu	Val	Thr	Leu	Lys	Gln	Thr	His	Leu	Asn	Ser	Gly	Val	
			90						95					100			
	ATT	CTC	TCA	ATT	CAC	CAC	TTG	GAT	CAC	AGG	TAC	CTG	GAA	CTG	GAA	AGT	450
30	Ile	Leu	Ser	Ile	His	His	Leu	Asp	His	Arg	Tyr	Leu	Glu	Leu	Glu	Ser	
		105						110					115				
	AGT	GGC	CAT	OGG	AAT	GAA	GTG	AGA	CTG	CAT	TAC	CGC	TCA	GGC	AGT	CAC	498
	Ser	Gly	His	Arg	Asn	Glu	Val	Arg	Leu	His	Tyr	Arg	Ser	Gly	Ser	His	
		120					125					130					
35	OGC	OCT	CAC	ACA	GAA	GTG	TTT	OCT	TAC	ATT	TTG	GCT	GAT	GAC	AAG	TGG	546
	Arg	Pro	His	Thr	Glu	Val	Phe	Pro	Tyr	Ile	Leu	Ala	Asp	Asp	Lys	Trp	
	135				140						145					150	
40	CAC	AAG	CTC	TOC	TTA	GOC	ATC	AGT	GCT	TOC	CAT	TTG	ATT	TTA	CAC	ATT	594
	His	Lys	Leu	Ser	Leu	Ala	Ile	Ser	Ala	Ser	His	Leu	Ile	Leu	His	Ile	
				155					160						165		
	GAC	TGC	AAT	AAA	ATT	TAT	GAA	AGG	GTA	GTA	GAA	AAG	CCC	TOC	ACA	GAC	642
45	Asp	Cys	Asn	Lys	Ile	Tyr	Glu	Arg	Val	Val	Glu	Lys	Pro	Ser	Thr	Asp	
			170					175						180			
	TTG	OCT	CTA	GGC	ACA	ACA	TTT	TGG	CTA	GGA	CAG	AGA	AAT	AAT	GCG	CAT	690
50	Leu	Pro	Leu	Gly	Thr	Thr	Phe	Trp	Leu	Gly	Gln	Arg	Asn	Asn	Ala	His	
			185					190					195				

5	GGA TAT TTT AAG GGT ATA ATG CAA GAT GTC CAA TTA CTT GTC ATG CCC Gly Tyr Phe Lys Gly Ile Met Gln Asp Val Gln Leu Leu Val Met Pro 200 205 210	738
10	CAG GGA TTT ATT GCT CAG TGC CCA GAT CTT AAT CGC ACC TGT CCA ACT Gln Gly Phe Ile Ala Gln Cys Pro Asp Leu Asn Arg Thr Cys Pro Thr 215 220 225 230	786
15	TGC AAT GAC TTC CAT GGA CTT GTG CAG AAA ATC ATG GAG CTA CAG GAT Cys Asn Asp Phe His Gly Leu Val Gln Lys Ile Met Glu Leu Gln Asp 235 240 245	834
20	ATT TTA GCC AAA ACA TCA GGC AAG CTG TCT CGA GCT GAA CAG CGA ATG Ile Leu Ala Lys Thr Ser Ala Lys Leu Ser Arg Ala Glu Gln Arg Met 250 255 260	882
25	AAT AGA TTG GAT CAG TGC TAT TGT GAA AGG ACT TGC ACC ATG AAG GGA Asn Arg Leu Asp Gln Cys Tyr Cys Glu Arg Thr Cys Thr Met Lys Gly 265 270 275	930
30	ACC ACC TAC CGA GAA TTT GAG TCC TGG ATA GAC GGC TGT AAG AAC TGC Thr Thr Tyr Arg Glu Phe Glu Ser Trp Ile Asp Gly Cys Lys Asn Cys 280 285 290	978
35	ACA TGC CTG AAT GGA ACC ATC CAG TGT GAA ACT CTA ATC TGC CCA AAT Thr Cys Leu Asn Gly Thr Ile Gln Cys Glu Thr Leu Ile Cys Pro Asn 295 300 305 310	1026
40	CGT GAC TGC CCA CTT AAG TCG GCT CTT GCG TAT GTG GAT GGC AAA TGC Pro Asp Cys Pro Leu Lys Ser Ala Leu Ala Tyr Val Asp Gly Lys Cys 315 320 325	1074
45	TGT AAG GAA TGC AAA TCG ATA TGC CAA TTT CAA GGA CGA ACC TAC TTT Cys Lys Glu Cys Lys Ser Ile Cys Gln Phe Gln Gly Arg Thr Tyr Phe 330 335 340	1122
50	GAA GGA GAA AGA AAT ACA GTC TAT TCC TCT TCT GGA GTA TGT GTT CTC Glu Gly Glu Arg Asn Thr Val Tyr Ser Ser Ser Gly Val Cys Val Leu 345 350 355	1170
55	TAT GAG TGC AAG GAC CAG ACC ATG AAA CTT GTT GAG AGT TCA GGC TGT Tyr Glu Cys Lys Asp Gln Thr Met Lys Leu Val Glu Ser Ser Gly Cys 360 365 370	1218
60	CCA GCT TTG GAT TGT CCA GAG TCT CAT CAG ATA ACC TTG TCT CAC AGC Pro Ala Leu Asp Cys Pro Glu Ser His Gln Ile Thr Leu Ser His Ser 375 380 385 390	1266
65	TGT TGC AAA GTT TGT AAA GGT TAT GAC TTT TGT TCT GAA AGG CAT AAC Cys Cys Lys Val Cys Lys Gly Tyr Asp Phe Cys Ser Glu Arg His Asn 395 400 405 410	1314

	395										400					405					
5	TGC	ATG	GAG	AAT	TCC	ATC	TGC	AGA	AAT	CTG	AAT	GAC	AGG	GCT	GTT	TGT	1362				
	Cys	Met	Glu	Asn	Ser	Ile	Cys	Arg	Asn	Leu	Asn	Asp	Arg	Ala	Val	Cys					
				410					415					420							
10	AGC	TGT	CGA	GAT	GGT	TTT	AGG	GCT	CTT	CGA	GAG	GAT	AAT	GCC	TAC	TGT	1410				
	Ser	Cys	Arg	Asp	Gly	Phe	Arg	Ala	Leu	Arg	Glu	Asp	Asn	Ala	Tyr	Cys					
				425				430					435								
15	GAA	GAC	ATC	GAT	GAG	TGT	GCT	GAA	GGG	CGC	CAT	TAC	TGT	CGT	GAA	AAT	1458				
	Glu	Asp	Ile	Asp	Glu	Cys	Ala	Glu	Gly	Arg	His	Tyr	Cys	Arg	Glu	Asn					
				440			445					450									
20	ACA	ATG	TGT	GTC	AAC	ACC	CCG	GGT	TCT	TTT	ATG	TGC	ATC	TGC	AAA	ACT	1506				
	Thr	Met	Cys	Val	Asn	Thr	Pro	Gly	Ser	Phe	Met	Cys	Ile	Cys	Lys	Thr					
						460					465					470					
25	GGA	TAC	ATC	AGA	ATT	GAT	GAT	TAT	TCA	TGT	ACA	GAA	CAT	GAT	GAG	TGT	1554				
	Gly	Tyr	Ile	Arg	Ile	Asp	Asp	Tyr	Ser	Cys	Thr	Glu	His	Asp	Glu	Cys					
					475					480					485						
30	ATC	ACA	AAT	CAG	CAC	AAC	TGT	GAT	GAA	AAT	GCT	TTA	TGC	TTC	AAC	ACT	1602				
	Ile	Thr	Asn	Gln	His	Asn	Cys	Asp	Glu	Asn	Ala	Leu	Cys	Phe	Asn	Thr					
				490					495				500								
35	GTT	GGA	GGA	CAC	AAC	TGT	GTT	TGC	AAG	CCG	GGC	TAT	ACA	GGG	AAT	GGA	1650				
	Val	Gly	Gly	His	Asn	Cys	Val	Cys	Lys	Pro	Gly	Tyr	Thr	Gly	Asn	Gly					
				505			510						515								
40	ACG	ACA	TGC	AAA	GCA	TTT	TGC	AAA	GAT	GGC	TGT	AGG	AAT	GGA	GGA	GCC	1698				
	Thr	Thr	Cys	Lys	Ala	Phe	Cys	Lys	Asp	Gly	Cys	Arg	Asn	Gly	Gly	Ala					
				520			525					530									
45	TGT	ATT	GCC	GCT	AAT	GTG	TGT	GCC	TGC	CCA	CAA	GGC	TTC	ACT	GGA	CCC	1746				
	Cys	Ile	Ala	Ala	Asn	Val	Cys	Ala	Cys	Pro	Gln	Gly	Phe	Thr	Gly	Pro					
						540					545				550						
50	AGC	TGT	GAA	ACG	GAC	ATT	GAT	GAA	TGC	TCT	GAT	GGT	TTT	GTT	CAA	TGT	1794				
	Ser	Cys	Glu	Thr	Asp	Ile	Asp	Glu	Cys	Ser	Asp	Gly	Phe	Val	Gln	Cys					
					555					560				565							
55	GAC	AGT	CGT	GCT	AAT	TGC	ATT	AAC	CTG	CCT	GGA	TGG	TAC	CAC	TGT	GAG	1842				
	Asp	Ser	Arg	Ala	Asn	Cys	Ile	Asn	Leu	Pro	Gly	Trp	Tyr	His	Cys	Glu					
				570					575					580							
60	TGC	AGA	GAT	GGC	TAC	CAT	GAC	AAT	GGG	ATG	TTT	TCA	CCA	AGT	GGA	GAA	1890				
	Cys	Arg	Asp	Gly	Tyr	His	Asp	Asn	Gly	Met	Phe	Ser	Pro	Ser	Gly	Glu					
				585				590					595								



5	TCG TGT GAA GAT ATT GAT GAG TGT GGG ACC GGG AGG CAC AGC TGT GGC Ser Cys Glu Asp Ile Asp Glu Cys Gly Thr Gly Arg His Ser Cys Ala 600 605 610	1938
10	AAT GAT ACC ATT TGC TTC AAT TTG GAT GGC GGA TAT GAT TGT OGA TGT Asn Asp Thr Ile Cys Phe Asn Leu Asp Gly Gly Tyr Asp Cys Arg Cys 615 620 625 630	1986
15	OCT CAT GGA AAG AAT TGC ACA GGG GAC TGC ATC CAT GAT GGA AAA GTT Pro His Gly Lys Asn Cys Thr Gly Asp Cys Ile His Asp Gly Lys Val 635 640 645	2034
20	AAG CAC AAT GGT CAG ATT TGG GTG TTG GAA AAT GAC AGG TGC TCT GTG Lys His Asn Gly Gln Ile Trp Val Leu Glu Asn Asp Arg Cys Ser Val 650 655 660	2082
25	TGC TCA TGT CAG AAT GGA TTC GTT ATG TGT OGA OGG ATG GTC TGT GAC Cys Ser Cys Gln Asn Gly Phe Val Met Cys Arg Arg Met Val Cys Asp 665 670 675	2130
30	TGT GAG AAT CCC ACA GTT GAT CTT TTT TGC TGC OCT GAA TGT GAC CCA Cys Glu Asn Pro Thr Val Asp Leu Phe Cys Cys Pro Glu Cys Asp Pro 680 685 690	2178
35	AGG CTT AGT AGT CAG TGC CTC CAT CAA AAT GGG GAA ACT TTG TAT AAC Arg Leu Ser Ser Gln Cys Leu His Gln Asn Gly Glu Thr Leu Tyr Asn 695 700 705 710	2226
40	AGT GGT GAC ACC TGG GTC CAG AAT TGT CAA CAG TGC CGC TGC TTG CAA Ser Gly Asp Thr Trp Val Gln Asn Cys Gln Gln Cys Arg Cys Leu Gln 715 720 725	2274
45	GGG GAA GTT GAT TGT TGG CCC CTG OCT TGC CCA GAT GTG GAG TGT GAA Gly Glu Val Asp Cys Trp Pro Leu Pro Cys Pro Asp Val Glu Cys Glu 730 735 740	2322
50	TTC AGC ATT CTC CCA GAG AAT GAG TGC TGC CCG OGC TGT GTC ACA GAC Phe Ser Ile Leu Pro Glu Asn Glu Cys Cys Pro Arg Cys Val Thr Asp 745 750 755	2370
55	OCT TGC CAG GCT GAC ACC ATC OGC AAT GAC ATC ACC AAG ACT TGC CTG Pro Cys Gln Ala Asp Thr Ile Arg Asn Asp Ile Thr Lys Thr Cys Leu 760 765 770	2418
60	GAC GAA ATG AAT GTG GTT CGC TTC AOC GGG TOC TCT TGG ATC AAA CAT Asp Glu Met Asn Val Val Arg Phe Thr Gly Ser Ser Trp Ile Lys His 775 780 785 790	2466
65	GGC ACT GAG TGT ACT CTC TGC CAG TGC AAG AAT GGC CAC ATC TGT TGC Gly Thr Glu Cys Thr Leu Cys Gln Cys Lys Asn Gly His Ile Cys Cys	2514

	795	800	805	
5	TCA GTG GAT CCA CAG TGC CTT CAG GAA CTG	TGAAGTTAAC	TGTCTCATGG	2564
	Ser Val Asp Pro Gln Cys Leu Gln Glu Leu			
	810	815		
	GAGATTTCTG	TTAAAAGAAT GTTCTTTTCAT	TAAAAGACCA AAAAGAAGTT AAAACTTAAA	2624
10	TTGGGTGATT	TGTGGGCAGC TAAATGCAGC	TTTGTTAATA GCTGAGTGAA CTTTCAATTA	2684
	TGAAATTTGT	GGAGCTTGAC AAAATCACAA AAGGAAAATT	ACTGGGGCAA AATTAGACCT	2744
	CAAGTCTGCC	TCTACTGTGT CTCACATCAC CATGTAGAAG	AATGGGGGTA CAGTATATAC	2804
15	CGTGACATCC	TGAACCTGG ATAGAAAGCC TGAGGCCATT	GGATCTGTGA AAGCCTCTAG	2864
	CTTCACTGGT	GCAGAAAATT TTCTCTAGA TCAGAATCTT	CAGAATCAGT TAGGTTCTCTC	2924
20	ACTGCAAGAA	ATAAAATGTC AGGCAGTGAA TGAATTATAT	TTTCAGAAGT AAAGCAAAGA	2984
	AGCTATAACA	TGTTATGTAC AGTACACTCT GAAAAGAAAT	CTGAAACAAG TTATTGTAAT	3044
	GATAAAAATA	ATGCACAGGC ATGGTTACTT AATATTTTCT	AACAGGAAAA GTCATCOCTA	3104
25	TTTCTTGTT	TTACTGCACT TAATATTATT TGGTTGAATT	TGTTCAGTAT AAGCTCGTTC	3164
	TTGTGCAAAA	TTAAATAAAT ATTTCTCTTA CCTT		3198

## (2) INFORMATION FOR SEQ ID NO:40:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 499 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:40:

Met	Glu	Leu	Ser	Glu	Pro	Val	Val	Glu	Asn	Gly	Glu	Val	Glu	Met	Ala
1				5				10					15		
Leu	Glu	Glu	Ser	Trp	Glu	His	Ser	Lys	Glu	Val	Ser	Glu	Ala	Glu	Pro
			20					25					30		
Gly	Gly	Gly	Ser	Ser	Gly	Asp	Ser	Gly	Pro	Pro	Glu	Glu	Ser	Gly	Gln
			35				40				45				

Glu Met Met Glu Glu Lys Glu Glu Ile Arg Lys Ser Lys Ser Val Ile  
 50 55 60  
 5 Val Pro Ser Gly Ala Pro Lys Lys Glu His Val Asn Val Val Phe Ile  
 65 70 75 80  
 Gly His Val Asp Ala Gly Lys Ser Thr Ile Gly Gly Gln Ile Met Phe  
 85 90 95  
 10 Leu Thr Gly Met Ala Asp Lys Arg Thr Leu Glu Lys Tyr Glu Arg Glu  
 100 105 110  
 15 Ala Glu Glu Lys Asn Arg Glu Thr Trp Tyr Leu Ser Trp Ala Leu Asp  
 115 120 125  
 Thr Asn Gln Glu Glu Arg Asp Lys Gly Lys Thr Val Glu Val Gly Arg  
 130 135 140  
 20 Ala Tyr Phe Glu Thr Glu Arg Lys His Phe Thr Ile Leu Asp Ala Pro  
 145 150 155 160  
 Gly His Lys Ser Phe Val Pro Asn Met Ile Gly Gly Ala Ser Gln Ala  
 165 170 175  
 25 Asp Leu Ala Val Leu Val Ile Ser Ala Arg Lys Gly Glu Phe Glu Thr  
 180 185 190  
 30 Gly Phe Glu Lys Gly Gly Gln Thr Arg Glu His Ala Met Phe Gly Lys  
 195 200 205  
 Thr Ala Gly Val Lys His Leu Ile Val Leu Ile Asn Lys Met Asp Asp  
 210 215 220  
 35 Pro Thr Val Asn Trp Gly Ile Glu Arg Tyr Glu Glu Cys Lys Glu Lys  
 225 230 235 240  
 Leu Val Pro Phe Leu Lys Lys Val Gly Phe Ser Pro Lys Lys Asp Ile  
 245 250 255  
 40 His Phe Met Pro Cys Ser Gly Leu Thr Gly Ala Asn Ile Lys Glu Gln  
 260 265 270  
 45 Ser Asp Phe Cys Pro Trp Tyr Thr Gly Leu Pro Phe Ile Pro Tyr Leu  
 275 280 285  
 Asn Asn Leu Pro Asn Phe Asn Arg Ser Ile Asp Gly Pro Ile Arg Leu  
 290 295 300  
 50 Pro Ile Val Asp Lys Tyr Lys Asp Met Gly Thr Val Val Leu Gly Lys  
 305 310 315 320  
 55

Leu Glu Ser Gly Ser Ile Phe Lys Gly Gln Gln Leu Val Met Met Pro  
 325 330 335  
 5 Asn Lys His Asn Val Glu Val Leu Gly Ile Leu Ser Asp Asp Thr Glu  
 340 345 350  
 Thr Asp Phe Val Ala Pro Gly Glu Asn Leu Lys Ile Arg Leu Lys Gly  
 355 360 365  
 10 Ile Glu Glu Glu Glu Ile Leu Pro Glu Phe Ile Leu Cys Asp Pro Ser  
 370 375 380  
 15 Asn Leu Cys His Ser Gly Arg Thr Phe Asp Val Gln Ile Val Ile Ile  
 385 390 395 400  
 Glu His Lys Ser Ile Ile Cys Pro Gly Tyr Asn Ala Val Leu His Ile  
 405 410 415  
 20 His Thr Cys Ile Glu Glu Val Glu Ile Thr Ala Leu Ile Ser Leu Val  
 420 425 430  
 Asp Lys Lys Ser Gly Glu Lys Ser Lys Thr Arg Pro Arg Phe Val Lys  
 435 440 445  
 25 Gln Asp Gln Val Cys Ile Ala Arg Leu Arg Thr Ala Gly Thr Ile Cys  
 450 455 460  
 Leu Glu Thr Phe Lys Asp Phe Pro Gln Met Gly Arg Phe Thr Leu Arg  
 465 470 475 480  
 30 Asp Glu Gly Lys Thr Ile Ala Ile Gly Lys Val Leu Lys Leu Val Pro  
 485 490 495  
 35 Glu Lys Asp

## (2) INFORMATION FOR SEQ ID NO:41:

40 (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 1497 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

45 (ii) MOLECULE TYPE: DNA(genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:41:

50 ATGGAAC TTT CAGAACTGT TGTAGAAAAT GGAGAGGTGG AAATGGOCCT AGAAGAATCA 60

55

TGGGAGCACA GTAAAGAAGT AAGTGAAGOC GAGCCTGGGG GTGGTTCCCTC GGGAGATTCA 120  
 5 GGGCCCCCAG AAGAAAGTGG CCAGGAAATG ATGGAGGAAA AAGAGGAAAT AAGAAAAATCC 180  
 AAATCTGTGA TOGTACCCCTC AGGTGCACCT AAGAAAGAAC ACGTAAATGT AGTATTCATT 240  
 GGCCATGTAG ACGCTGGCAA GTCAACCATC GGAGGACAGA TAATGTTTTT GACTGGAATG 300  
 10 GCTGACAAAA GAACACTGGA GAAATATGAA AGAGAAGCTG AGGAAAAAAA CAGAGAAAACC 360  
 TGGTATTTGT CCTGGGCTTT AGATACAAAT CAGGAGGAAC GAGACAAGGG TAAACAGTC 420  
 GAAGTGGGTC GTGCTATTTT TGAAACAGAA AGGAAACATT TCACAATTTT AGATGCCCCT 480  
 15 GGCCACAAGA GTTTTGTOCC AAATATGATT GGTGGTGCTT CTCAAGCTGA TTTGGCTGTG 540  
 CTGGTCATCT CTGOCAGGAA AGGAGAGTTT GAACTGGAT TTGAAAAAGG TGGACAGACA 600  
 20 AGAGAACATG CGATGTTTGG CAAAAOGGCA GGAGTAAAC ATTTAATAGT GCTTATTAAT 660  
 AAGATGGATG ATCCACAGT AAATTGGGGC ATCGAGAGAT ATGAAGAATG TAAAGAAAAA 720  
 CTGGTGCCCT TTTTGAAAAA AGTAGGCTTT AGTCAAAAAA AGGACATTCA CTTTATGCCC 780  
 25 TGCTCAGGAC TGACCGGAGC AAATATTAAA GAGCAGTCAG ATTTCTGCCC TTGGTACACT 840  
 GGATTACCAT TTATTOGTA TTTGAATAAC TTGCCAAACT TCAACAGATC AATTGATGGA 900  
 30 OCAATAAGAC TGCCAATTGT GGATAAGTAC AAAGATATGG GCACTGTGGT CCTGGGAAAG 960  
 CTGGAATCCG GGTCCATTTT TAAAGGCCAG CAGCTCGTGA TGATGCCAAA CAAGCACAAT 1020  
 GTAGAAGTTC TTGGAATACT TTCTGATGAT ACTGAACTG ATTTTGTAGC CCCAGGTGAA 1080  
 35 AAOCTCAAAA TCAGACTGAA GGAATTGAA GAAGAAGAGA TTCTTCAGA ATTCATACTT 1140  
 TGTGATCCTA GTAACCTCTG CCATTCTGGA CGCACGTTTG ATGTTTCAGAT AGTGATTATT 1200  
 GAGCACAAAT CCATCATCTG CCCAGGTTAT AATGCGGTGC TGCACATTCA TACTTGTATT 1260  
 40 GAGGAAGTTG AGATAACAGC GTTAATCTCC TTGGTAGACA AAAAATCAGG GGAAAAAAGT 1320  
 AAGACACGAC CCCGCTTGT GAAACAAGAT CAAGTATGCA TTGCTCGTTT AAGGACAGCA 1380  
 45 GGAACCATCT GCTCGAGAC GTTCAAAGAT TTCTCTCAGA TGGGTCGTTT TACTTTAAGA 1440  
 GATGAGGGTA AGACCATTC AATTGGAAAA GTTCTGAAAT TGGTCCAGA GAAGGAC 1497

50 (2) INFORMATION FOR SEQ ID NO:42:

55

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 2057 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA(genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vii) IMMEDIATE SOURCE:

(A) LIBRARY: Human fetal brain cDNA library

(B) CLONE: GEN-077A09

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 144..1640

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:42:

25	TCCCGGCGCG CTCCGGCAGC AACGATGAAG OCTGCAACGG CGCGGGATAC OCTCAAGGTA	60
	AAAGGATGGG ACGGGGGGCA OCTGTGGAAC CTTCOCGAGA GGAACCGTTA GTGTGCTTG	120
	AAGGTTCCAA TTCAGCGGTT ACC ATG GAA CTT TCA GAA OCT GTT GTA GAA	170
30	Met Glu Leu Ser Glu Pro Val Val Glu	
	1 5	
	AAT GGA GAG GTG GAA ATG GCC CTA GAA GAA TCA TGG GAG CAC AGT AAA	218
35	Asn Gly Glu Val Glu Met Ala Leu Glu Glu Ser Trp Glu His Ser Lys	
	10 15 20 25	
	GAA GTA AGT GAA GCC GAG OCT GGG GGT GGT TCC TCG GGA GAT TCA GGG	266
	Glu Val Ser Glu Ala Glu Pro Gly Gly Gly Ser Ser Gly Asp Ser Gly	
	30 35 40	
40	CCC OCA GAA GAA AGT GGC CAG GAA ATG ATG GAG GAA AAA GAG GAA ATA	314
	Pro Pro Glu Glu Ser Gly Gln Glu Met Met Glu Glu Lys Glu Glu Ile	
	45 50 55	
45	AGA AAA TCC AAA TCT GTG ATC GTA CCC TCA GGT GCA CCT AAG AAA GAA	362
	Arg Lys Ser Lys Ser Val Ile Val Pro Ser Gly Ala Pro Lys Lys Glu	
	60 65 70	
	CAC GTA AAT GTA GTA TTC ATT GGC CAT GTA GAC GCT GGC AAG TCA ACC	410
50	His Val Asn Val Val Phe Ile Gly His Val Asp Ala Gly Lys Ser Thr	
	75 80 85	

5	ATC GGA GGA CAG ATA ATG TTT TTG ACT GGA ATG GCT GAC AAA AGA ACA Ile Gly Gly Gln Ile Met Phe Leu Thr Gly Met Ala Asp Lys Arg Thr 90 95 100 105	458
10	CTG GAG AAA TAT GAA AGA GAA GCT GAG GAA AAA AAC AGA GAA AOC TGG Leu Glu Lys Tyr Glu Arg Glu Ala Glu Glu Lys Asn Arg Glu Thr Trp 110 115 120	506
15	TAT TTG TOC TGG GOC TTA GAT ACA AAT CAG GAG GAA CGA GAC AAG GGT Tyr Leu Ser Trp Ala Leu Asp Thr Asn Gln Glu Glu Arg Asp Lys Gly 125 130 135	554
20	AAA ACA GTC GAA GTG GGT CGT GOC TAT TTT GAA ACA GAA AGG AAA CAT Lys Thr Val Glu Val Gly Arg Ala Tyr Phe Glu Thr Glu Arg Lys His 140 145 150	602
25	TTC ACA ATT TTA GAT GOC OCT GGC CAC AAG AGT TTT GTC CCA AAT ATG Phe Thr Ile Leu Asp Ala Pro Gly His Lys Ser Phe Val Pro Asn Met 155 160 165	650
30	ATT GGT GGT GCT TCT CAA GCT GAT TTG GCT GTG CTG GTC ATC TCT GCC Ile Gly Gly Ala Ser Gln Ala Asp Leu Ala Val Leu Val Ile Ser Ala 170 175 180 185	698
35	AGG AAA GGA GAG TTT GAA ACT GGA TTT GAA AAA GGT GGA CAG ACA AGA Arg Lys Gly Glu Phe Glu Thr Gly Phe Glu Lys Gly Gly Gln Thr Arg 190 195 200	746
40	GAA CAT GCG ATG TTT GGC AAA ACG GCA GGA GTA AAA CAT TTA ATA GTG Glu His Ala Met Phe Gly Lys Thr Ala Gly Val Lys His Leu Ile Val 205 210 215	794
45	CTT ATT AAT AAG ATG GAT GAT CCC ACA GTA AAT TGG GGC ATC GAG AGA Leu Ile Asn Lys Met Asp Asp Pro Thr Val Asn Trp Gly Ile Glu Arg 220 225 230	842
50	TAT GAA GAA TGT AAA GAA AAA CTG GTG CCC TTT TTG AAA AAA GTA GGC Tyr Glu Glu Cys Lys Glu Lys Leu Val Pro Phe Leu Lys Lys Val Gly 235 240 245	890
55	TTT AGT CCA AAA AAG GAC ATT CAC TTT ATG CCC TGC TCA GGA CTG AOC Phe Ser Pro Lys Lys Asp Ile His Phe Met Pro Cys Ser Gly Leu Thr 250 255 260 265	938
60	GGA GCA AAT ATT AAA GAG CAG TCA GAT TTC TGC OCT TGG TAC ACT GGA Gly Ala Asn Ile Lys Glu Gln Ser Asp Phe Cys Pro Trp Tyr Thr Gly 270 275 280	986
65	TTA CCA TTT ATT CCG TAT TTG AAT AAC TTG CCA AAC TTC AAC AGA TCA Leu Pro Phe Ile Pro Tyr Leu Asn Asn Leu Pro Asn Phe Asn Arg Ser	1034

	285	290	295	
5	ATT GAT GGA OCA ATA AGA CTG OCA ATT GTG GAT AAG TAC AAA GAT ATG Ile Asp Gly Pro Ile Arg Leu Pro Ile Val Asp Lys Tyr Lys Asp Met 300 305 310			1082
10	GGC ACT GTG GTC CTG GGA AAG CTG GAA TOC GGG TOC ATT TTT AAA GGC Gly Thr Val Val Leu Gly Lys Leu Glu Ser Gly Ser Ile Phe Lys Gly 315 320 325			1130
15	CAG CAG CTC GTG ATG ATG OCA AAC AAG CAC AAT GTA GAA GTT CTT GGA Gln Gln Leu Val Met Met Pro Asn Lys His Asn Val Glu Val Leu Gly 330 335 340 345			1178
20	ATA CTT TCT GAT GAT ACT GAA ACT GAT TTT GTA GGC OCA GGT GAA AAC Ile Leu Ser Asp Asp Thr Glu Thr Asp Phe Val Ala Pro Gly Glu Asn 350 355 360			1226
25	CTC AAA ATC AGA CTG AAG GGA ATT GAA GAA GAA GAG ATT CTT OCA GAA Leu Lys Ile Arg Leu Lys Gly Ile Glu Glu Glu Glu Ile Leu Pro Glu 365 370 375			1274
30	TTC ATA CTT TGT GAT OCT AGT AAC CTC TGC CAT TCT GGA CGC ACG TTT Phe Ile Leu Cys Asp Pro Ser Asn Leu Cys His Ser Gly Arg Thr Phe 380 385 390			1322
35	GAT GTT CAG ATA GTG ATT ATT GAG CAC AAA TOC ATC ATC TGC OCA GGT Asp Val Gln Ile Val Ile Ile Glu His Lys Ser Ile Ile Cys Pro Gly 395 400 405			1370
40	TAT AAT GCG GTG CTG CAC ATT CAT ACT TGT ATT GAG GAA GTT GAG ATA Tyr Asn Ala Val Leu His Ile His Thr Cys Ile Glu Glu Val Glu Ile 410 415 420 425			1418
45	ACA GCG TTA ATC TOC TTG GTA GAC AAA AAA TCA GGG GAA AAA AGT AAG Thr Ala Leu Ile Ser Leu Val Asp Lys Lys Ser Gly Glu Lys Ser Lys 430 435 440			1466
50	ACA OGA OCC CGC TTC GTG AAA CAA GAT CAA GTA TGC ATT GCT CGT TTA Thr Arg Pro Arg Phe Val Lys Gln Asp Gln Val Cys Ile Ala Arg Leu 445 450 455			1514
55	AGG ACA GCA GGA ACC ATC TGC CTC GAG ACG TTC AAA GAT TTT OCT CAG Arg Thr Ala Gly Thr Ile Cys Leu Glu Thr Phe Lys Asp Phe Pro Gln 460 465 470			1562
	ATG GGT CGT TTT ACT TTA AGA GAT GAG GGT AAG ACC ATT GCA ATT GGA Met Gly Arg Phe Thr Leu Arg Asp Glu Gly Lys Thr Ile Ala Ile Gly 475 480 485			1610



AAA GTT CTG AAA TTG GTC CCA GAG AAG GAC TAAGCAATTT TCTTGATGCC 1660  
 Lys Val Leu Lys Leu Val Pro Glu Lys Asp  
 490 495

TCTGCAAGAT ACTGTGAGGA GAATTGACAG CAAAAGTTCA CCACCTACTC TTATTTACTG 1720

OCCATTGATT GACTTTTCTT CATATTTTGC AAAGAGAAAT TTCACAGCAA AAATTCATGT 1780

TTTGTCAGCT TTCTCATGTT GAGATCTGTT ATGTCACCTGA TGAATTTACC CTCAAGTTTC 1840

CTTCTCTGT ACCACTCTGC TTCTTGGAC AATATCAGTA ATAGCTTTGT AAGTGATGTG 1900

GACGTAATTG OCTACAGTAA TAAAAAATA ATGTACTTTA ATTTTTCATT TTCTTTTAGG 1960

ATATTTAGAC CACCTTGTGTT CCACGCAAAC CAGAGTGTGT CAGTGTTTGT GTGTGTGTTA 2020

AAATGATAAC TAACATGTGA ATAAAATACT CCATTTG 2057

## Claims

1. A GDP dissociation stimulating protein gene comprises a nucleotide sequence coding for the amino acid sequence shown under SEQ ID NO:1.
2. A GDP dissociation stimulating protein gene comprises the nucleotide sequence shown under SEQ ID NO:2.
3. A GDP dissociation stimulating protein gene as defined in Claim 2 which has the nucleotide sequence shown under SEQ ID NO:3.
4. A brain-specific nucleosome assembly protein gene comprises a nucleotide sequence coding for the amino acid sequence shown under SEQ ID NO:19.
5. A brain-specific nucleosome assembly protein gene comprises a nucleotide sequence shown under SEQ ID NO:20.
6. A brain-specific nucleosome assembly protein gene as defined in Claim 5 which has the nucleotide sequence shown under SEQ ID NO:21.
7. A human skeletal muscle-specific ubiquitin-conjugating enzyme gene comprises a nucleotide sequence coding for the amino acid sequence shown under SEQ ID NO:22.
8. A human skeletal muscle-specific ubiquitin-conjugating enzyme gene comprises the nucleotide sequence shown under SEQ ID NO:23.
9. A human skeletal muscle-specific ubiquitin-conjugating enzyme gene as defined in Claim 8 which has the nucleotide sequence shown under SEQ ID NO:24.
10. A TMP-2 gene comprises a nucleotide sequence coding for the amino acid sequence shown under SEQ ID NO:25.
11. A TMP-2 gene comprises the nucleotide sequence shown under SEQ ID NO:26.
12. A TMP-2 gene as defined in Claim 11 which has the nucleotide sequence shown under SEQ ID NO:27.
13. A human NPIK gene comprises a nucleotide sequence coding for the amino acid sequence shown under SEQ ID NO:28.

14. A human NPIK gene comprises the nucleotide sequence shown under SEQ ID NO:29.
15. A human NPIK gene as defined in Claim 14 which has the nucleotide sequence shown under SEQ ID NO:30.
- 5 16. A human NPIK gene comprises a nucleotide sequence coding for the amino acid sequence shown under SEQ ID NO:31.
17. A human NPIK gene comprises the nucleotide sequence shown under SEQ ID NO:32.
- 10 18. A human NPIK gene as defined in Claim 17 which has the nucleotide sequence shown under SEQ ID NO:33.
19. A nel-related protein type 1 gene comprises a nucleotide sequence coding for the amino acid sequence shown under SEQ ID NO:34.
- 15 20. A nel-related protein type 1 gene comprises the nucleotide sequence shown under SEQ ID NO:35.
21. A nel-related protein type 1 gene as defined in Claim 20 which has the nucleotide sequence shown under SEQ ID NO:36.
- 20 22. A nel-related protein type 2 gene comprises a nucleotide sequence coding for the amino acid sequence shown under SEQ ID NO:37.
23. A nel-related protein type 2 gene comprises the nucleotide sequence shown under SEQ ID NO:38.
- 25 24. A nel-related protein type 2 gene as defined in Claim 23 which has the nucleotide sequence shown under SEQ ID NO:39.
25. A method for the in vitro diagnosis of hereditary diseases and cancer, characterized by employing any of the nucleotide or amino acid sequences as given in claims 1-24.
- 30 26. The use of any of the nucleotide or amino acid sequences as given in claims 1 - 24 for in vitro diagnosis as well as for the preparation of a pharmaceutical for the treatment of diseases.

35

40

45

50

55

FIG. 1






FIG. 2





Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

**EP 0 796 913 A3**

(12)

**EUROPEAN PATENT APPLICATION**

(88) Date of publication A3:  
31.01.2001 Bulletin 2001/05

(43) Date of publication A2:  
24.09.1997 Bulletin 1997/39

(21) Application number: 97104842.6

(22) Date of filing: 19.03.1997

(51) Int. Cl.<sup>7</sup>: **C12N 15/12**, C12N 15/54,  
C12N 15/55, C07K 14/47,  
C12N 9/12, C12N 9/00,  
C12N 9/64, C12Q 1/68,  
A61K 38/17, A61K 38/45,  
A61K 38/53

(84) Designated Contracting States:  
**AT BE CH DE DK ES FI FR GB GR IE IT LI LU MC  
NL PT SE**

(30) Priority: 19.03.1996 JP 6341096  
05.03.1997 JP 6916397

(71) Applicant:  
**OTSUKA PHARMACEUTICAL CO., LTD.**  
Chiyoda-ku Tokyo 101 (JP)

(72) Inventors:  
• Fujiwara, Tsutomu  
Naruto-shi, Tokushima-ken (JP)

• Watanabe, Takeshi  
Aizumi-cho, Itano-gun, Tokushima-ken (JP)  
• Horie, Masato  
Tokushima-shi, Tokushima-ken (JP)

(74) Representative:  
**Hansen, Bernd, Dr. Dipl.-Chem. et al**  
**Hoffmann Eitle,**  
**Patent- und Rechtsanwälte,**  
**Arabellastrasse 4**  
**81925 München (DE)**

(54) **GDP dissociation stimulating protein, brain-specific nucleosome assembly protein, skeletal muscle specific ubiquitin-conjugating enzyme, cell proliferation protein, phosphatidylinositolkinase, nel related proteins**

(57) The present invention provides human genes, for example human genes comprising nucleotide sequences coding for amino acid sequences of GDP dissociation stimulating protein, brain-specific nucleosome assembly protein, skeletal muscle specific ubiquitin-conjugating enzyme, cell proliferation protein, phosphatidylinositolkinase, nel related proteins. Analysis of diseases associated with the genes, for example, hereditary diseases and cancer, and diagnosis and treatment of such diseases.

**EP 0 796 913 A3**



European Patent  
Office

# EUROPEAN SEARCH REPORT

Application Number  
EP 97 10 4842

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
Y	SPAARGAREN MARCEL ET AL: "Identification of the guanine nucleotide dissociation stimulator for Ral as a putative effector molecule of R-ras, H-ras, K-ras, and rap." PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE UNITED STATES, vol. 91, no. 26, 1994, pages 12609-12613, XP002145498 1994 ISSN: 0027-8424 * the whole document *	1-3,25, 26	C12N15/12 C12N15/54 C12N15/55 C07K14/47 C12N9/12 C12N9/00 C12N9/64 C12Q1/68 A61K38/17 A61K38/45
Y	HILLIER L. ET AL.: "yd85f10.r1 cDNA clone - similar to mouse guanine nucleotide dissociation stimulator" EMBL DATABASE, ACCESSION NUMBER HS60650, 1 April 1995 (1995-04-01), XP002145499 * the whole document *	1-3,25, 26	
Y	HILLIER ET AL.: "yx19d04.r1 cDNA clone" EMBL DATABASE, ACCESSION NUMBER HS045260, 30 December 1995 (1995-12-30), XP002145500 * the whole document *	1-3,25, 26	TECHNICAL FIELDS SEARCHED (Int.Cl.6) C07K C12N
P,X	ISOMURA M ET AL: "Isolation and mapping of RAB2L, a human cDNA that encodes a protein homologous to RaIGDS." CYTOGENETICS AND CELL GENETICS, vol. 74, no. 4, 1996, pages 263-265, XP000938401 ISSN: 0301-0171 * the whole document *	1-3	
-/--			
-The present search report has been drawn up for all claims			
Place of search <b>THE HAGUE</b>		Date of completion of the search <b>23 August 2000</b>	Examiner <b>Gurdjian, D</b>
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	

EPO FORM 1503 03.82 (P04C01)



European Patent  
Office

# EUROPEAN SEARCH REPORT

Application Number  
EP 97 10 4842

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
P,X	<p>PETERSON SCOTT N ET AL: "Identification of a novel RalGDS-related protein as a candidate effector for Ras and Rap1." JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 271, no. 47, 1996, pages 29903-29908, XP002145501.</p> <p>ISSN: 0021-9258</p> <p>* the whole document *</p> <p>-----</p>	1-3	
			TECHNICAL FIELDS SEARCHED (Int.Cl.6)
<p><del>The present search report has been drawn up for all claims</del></p>			
Place of search		Date of completion of the search	Examiner
THE HAGUE		23 August 2000	Gurdjian, D
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone</p> <p>Y : particularly relevant if combined with another document of the same category</p> <p>A : technological background</p> <p>O : non-written disclosure</p> <p>P : intermediate document</p> <p>T : theory or principle underlying the invention</p> <p>E : earlier patent document, but published on, or after the filing date</p> <p>D : document cited in the application</p> <p>L : document cited for other reasons</p> <p>&amp; : member of the same patent family, corresponding document</p>			

EPO FORM 1503 03/82 (P04C01)



European Patent  
Office

Application Number

EP 97 10 4842

### CLAIMS INCURRING FEES

The present European patent application comprised at the time of filing more than ten claims.

☐ Only part of the claims have been paid within the prescribed time limit. The present European search report has been drawn up for the first ten claims and for those claims for which claims fees have been paid, namely claim(s):

☐ No claims fees have been paid within the prescribed time limit. The present European search report has been drawn up for the first ten claims.

### LACK OF UNITY OF INVENTION

The Search Division considers that the present European patent application does not comply with the requirements of unity of invention and relates to several inventions or groups of inventions, namely:

see sheet B

☐ All further search fees have been paid within the fixed time limit. The present European search report has been drawn up for all claims.

☐ As all searchable claims could be searched without effort justifying an additional fee, the Search Division did not invite payment of any additional fee.

☐ Only part of the further search fees have been paid within the fixed time limit. The present European search report has been drawn up for those parts of the European patent application which relate to the inventions in respect of which search fees have been paid, namely claims:

☒ None of the further search fees have been paid within the fixed time limit. The present European search report has been drawn up for those parts of the European patent application which relate to the invention first mentioned in the claims, namely claims:

25, 26 partially and 1-3



European Patent  
Office

**LACK OF UNITY OF INVENTION  
SHEET B**

Application Number

EP 97 10 4842

The Search Division considers that the present European patent application does not comply with the requirements of unity of invention and relates to several inventions or groups of inventions, namely:

**1. Claims: 25,26 partially and 1-3**

GDP-dissociation-stimulating protein gene , and  
corresponding method for in vitro diagnosing and use  
for the preparation of a pharmaceutical

**2. Claims: 25,26 partially and 4-6**

brain-specific nucleosome assembly protein gene ,  
and corresponding method for in vitro diagnosing and use  
for the preparation of a pharmaceutical

**3. Claims: 25,26 partially and 7-9**

human skeletal-muscle-specific ubiquitin-conjugating enzyme  
gene ,and corresponding method for in vitro diagnosing and  
use for the preparation of a pharmaceutical

**4. Claims: 25,26 partially and 10-12**

TMP-2 cell proliferation gene and corresponding method for  
in vitro diagnosing and use for the preparation of a  
pharmaceutical

**5. Claims: 25,26 partially and 13-18**

human NPIK phosphatidylinositolkinase genes and  
corresponding method for in vitro diagnosing and use for the  
preparation of a pharmaceutical

**6. Claims: 25,26 partially and 19-24**

nel-related protein genes and corresponding method for in  
vitro diagnosing and use for the preparation of a  
pharmaceutical